

indicate the risk of its occurrence in these regions, where *H. longicornis* is widely distributed (10). More extensive investigation to clarify the natural circulation of NSDV among ticks should be conducted and surveillance of sheep improved to prevent outbreaks of Nairobi sheep disease in China and East Asia.

This work was supported by the Science and Technology Basic Work Program from the Ministry of Science and Technology of China (2013FY113600), and the Military Medical Innovation Program of Academy of Military Medical Sciences (2012CXJJ019).

## References

1. Montgomery E. On a tick-borne gastro-enteritis of sheep and goats occurring in East Africa. *J Comp Pathol Ther.* 1917;30:28–57. [http://dx.doi.org/10.1016/S0368-1742\(17\)80002-3](http://dx.doi.org/10.1016/S0368-1742(17)80002-3)
2. Marczinke BI, Nichol ST. Nairobi sheep disease virus, an important tick-borne pathogen of sheep and goats in Africa, is also present in Asia. *Virology.* 2002;303:146–51. <http://dx.doi.org/10.1006/viro.2002.1514>
3. Weinbren MP, Gourlay RN, Lumsden WHR, Weinbren WM. An epizootic of Nairobi sheep disease in Uganda. *J Comp Pathol Ther.* 1958;68:174–87. [http://dx.doi.org/10.1016/S0368-1742\(58\)80018-1](http://dx.doi.org/10.1016/S0368-1742(58)80018-1)
4. Lasecka L, Baron MD. The nairovirus Nairobi sheep disease virus/Ganjam virus induces the translocation of protein disulphide isomerase-like oxidoreductases from the endoplasmic reticulum to the cell surface and the extracellular space. *PLoS ONE.* 2014;9:e94656. <http://dx.doi.org/10.1371/journal.pone.0094656>
5. Yadav PD, Vincent MJ, Khristova M, Kale C, Nichol ST, Mishra AC, et al. Genomic analysis reveals Nairobi sheep disease virus to be highly diverse and present in both Africa, and in India in the form of the Ganjam virus variant. *Infect Genet Evol.* 2011;11:1111–20. <http://dx.doi.org/10.1016/j.meegid.2011.04.001>
6. Perera LP, Peiris JSM, Weilgama DJ, Calisher CH, Shope RE. Nairobi sheep disease virus isolated from *Haemaphysalis intermedia* ticks collected in Sri Lanka. *Ann Trop Med Parasitol.* 1996;90:91–3.
7. Liu Q, He B, Huang SY, Wei F, Zhu XQ. Severe fever with thrombocytopenia syndrome, an emerging tick-borne zoonosis. *Lancet Infect Dis.* 2014;14:763–72. [http://dx.doi.org/10.1016/S1473-3099\(14\)70718-2](http://dx.doi.org/10.1016/S1473-3099(14)70718-2)
8. Rajagopalan PK, Sreenivasan MA, Paul SD. Isolation of Ganjam virus from the bird tick *Haemaphysalis wellingtoni* Nuttall and Warburton 1907. *Indian J Med Res.* 1970;58:1195–6.
9. Joshi MV, Geevarghese G, Joshi GD, Ghodke YS, Mourya DT, Mishra AC. Isolation of Ganjam virus from ticks collected of domestic animals around Pune, Maharashtra, India. *J Med Entomol.* 2005;42:204–6. <http://dx.doi.org/10.1093/jmedent/42.2.204>
10. Hoogstraal H, Roberts FH, Kohls GM, Tipton VJ. Review of *Haemaphysalis* (Kaiseriana) *longicornis* Neumann (resurrected) of Australia, New Zealand, New Caledonia, Fiji, Japan, Korea, and Northeastern China and USSR, and its parthenogenetic and bisexual populations (Ixodoidea, Ixodidae). *J Parasitol.* 1968;54:1197–213. <http://dx.doi.org/10.2307/3276992>

Address for correspondence: Quan Liu, Key Laboratory of Jilin Province for Zoonosis Prevention and Control, Military Veterinary Institute, Academy of Military Medical Sciences, 666 Liuying West Rd, Jingyue Economic Development Zone, Changchun, 130122, People's Republic of China; email: liuquan1973@hotmail.com

## Avian Influenza A(H10N7) Virus–Associated Mass Deaths among Harbor Seals

Rogier Bodewes, Theo M. Bestebroer, Erhard van der Vries, Josanne H. Verhagen, Sander Herfst, Marion P. Koopmans, Ron A.M. Fouchier, Vanessa M. Pfankuche, Peter Wohlsein, Ursula Siebert, Wolfgang Baumgärtner, Albert D.M.E. Osterhaus

Author affiliations: Erasmus Medical Centre, Rotterdam, the Netherlands (R. Bodewes, T.M. Bestebroer, E. van der Vries, J.H. Verhagen, S. Herfst, M.P. Koopmans, R.A.M. Fouchier, A.D.M.E. Osterhaus); University of Veterinary Medicine, Hannover, Germany (V.M. Pfankuche, P. Wohlsein, U. Siebert, W. Baumgärtner, A.D.M.E. Osterhaus); Artemis One Health, Utrecht, the Netherlands (A.D.M.E. Osterhaus)

DOI: <http://dx.doi.org/10.3201/eid2104.141675>


**To the Editor:** Avian influenza A viruses occasionally cross the species barrier; influenza A(H5N1) virus and the recently emerged influenza A(H7N9) virus are prime examples of bird-to-human transmission (1,2). In addition, avian influenza A viruses can cross to various other mammalian species, including pinnipeds (e.g., seals) (3,4).

Recently, mass deaths have occurred among harbor seals (*Phoca vitulina*); hundreds of carcasses washed up the shores of Sweden (March 2014), Denmark (July 2014), and Germany (October 2014). Approximately 1,400 dead harbor seals were seen in the coastal waters of Schleswig-Holstein in Germany alone, where the population is ≈12,000 animals.

We report the investigation of the deaths of 17 seals from different age groups that were found dead on the islands of Helgoland and Sylt, Germany, during the second week of October 2014. Complete necropsies were performed on the carcasses, which were in variable nutritional conditions, ranging from very poor to good. Necropsy results showed consistently poorly retracted lungs with severe congestion, occasional diffuse consolidation, and multifocal firm nodular areas of gray-yellow discoloration with varying numbers of metazoic parasites. Histologic examinations showed acute necrotizing bronchitis and adenitis of bronchial glands with sloughing of epithelial cells (Figure, panel A). Occasionally, mild interstitial pneumonia was found. Multifocal pyogranulomatous to necrotizing pneumonia was associated with parasite infestation. A few animals had suppurative to necrotizing or nonsuppurative rhinitis and tracheitis.

Because mass deaths among seals were caused by phocine distemper virus in the same area in 1988 and 2002,

we tested lung and throat swab samples for morbillivirus using reverse transcription PCR (RT-PCR) and immunohistochemical analysis (5). In addition, real-time RT-PCR targeting the influenza A virus matrix gene was performed (6). No indications for the morbillivirus were detected by



Histopathologic and phylogenetic analyses of necropsy samples from harbor seals infected with avian influenza A(H10N7) virus, Germany, 2014. A) Lung of harbor seal showing marked necrosis and sloughing of epithelial cells in bronchial glands (arrows); c = bronchial cartilage; hematoxylin and eosin stain. Scale bar indicates 50  $\mu$ m B) Immunohistochemical labeling of influenza A nucleoprotein in bronchial epithelial cells (arrowheads) and glandular epithelial cells (arrows); c = bronchial cartilage; avidin-biotin-peroxidase complex method. Scale bar indicates 50  $\mu$ m. C) Maximum-likelihood phylogenetic tree of the partial hemagglutinin gene (1,577 nt) of the influenza A/harbor seal/Germany/1/2014 (H10N7) isolate and various other closely related viruses. GenBank accession numbers are provided in online Technical Appendix Table 1 (<http://wwwnc.cdc.gov/EID/article/21/4/14-1675-Techapp1.pdf>). Scale bar indicates nucleotide substitutions per site.

RT-PCR and immunohistochemistry; however, in lung lesions and throat swab samples of 11 animals, a positive signal was observed by the influenza A matrix gene real-time RT-PCR (cycle threshold values 15.0–33.9). Influenza A virus (A/harbor seal/Germany/1/2014) was subsequently isolated from lung and throat swab samples; the virus replicated to high titers in 11-day-old embryonated chicken eggs and on MDCK cells. By PCR using specific primers and subsequent Sanger sequencing of the hemagglutinin and neuraminidase genes, this virus was characterized as an influenza A virus of the H10N7 subtype, commonly found in migratory waterfowl (6). In addition, genetic analyses of all other gene segments indicated that the influenza virus A/harbor seal/Germany/1/2014 was most closely related to various influenza A viruses detected in wild birds. Specifically, the hemagglutinin and neuraminidase genes were genetically most closely related to subtype H10N7 viruses recently found in migratory ducks in Georgia, Egypt, and the Netherlands (Figure, panel C) (7). Genetic analyses were based on BLAST analyses using public databases available as of October 17, 2014 (<http://www.ncbi.nlm.nih.gov>, <http://www.gisaid.com>) and supplemented with H10 and N7 sequences from the international wild bird surveillance program of Erasmus Medical Center (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/21/4/14-1675-Techapp1.pdf>). A maximum-likelihood phylogenetic tree of the hemagglutinin gene was generated by using PhyML version 3.1 (8) with the general time reversible +I+G model of nucleotide substitution; a full heuristic search and subtree pruning and regrafting searches were performed. The tree was visualized by using Figtree version 1.4.0 (<http://tree.bio.ed.ac.uk/software/figtree>).

To further elucidate the role of influenza A(H10N7) virus in the pathogenesis of the disease causing deaths among the seals, we conducted immunohistochemical analysis on the lungs using an influenza A virus nucleoprotein-specific monoclonal antibody (9). Evaluation of the lung tissues of the dead seals showed viral antigen in cytoplasm and nuclei of epithelial cells of bronchi and bronchial glands of affected lung areas (Figure, panel B), which suggests that this virus played a major role in the deaths. Immunohistochemical analysis performed on various organs (including brain and olfactory bulb) indicated that viral antigen was restricted to the respiratory tract.

Although avian influenza A virus infections previously have caused mass deaths in seals (3,4,10), subtype H10N7 has not been associated with such events. We can speculate that the ongoing deaths could eventually affect all harbor seal populations of northwestern Europe and have consequences for wildlife management and seal rehabilitation activities. In addition, preliminary analysis of the hemagglutinin sequence of the influenza A(H10N7) virus suggests the presence of molecular determinants that indicate

mammalian adaptation. Various analyses are ongoing to answer questions about the route of transmission among seals and possible transmissibility to humans.

Note added in proof: Zohari et al. also recently reported the involvement of avian influenza A(H10N7) virus in mass deaths of harbor seals in Sweden (Euro Surveill. 2014;19:pii: 20967).

This work was supported in part by a grant from the Niedersachsen-Research Network on Neuroinfectiology of the Ministry of Science and Culture of Lower Saxony, Germany, and the European Research Council project FLUPLAN (250136).

## References

1. Claas EC, Osterhaus AD, van Beek R, De Jong JC, Rimmelzwaan GF, Senne DA, et al. Human influenza A H5N1 virus related to a highly pathogenic avian influenza virus. *Lancet*. 1998;351:472–7. [http://dx.doi.org/10.1016/S0140-6736\(97\)11212-0](http://dx.doi.org/10.1016/S0140-6736(97)11212-0)
2. Gao R, Cao B, Hu Y, Feng Z, Wang D, Hu W, et al. Human infection with a novel avian-origin influenza A (H7N9) virus. *N Engl J Med*. 2013;368:1888–97. <http://dx.doi.org/10.1056/NEJMoa1304459>
3. Reperant LA, Rimmelzwaan GF, Kuiken T. Avian influenza viruses in mammals. *Rev Sci Tech*. 2009;28:137–59.
4. Anthony SJ, St Leger JA, Puglianes K, Ip HS, Chan JM, Carpenter ZW, et al. Emergence of fatal avian influenza in New England harbor seals. *MBio*. 2012;3:e00166-12.
5. Jensen T, van de Bildt M, Dietz HH, Andersen TH, Hammer AS, Kuiken T, et al. Another phocine distemper outbreak in Europe. *Science*. 2002;297:209. <http://dx.doi.org/10.1126/science.1075343>
6. Munster VJ, Baas C, Lexmond P, Waldenstrom J, Wallensten A, Fransson T, et al. Spatial, temporal, and species variation in prevalence of influenza A viruses in wild migratory birds. *PLoS Pathog*. 2007;3:e61. <http://dx.doi.org/10.1371/journal.ppat.0030061>
7. Lewis NS, Javakhishvili Z, Russell CA, Machabishvili A, Lexmond P, Verhagen JH, et al. Avian influenza virus surveillance in wild birds in Georgia: 2009–2011. *PLoS ONE*. 2013;8:e58534.
8. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol*. 2010;59:307–21. <http://dx.doi.org/10.1093/sysbio/syq010>
9. de Boer GF, Back W, Osterhaus AD. An ELISA for detection of antibodies against influenza A nucleoprotein in humans and various animal species. *Arch Virol*. 1990;115:47–61. <http://dx.doi.org/10.1007/BF01310622>
10. Hinshaw VS, Bean WJ, Webster RG, Rehg JE, Fiorelli P, Early G, et al. Are seals frequently infected with avian influenza viruses? *J Virol*. 1984;51:863–5.

Address for correspondence: Albert D.M.E. Osterhaus, University of Veterinary Medicine—Department of Pathology, Buenteweg 17, Lower Saxony, Hannover D-30559, Germany; email: [albert.osterhaus@tiho-hannover.de](mailto:albert.osterhaus@tiho-hannover.de)

## Zika Virus Infection, Philippines, 2012

**Maria Theresa Alera, Laura Hermann, Ilya A. Tac-An, Chonticha Klungthong, Wiriya Rutvisuttinunt, Wudtichai Manasatienkij, Daisy Villa, Butsaya Thaisomboonsuk, John Mark Velasco, Piyawan Chinnawirotpisan, Catherine B. Lago, Vito G. Roque, Jr., Louis R. Macareo, Anon Srikiatkachorn, Stefan Fernandez, In-Kyu Yoon**

Author affiliations: Philippines-AFRIMS (Armed Forces Research Institute of Medical Sciences), Cebu City, Philippines (M.T. Alera, J.M. Velasco, C.B. Lago); University of Toronto, Toronto, Ontario, Canada (L. Hermann); AFRIMS, Bangkok, Thailand (L. Hermann, C. Klungthong, W. Rutvisuttinunt, W. Manasatienkij, B. Thaisomboonsuk, P. Chinnawirotpisan, L.R. Macareo, S. Fernandez, I.-K. Yoon); Cebu City Health Department, Cebu City (I.A. Tac-An, D. Villa); Department of Health, Manila, Philippines (V.G. Roque, Jr.); University of Massachusetts Medical School, Worcester, Massachusetts, USA (A. Srikiatkachorn)

DOI: <http://dx.doi.org/10.3201/eid2104.141707>

**To the Editor:** Zika virus (ZIKV), a mosquito-borne flavivirus, was first isolated from a rhesus monkey in Uganda in 1947 (1). This positive-sense, single-stranded RNA virus (family *Flaviviridae*, genus *Flavivirus*) has a 10,794-nt genome and is most closely related to Spondweni virus (2,3). Phylogenetic analyses have revealed 2 major lineages: Asian and African (2–4).

The first human infection with ZIKV was reported in Nigeria in 1954 (5). The virus caused only sporadic infections until 2007, when a large outbreak occurred on Yap, an island in the Federated States of Micronesia (6). In October 2013, ZIKV was detected in French Polynesia; since then, >400 laboratory-confirmed cases have been reported (7). ZIKV has spread across the South Pacific, and autochthonous cases have been reported in New Caledonia, Easter Island, and the Cook Islands. Several cases of ZIKV infections have been reported in travelers to Southeast Asia (4,8) and French Polynesia (3,7).

In March 2012, a prospective longitudinal cohort study, which included active surveillance for acute febrile illness, was initiated in Cebu City, Philippines (I. Yoon, unpub. data). Participants contacted study staff to report fever and were also contacted weekly by staff to determine if they had fever during the previous 7 days. Fever episodes triggered an acute-illness visit by a study nurse, who performed a clinical assessment of the patient and collected an acute-phase blood sample. During the first year of surveillance, 270 acute febrile

# Avian Influenza A(H10N7) Virus–Associated Mass Deaths among Harbor Seals

## Technical Appendix

Technical Appendix Table. Details of hemagglutinin sequences shown in the Figure\*

Isolate name	Accession no.	Online database
A/seal/Germany/EMC-1/2014(H10N7)	EPI544351	GISAID EpiFlu
A/mallard/Netherlands/1/2014(H10N7)	EPI552751	GISAID EpiFlu
A/mallard/Sweden/133546/2011(H10N4)	CY183991.1	GenBank
A/domestic duck/RepublicofGeorgia/2/2010(H10N7)	CY185457.1	GenBank
A/domestic duck/RepublicofGeorgia/1/2010(H10(N7)	CY185449.1	GenBank
A/mallard/Sweden/104746/2009(H10N1)	CY183855.1	GenBank
A/mallard/Sweden/105465/2009(H10N1)	CY183927.1	GenBank
A/mallard/Sweden/102260/2009(H10N1)	CY183847.1	GenBank
A/mallard/Sweden/105323/2009(H10N1)	CY183887.1	GenBank
A/mallard/Sweden/105402/2009(H10N1)	CY183911.1	GenBank
A/mallard/Sweden/105522/2009(H10N1)	JX566079.1	GenBank
A/mallard/Netherlands/50/2010(H10N7)	EPI552752	GISAID EpiFlu
A/mallard/Netherlands/47/2010(H10N7)	EPI552753	GISAID EpiFlu
A/mallard/Republic of Georgia/14/2011(H10N7)	CY185689.1	GenBank
A/mallard/Republic of Georgia/15/2011(H10N7)	CY185385.1	GenBank
A/northern pintail/Egypt/EMC-1/2012(H10N7)	EPI552754	GISAID EpiFlu
A/mallard/Egypt/EMC-4/2012(H10N7)	EPI552755	GISAID EpiFlu
A/mallard/Netherlands/1/2012(H10N7)	EPI552756	GISAID EpiFlu
A/shoveler/Egypt/01198-NAMRU3/2007(H10N7)	EPI372402†	GISAID EpiFlu
A/Jiangxi-Donghu/346/2013(H10N8)	EPI497477‡	GISAID EpiFlu
A/chicken/Jiangxi/102/2013(H10N8)	EPI530542§	GISAID EpiFlu

\*We gratefully acknowledge the authors, originating and submitting laboratories of the sequences from the Global Initiative on Sharing Avian Influenza Data (GISAID) EpiFlu™ database on which this research is based. All submitters of data may be contacted directly through the GISAID website.

†Originating laboratory: US Naval Medical Research Unit No.3, Egypt; Submitting laboratory: Centers for Disease Control and Prevention, Atlanta, GA, USA (1).

‡Originating and submitting laboratory: World Health Organization Chinese National Influenza Center, Beijing, China (2).

§Originating laboratory: South China Agricultural University, Guangzhou, China; Submitting laboratory: South China Agricultural University, Guangzhou, China (3).

## References

1. Gerloff NA, Jones J, Simpson N, Balish A, Elbadry MA, Baghat V, et al. A high diversity of Eurasian lineage low pathogenicity avian influenza A viruses circulate among wild birds sampled in Egypt. *PLoS ONE*. 2013;8:e68522. [PubMed http://dx.doi.org/10.1371/journal.pone.0068522](http://dx.doi.org/10.1371/journal.pone.0068522)
2. Chen H, Yuan H, Gao R, Zhang J, Wang D, Xiong Y, et al. Clinical and epidemiological characteristics of a fatal case of avian influenza A H10N8 virus infection: a descriptive study. *Lancet*. 2014;383:714–21. [PubMed http://dx.doi.org/10.1016/S0140-6736\(14\)60111-2](http://dx.doi.org/10.1016/S0140-6736(14)60111-2)
3. Qi W, Zhou X, Shi W, Huang L, Xia W, Liu D, et al. Genesis of the novel human-infecting influenza A(H10N8) virus and potential genetic diversity of the virus in poultry, China. *Euro Surveill*. 2014;19:pii: 20841. [PubMed](http://dx.doi.org/10.2807/1564-5618.2014.19.20841)