

Third Wave of Influenza A(H7N9) Virus from Poultry, Guangdong Province, China, 2014–2015

Shumin Xie,¹ Weixin Jia,¹ Yicun Lin,
Kaixiang Xing, Xingxing Ren,
Wenbao Qi, Ming Liao

Fourteen influenza A(H7N9) viruses were isolated from poultry or the environment in live poultry markets in Guangdong Province, China during 2014–2015. Phylogenetic analysis showed that all viruses were descended from viruses of the second wave of influenza A(H7N9) virus infections during 2013. These viruses can be divided into 2 branches.

A new influenza A(H7N9) virus was detected in China on February 19, 2013, and has caused worldwide concern (1). Since 2013, the outbreak of this virus in humans has occurred in 3 waves. The third wave began when 2 additional laboratory-confirmed cases of human infection with this virus were detected in Xinjiang Province, China, on September 2, 2014. This wave has continued with increasing numbers of human cases during 2015, including infections in Fujian, Hong Kong, Guizhou, Jiangsu, and Guangdong Provinces. The largest number of human cases has been reported in southern China; >50 infected patients were detected in Guangdong Province January and February (2).

The virus has been identified as a novel triple reassortant of avian influenza A(H7N3), A(H7N9), and A(H9N2) viruses and has low pathogenicity in poultry (3–5). Influenza A(H7N9) virus is now endemic to China, and its continuing reassortment in poultry makes it probable that humans will continue to be infected sporadically.

Because influenza A(H7N9) virus-contaminated live poultry markets (LPMs) are regarded as major sources of human infections with this virus (6–8), we implemented LPM sampling programs in Guangdong Province and analyzed the evolution of the virus during the third wave. In this study, we also collected samples from chicken farms and integrated epidemiologic and sequence data to infer the genetic diversity and evolution of influenza A(H7N9) viruses found in poultry in Guangdong Province, China.

The Study

Poultry surveillance for influenza A(H7N9) virus was conducted at LPMs and chicken farms in Guangdong Province

Author affiliation: South China Agricultural University, Guangzhou, China

DOI: <http://dx.doi.org/10.3201/eid2109.150635>

(4 LPMs in Guangzhou, 4 LPMs in Dongguan, 1 LPM in Shanwei, 1 LPM in Chaozhou, 2 farms in Huizhou, and 1 farm in Foshan) during September 1, 2014–February 28, 2015. Throat and cloacal swab specimens were collected every 2 weeks. Specific pathogen-free embryonated chicken eggs were used for virus isolation. Hemagglutination-positive isolates, based upon the agglutination of erythrocytes, were collected and were further subtyped by using hemagglutination inhibition assays and reverse transcription PCR.

Fourteen influenza A(H7N9) virus-positive isolates (Figure 1; online Technical Appendix Table 1, <http://wwwnc.cdc.gov/EID/article/21/9/15-0635-Techapp1.pdf>) were sequenced. Full-genome sequences generated in this study were submitted to the Global Initiative on Sharing All Influenza Data (GISAID; <http://platform.gisaid.org/epi3/frontend#41ab15>) under accession nos. EPI_ISL_176816–176820, 176824, 176828, 176830, and 176832–176837.

To understand the molecular epidemiology of these viruses, we compared our data with gene sequences of influenza A(H7N9) viruses in public databases at the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>) and GISAID on March 1, 2015. These data included all available complete gene sequences from influenza A(H7N9) viruses and sequences with high degrees of homology from other subtype virus gene sequences (hemagglutinin [HA], n = 323; neuraminidase [NA], n = 301; polymerase basic [PB] 2, n = 380; PB1, n = 286; polymerase acidic [PA], n = 286; nonstructural [NS], n = 326; nucleoprotein [NP], n = 311; and matrix [M], n = 316).

Maximum-likelihood trees were estimated for all 8 gene segments by using MEGA version 5.01 (<http://www.megasoftware.net>). To assess the robustness of individual nodes on phylogenetic trees, a bootstrap resampling process (1,000 replications), the neighbor-joining method, and the maximum composite likelihood model were used.

Phylogenetic analyses of HA genes confirmed that all third-wave influenza A(H7N9) viruses in Guangdong Province were descended from viruses of the second wave (Figure 2). It is clear that 2 H7N9 lineages co-circulate in Guangdong because third-wave viruses clustered into 2 major clades designated W3-a and W3-b, both of which emerged from the wave 2 clade. The W3-a clade contains viruses detected in Dongguan, Guangzhou, and Huizhou, and clusters of viruses

¹These authors contributed equally to this article.




Figure 1. Distribution of influenza A(H7N9) viruses, Guangdong Province, China. A) Shading indicates locations where viruses were isolated from patients during the third wave of the virus mapped according to data from the World Health Organization as of March 1, 2015. B) Circles indicate locations where influenza A(H7N9) viruses were isolated from poultry in Guangdong Province, China, during 2014–2015 (this study).

from Guangdong, Hong Kong, and Guangxi, which suggests that W3-a viruses from poultry were simultaneously prevalent in humans residing in these localities. In contrast, A/chicken/Guangdong/GZ068/2015 (H7N9) virus showed major genetic divergence from these viruses.

The W3-b clade contains viruses detected in Shanwei and Chaozhou, including A/chicken/Guangdong/CZ145/2015 (H7N9), A/chicken/Guangdong/SW153/2015(H7N9), and A/chicken/Guangdong/SW154/2015(H7N9), that clustered with strains detected in Xinjiang, Fujian, Guizhou, and Jiangsu from humans or the environment during the third wave. These data suggest regional spread of the viruses, probably by regional transport of poultry or by migratory bird populations. Phylogenetic analysis of N9 NA genes showed a topology similar to that of H7 HA genes.

Phylogenetic trees were constructed for each internal gene segment against all currently available H7N9 subtype and other subtype virus sequences (highest homology strains from BLAST [<http://blast.ncbi.nlm.nih.gov/Blast.cgi>]) from the National Center for Biotechnology Information and GISAID. Phylogenetic analysis of the whole-genome sequences showed that all 6 internal genes of DG478/2014, DG592/2014, DG593/2014, DG479/2014, DG527/2014, HZ098/2015, DG120/2015, and DG127/2015, and the PB1, PB2, PA, and NP genes of DG103/2015, DG104/2015 clustered with strains A/Guangdong/02496/2014(H7N9) and A/Hong Kong/8130773/2015(H7N9) from humans. The NS gene of DG103/2015 clustered with A/Guangdong/15SF018/2015(H7N9). The M gene clustered with A/Hong Kong/8122430/2014(H7N9).

The internal genes of CZ145/2015, SW153/2015, and SW154/2015 showed different genetic characteristics.

PB1, PB2, NP, and NS genes of SW153/2015 and SW154/2015 clustered with A/Taiwan/2/2014(H7N9), and M and PA genes were closely related to those of strains isolated in eastern China during the second wave. Internal genes, except for the PA gene of CZ145/2015, clustered with strains isolated from humans in Xinjiang. The PA gene also has a close genetic relationship with the PA gene of an H9N2 subtype strain (A/chicken/Suzhou/097–2/2013).

We conjecture that DG103/2015, CZ145/2015, SW153/2015, SW154/2015, and GZ068/2015 viruses might have undergone additional reassortment, but we cannot infer from our dataset the time, place, or with which other strains these isolates reassorted. Phylogenetic analysis of internal genes also suggested that evolution of wave 3 influenza A(H7N9) viruses resulted in a major increase in genetic diversity and sequential reassortment events with local H9N2 subtype or other subtype viruses (online Technical Appendix Figures 1–6).

We conducted mutation analyses of critical and apparent amino acid residues of influenza A(H7N9) virus isolates. All H7N9 subtype viruses isolated have an amino acid PB2-627E, PB2-701D, HA-226L(H3 numbering), NA-289R (N9 numbering), M2-31N, and HA-cleavage sites–PEIPKGRG (online Technical Appendix Table 2). These amino acid residues showed no changes when compared with those of other virus isolates from poultry. All viruses have M2-31N, which might be involved in resistance to adamantane (9). Four H7N9 subtype viruses have HA-186V (H3 numbering) and other viruses have HA-186A (H3 numbering). HA-186V may increase binding affinity for the α (2–6)-linked sialic acid receptor (10,11).




Figure 2. Phylogenetic relationships of influenza A(H7N9) virus hemagglutinin (HA) and neuraminidase (NA) genes isolated from poultry, Guangdong Province, China, 2014–2015. Phylogenetic trees were constructed by using the neighbor-joining method in MEGA software (<http://www.megasoftware.net/>). B and C are enlargements of A. Branches of the first, second, and third influenza A(H7N9) virus waves are shown in black, green, and red, respectively. Black triangles indicate newly sequenced viruses isolated from poultry in Guangdong during the third wave. Scale bars indicate nucleotide substitutions per site.

PB2-627K can enhance viral replication and virulence in a mice model (12), but all H7N9 subtype viruses in our study have PB2-627E. Thus, these strains might be less able to replicate and cause disease in mammals. Although most of the phenotypes associated with the amino acid substitutions have been demonstrated for subtypes other than H7N9, we cannot be sure that these phenotypes are also present in H7N9 subtype viruses.

Conclusions

Fourteen influenza A(H7N9) viruses were isolated from poultry or environment in LPMS in Guangdong Province,

China, during 2014–2015. Phylogenetic analyses of HA and NA genes confirmed that all third-wave influenza A(H7N9) viruses in Guangdong Province were descended from viruses of the second wave. Two H7N9 lineages from poultry co-circulated in Guangdong Province during the third wave, and both are closely related to H7N9 strains isolated from humans in local or adjacent regions. These data suggest that the dominant H7N9 strains have a dynamic evolutionary process for adapting to the local environment. Their internal genes show more regional characteristics, which might be related to transportation of live birds across provinces or to migratory birds.

The results of our study are limited by the number of samples obtained and locations of sampling. However, our findings serve as a warning to public health officials to be aware of the risk of poultry farms being infected with influenza A(H7N9) virus.

Acknowledgment

The study was performed in the National and Regional Joint Engineering Laboratory for Medicament of Zoonosis Prevention and Control, the Key Laboratory of Zoonoses Control and Prevention of Guangdong, the Key Laboratory of Animal Vaccine Development (Ministry of Agriculture).

This study was supported by grants from Program for National Broiler Industry (CARS-42-G09), the Science and Technology Projects of Guangdong Province (2013B020202001, 20140224), and the Science and Technology Projects of Dongguan (2014108101041).

Ms. Xie is a research assistant at South China Agricultural University, Guangzhou, China. Her research interest is the epidemiology of avian influenza.

References

- Chen Y, Liang W, Yang S, Wu N, Gao H, Sheng J, et al. Human infections with the emerging avian influenza A H7N9 virus from wet market poultry: clinical analysis and characterisation of viral genome. *Lancet*. 2013;381:1916–25. [http://dx.doi.org/10.1016/S0140-6736\(13\)60903-4](http://dx.doi.org/10.1016/S0140-6736(13)60903-4)
- World Health Organization. Situation updates—avian influenza [cited 2015 Apr 4]. http://www.who.int/entity/influenza/human_animal_interface/avian_influenza/archive/en/index.html
- Wu A, Su C, Wang D, Peng Y, Liu M, Hua S, et al. Sequential reassortments underlie diverse influenza H7N9 genotypes in China. *Cell Host Microbe*. 2013;14:446–52. <http://dx.doi.org/10.1016/j.chom.2013.09.001>
- Lam TT, Wang J, Shen Y, Zhou B, Duan L, Cheung CL, et al. The genesis and source of the H7N9 influenza viruses causing human infections in China. *Nature*. 2013;502:241–4. <http://dx.doi.org/10.1038/nature12515>
- Yu X, Jin T, Cui Y, Pu X, Li J, Xu J, et al. Influenza H7N9 and H9N2 viruses: coexistence in poultry linked to human H7N9 infection and genome characteristics. *J Virol*. 2014;88:3423–31. <http://dx.doi.org/10.1128/JVI.02059-13>
- Yu H, Wu JT, Cowling BJ, Liao Q, Fang VJ, Zhou S, et al. Effect of closure of live poultry markets on poultry-to-person transmission of avian influenza A H7N9 virus: an ecological study. *Lancet*. 2014;383:541–8. [http://dx.doi.org/10.1016/S0140-6736\(13\)61904-2](http://dx.doi.org/10.1016/S0140-6736(13)61904-2)
- Wu P, Jiang H, Wu JT, Chen E, He J, Zhou H, et al. Poultry market closures and human infection with influenza A(H7N9) virus, China, 2013–14. *Emerg Infect Dis*. 2014;20:1891–4. <http://dx.doi.org/10.3201/eid2011.140556>
- Yu X, Jin T, Cui Y, Pu X, Li J, Xu J, et al. Influenza H7N9 and H9N2 viruses: coexistence in poultry linked to human H7N9 infection and genome characteristics. *J Virol*. 2014;88:3423–31. <http://dx.doi.org/10.1128/JVI.02059-13>
- Pinto LH, Holsinger LJ, Lamb RA. Influenza virus M2 protein has ion channel activity. *Cell*. 1992;69:517–28. [http://dx.doi.org/10.1016/0092-8674\(92\)90452-1](http://dx.doi.org/10.1016/0092-8674(92)90452-1)
- Gambaryan AS, Matrosovich TY, Philipp J, Munster VJ, Fouchier RA, Cattoli G, et al. Receptor-binding profiles of H7 subtype influenza viruses in different host species. *J Virol*. 2012;86:4370–9. <http://dx.doi.org/10.1128/JVI.06959-11>
- Xiong X, Martin SR, Haire LF, Wharton SA, Daniels RS, Bennett MS, et al. Receptor binding by an H7N9 influenza virus from humans. *Nature*. 2013;499:496–9. <http://dx.doi.org/10.1038/nature12372>
- Hatta M, Gao P, Halfmann P, Kawaoka Y. Molecular basis for high virulence of Hong Kong H5N1 influenza A viruses. *Science*. 2001;293:1840–2. <http://dx.doi.org/10.1126/science.1062882>

Address for correspondence: Ming Liao, College of Veterinary Medicine, South China Agricultural University, Guangzhou, China; email: mliao@scau.edu.cn

UPDATE MY ADDRESS

Full text free online at
www.cdc.gov/eid

EMERGING INFECTIOUS DISEASES®

The print journal is available at no charge to public health professionals.

Yes, I still want the journal. Please send it to me at the address below.



Number on mailing label: (required) _____

Name:

Full mailing address: (BLOCK LETTERS)

Return:

Email: eideditor@cdc.gov

Fax: 404-639-1954

or mail to:

EID Editor
CDC/OID/NCEZID
1600 Clifton Rd NE
Mailstop D61
Atlanta, GA 30329-4027

Third Wave of Influenza A(H7N9) Virus from Poultry, Guangdong Province, China, 2014–2015

Technical Appendix

Additional Information on Influenza A(H7N9) Viruses and their Internal Genes, Guangdong Province, China, 2014–2015

Technical Appendix Table 1. Influenza A(H7N9) viruses isolated and sequenced in this study, Guangdong Province, China*

ID	Virus	Site	Collection date	Host
EPI_ISL_176816	A/Chicken/Guangdong/DG478/2014(H7N9)	LPM, Houjie, Dongguan	2014 Nov 25	Chicken
EPI_ISL_176817	A/Chicken/Guangdong/DG479/2014(H7N9)	LPM, Guancheng, Dongguan	2014 Nov 30	Chicken
EPI_ISL_176818	A/Duck/Guangdong/DG527/2014(H7N9)	LPM, Houjie, Dongguan	2014 Dec 2	Duck
EPI_ISL_176819	A/Chicken/Guangdong/DG592/2014(H7N9)	LPM, Houjie, Dongguan	2014 Dec 8	Chicken
EPI_ISL_176820	A/Chicken/Guangdong/DG593/2014(H7N9)	LPM, Guancheng, Dongguan	2014 Dec 12	Chicken
EPI_ISL_176824	A/Chicken/Guangdong/HZ098/2015(H7N9)	Farm, Boluo, Huizhou	2015 Jan 1	Chicken
EPI_ISL_176828	A/Duck/Guangdong/DG103/2015(H7N9)	LPM, Humen, Dongguan	2015 Jan 7	Duck
EPI_ISL_176830	A/Environment/Guangdong/DG104/2015(H7N9)	LPM, Houjie, Dongguan	2015 Jan 16	Environment
EPI_ISL_176832	A/Chicken/Guangdong/DG120/2015(H7N9)	LPM, Guancheng, Dongguan	2015 Jan 15	Chicken
EPI_ISL_176833	A/Environment/Guangdong/DG127/2015(H7N9)	LPM, Humen, Dongguan	2015 Feb 20	Environment
EPI_ISL_176834	A/Chicken/Guangdong/GZ068/2015(H7N9)	LPM, Jianggao, Guangzhou	2015 Jan 1	Chicken
EPI_ISL_176835	A/Chicken/Guangdong/CZ145/2015(H7N9)	LPM, Chaozhou	2015 Feb 1	Chicken
EPI_ISL_176836	A/Chicken/Guangdong/SW153/2015(H7N9)	LPM, Shanwei	2015 Feb 5	Chicken
EPI_ISL_176837	A/Chicken/Guangdong/SW154/2015(H7N9)	LPM, Shanwei	2015 Feb 10	Chicken

*ID, identification; LPM, live poultry market.


Technical Appendix Table 2. Mutation analysis of critical and apparent amino acid residues of influenza A(H7N9) virus isolates, Guangdong Province, China*

Virus	HA		Cleavage sites	NA	PB2		M2
	186 (H3 numbering)	226 (H3 numbering)		289 (N9 numbering)	627	701	31
A/Chicken/Guangdong/DG478/2014(H7N9)	A	L	PEIPKGRG	R	E	D	N
A/Chicken/Guangdong/DG479/2014(H7N9)	A	L	PEIPKGRG	R	E	D	N
A/Duck/Guangdong/DG527/2014(H7N9)	A	L	PEIPKGRG	R	E	D	N
A/Chicken/Guangdong/DG592/2014(H7N9)	A	L	PEIPKGRG	R	E	D	N
A/Chicken/Guangdong/DG593/2014(H7N9)	A	L	PEIPKGRG	R	E	D	N
A/Chicken/Guangdong/HZ098/2015(H7N9)	A	L	PEIPKGRG	R	E	D	N
A/Duck/Guangdong/DG103/2015(H7N9)	A	L	PEIPKGRG	R	E	D	N
A/Environment/Guangdong/DG104/2015(H7N9)	A	L	PEIPKGRG	R	E	D	N
A/Chicken/Guangdong/DG120/2015(H7N9)	A	L	PEIPKGRG	R	E	D	N
A/Environment/Guangdong/DG127/2015(H7N9)	A	L	PEIPKGRG	R	E	D	N
A/Chicken/Guangdong/GZ068/2015(H7N9)	V	L	PEIPKGRG	R	E	D	N
A/Chicken/Guangdong/CZ145/2015(H7N9)	V	L	PEIPKGRG	R	E	D	N
A/Chicken/Guangdong/SW153/2015(H7N9)	V	L	PEIPKGRG	R	E	D	N
A/Chicken/Guangdong/SW154/2015(H7N9)	V	L	PEIPKGRG	R	E	D	N


*HA, hemagglutinin; NA, neuraminidase; PB2, polymerase basic 2; M2, matrix 2. HA 186: G→V increases binding affinity for $\alpha(2-6)$ -linked sialic acid receptor (1,2). HA 226: Q→L increases binding affinity for $\alpha(2-6)$ -linked sialic acid receptor (3-5). NA 289: R→K reduces susceptibility to oseltamivir (6). PB2 627: E→K enhances viral replication and virulence in mice model (7). PB2 701: D→N causes mammalian adaptation (8). M2 31: S→N causes resistance to adamantanes (9).

References


1. Gambaryan AS, Matrosovich TY, Philipp J, Munster VJ, Fouchier RA, Cattoli G, et al. Receptor-binding profiles of H7 subtype influenza viruses in different host species. *J Virol*. 2012;86:4370–9. [PubMed](#) <http://dx.doi.org/10.1128/JVI.06959-11>
2. Xiong X, Martin SR, Haire LF, Wharton SA, Daniels RS, Bennett MS, et al. Receptor binding by an H7N9 influenza virus from humans. *Nature*. 2013;499:496–9. [PubMed](#) <http://dx.doi.org/10.1038/nature12372>
3. Connor RJ, Kawaoka Y, Webster RG, Paulson JC. Receptor specificity in human, avian, and equine H2 and H3 influenza virus isolates. *Virology*. 1994;205:17–23. [PubMed](#) <http://dx.doi.org/10.1006/viro.1994.1615>
4. Ha Y, Stevens DJ, Skehel JJ, Wiley DC. X-ray structures of H5 avian and H9 swine influenza virus hemagglutinins bound to avian and human receptor analogs. *Proc Natl Acad Sci U S A*. 2001;98:11181–6. [PubMed](#) <http://dx.doi.org/10.1073/pnas.201401198>
5. Herfst S, Schrauwen EJ, Linster M, Chutinimitkul S, de Wit E, Munster VJ, et al. Airborne transmission of influenza A/H5N1 virus between ferrets. *Science*. 2012;336:1534–41. [PubMed](#) <http://dx.doi.org/10.1126/science.1213362>
6. Colman PM, Varghese JN, Laver WG. Structure of the catalytic and antigenic sites in influenza virus neuraminidase. *Nature*. 1983;303:41–4. [PubMed](#) <http://dx.doi.org/10.1038/303041a0>
7. Hatta M, Gao P, Halfmann P, Kawaoka Y. Molecular basis for high virulence of Hong Kong H5N1 influenza A viruses. *Science*. 2001;293:1840–2. [PubMed](#) <http://dx.doi.org/10.1126/science.1062882>
8. Gao Y, Zhang Y, Shinya K, Deng G, Jiang Y, Li Z, et al. Identification of amino acids in HA and PB2 critical for the transmission of H5N1 avian influenza viruses in a mammalian host. *PLoS Pathog*. 2009;5:e1000709. [PubMed](#) <http://dx.doi.org/10.1371/journal.ppat.1000709>
9. Pinto LH, Holsinger LJ, Lamb RA. Influenza virus M2 protein has ion channel activity. *Cell*. 1992;69:517–28. [PubMed](#) [http://dx.doi.org/10.1016/0092-8674\(92\)90452-I](http://dx.doi.org/10.1016/0092-8674(92)90452-I)



Technical Appendix Figure 1. Phylogenetic relationships of polymerase basic 2 genes of influenza A(H7N9) viruses from poultry, Guangdong Province, China, 2014–2015. Trees were constructed by using the neighbor-joining method in MEGA (<http://www.megasoftware.net/>). Red triangles indicate newly sequenced viruses isolated from poultry in Guangdong during the third wave. Scale bar indicates nucleotide substitutions per site.




Technical Appendix Figure 2. Phylogenetic relationships of polymerase basic 1 genes of influenza A(H7N9) viruses from poultry, Guangdong Province, China, 2014–2015. Trees were constructed by using the neighbor-joining method in MEGA (<http://www.megasoftware.net/>). Red triangles indicate newly sequenced viruses isolated from poultry in Guangdong during the third wave. Scale bar indicates nucleotide substitutions per site.




Technical Appendix Figure 3. Phylogenetic relationships of polymerase acidic genes of influenza A(H7N9) viruses from poultry, Guangdong Province, China, 2014–2015. Trees were constructed by using the neighbor-joining method in MEGA (<http://www.megasoftware.net/>). Red triangles indicate newly sequenced viruses isolated from poultry in Guangdong during the third wave. Scale bar indicates nucleotide substitutions per site.



Technical Appendix Figure 4. Phylogenetic relationships of nonstructural protein genes of influenza A(H7N9) viruses from poultry, Guangdong Province, China, 2014–2015. Trees were constructed by using the neighbor-joining method in MEGA (<http://www.megasoftware.net/>). Red triangles indicate newly sequenced viruses isolated from poultry in Guangdong during the third wave. Scale bar indicates nucleotide substitutions per site.



Technical Appendix Figure 5. Phylogenetic relationships of nucleoprotein genes of influenza A(H7N9) viruses from poultry, Guangdong Province, China, 2014–2015. Trees were constructed by using the neighbor-joining method in MEGA (<http://www.megasoftware.net/>). Red triangles indicate newly sequenced viruses isolated from poultry in Guangdong during the third wave. Scale bar indicates nucleotide substitutions per site.



Technical Appendix Figure 6. Phylogenetic relationships of matrix genes of influenza A(H7N9) viruses from poultry, Guangdong Province, China, 2014–2015. Trees were constructed by using the neighbor-joining method in MEGA (<http://www.megasoftware.net/>). Red triangles indicate newly sequenced viruses isolated from poultry in Guangdong during the third wave. Scale bar indicates nucleotide substitutions per site.