
Risk Factors for MERS-CoV Seropositivity among Animal Market and Slaughterhouse Workers, Abu Dhabi, United Arab Emirates, 2014–2017

Ahmed Khudhair,¹ Marie E. Killerby,¹ Mariam Al Mulla, Kheir Abou Elkheir, Wassim Ternanni, Ziad Bandar, Stefan Weber, Mary Khoury, George Donnelly, Salama Al Muhairi, Abdelmalik I. Khalafalla, Suvang Trivedi, Azaibi Tamin, Natalie J. Thornburg, John T. Watson, Susan I. Gerber, Farida Al Hosani,² Aron J. Hall²

Camel contact is a recognized risk factor for Middle East respiratory syndrome coronavirus (MERS-CoV) infection. Because specific camel exposures associated with MERS-CoV seropositivity are not fully understood, we investigated worker–camel interactions and MERS-CoV seroprevalence. We assessed worker seroprevalence in 2 slaughterhouses and 1 live-animal market in Abu Dhabi, United Arab Emirates, during 2014–2017 and administered an epidemiologic survey in 2016 and 2017. Across 3 sampling rounds during 2014–2017, we sampled 100–235 workers, and 6%–19% were seropositive for MERS-CoV at each sampling round. One (1.4%) of 70 seronegative workers tested at multiple rounds seroconverted. On multivariable analyses, working as a camel salesman, handling live camels or their waste, and having diabetes were associated with seropositivity among all workers, whereas handling live camels and either administering medications or cleaning equipment was associated with seropositivity among market workers. Characterization of high-risk exposures is critical for implementation of preventive measures.

Middle East respiratory syndrome (MERS) coronavirus (MERS-CoV) was first identified as a cause of severe respiratory tract infections in Saudi Arabia in October 2012 (1). The clinical spectrum of MERS ranges from asymptomatic infection to acute respiratory distress

syndrome and death (2). As of April 3, 2019, a total of 2,374 laboratory-confirmed cases of infection have been reported by 27 countries to the World Health Organization (WHO); the reported case-fatality rate is 35% (2). All reported cases have an epidemiologic link to the Arabian Peninsula, and imported cases have been reported in Europe, Asia, North America, and Africa. The United Arab Emirates has reported the third-highest number of MERS cases since 2012 (3).

MERS-CoV is a zoonotic virus, and dromedaries (camels) are recognized as a major virus reservoir for spillover to humans (4). Multiple studies have isolated MERS-CoV or MERS-CoV RNA from camels across the Arabian Peninsula and Africa (5–11). Serologic studies of camels in the Middle East and Africa have revealed MERS-CoV seroprevalence of >90%–97% (8, 11–13). In natural infection, camels have been found to shed MERS-CoV in respiratory secretions and to a lesser extent in stool (14, 15). Evidence of virus RNA has also been found in milk collected by traditional milking procedures, which involve calf suckling as a stimulus for milk letdown (15).

Epidemiologic links between infected camels and human MERS-CoV infections have been shown, with identical or nearly identical MERS-CoV genomes found in human cases and in camels with which they had direct contact (16–18). Also, a case–control study identified exposure to camels as a risk factor for human MERS-CoV infection (19). Human seroprevalence studies also support the association between MERS-CoV infection and camel contact; in Saudi Arabia MERS-CoV seroprevalence was found to be 15 times greater in camel shepherds and 23 times greater in slaughterhouse workers compared with the

Author affiliations: Abu Dhabi Department of Health, Abu Dhabi, United Arab Emirates (A. Khudhair, M. Al Mulla, K.A. Elkheir, W. Ternanni, Z. Bandar, F. Al Hosani); Sheikh Khalifa Medical City, Abu Dhabi (M.E. Killerby, S. Weber, M. Khoury, G. Donnelly); Centers for Disease Control and Prevention, Atlanta, Georgia, USA (M.E. Killerby, S. Trivedi, A. Tamin, N.J. Thornburg, J.T. Watson, S.I. Gerber, A.J. Hall); Abu Dhabi Food Control Authority, Abu Dhabi (S. Al Muhairi, A.I. Khalafalla)

DOI: <https://doi.org/10.3201/eid2505.181728>

¹These authors contributed equally to this article.

²These authors contributed equally to this article.

general population (20). Further studies have also shown high seroprevalence in specific occupational groups with various camel exposures (e.g., seropositivity was detected in 6.8% of a cohort of 294 camel workers in Qatar [21] and in 53% of a cohort of 30 camel workers in Saudi Arabia [22]).

Although multiple lines of evidence suggest camel exposure is associated with human MERS-CoV infection, the exact mechanisms of transmission are not fully understood. Information on specific risk factors relating to camel interactions are needed to further understand how the virus might be transmitted from camels to humans and to guide interventions to prevent zoonotic transmission, including changes to camel management practices. Because MERS-CoV vaccines are currently in development and have reported success in phase I clinical trials (23), knowledge of groups at risk for MERS-CoV infection might also be useful when considering future vaccine use. Our study aimed to identify risk factors for MERS-CoV seropositivity among live-animal market and slaughterhouse workers.

Methods

Study Site and Population

The study sites consisted of an open-air animal market and 2 slaughterhouses (1 commercial and 1 public). All 3 facilities housed camels, goats, sheep, and cattle (Figure 1). Typically during the study period, approximately 460 persons worked at the market, 101 at the public slaughterhouse, and 29 at the commercial slaughterhouse. The market investigated in this study was linked to a human MERS case in 2015 (24). Prior investigation showed a large diversity of MERS-CoVs circulating among camels at the market; 109 (29%) of 276 screened camels had detectable MERS-CoV RNA in nasal swab specimens in the spring of 2015 (25).

Serum Sampling and Data Collection

We conducted 3 rounds of worker serum sampling. The first round was conducted during May 11–14, 2014, and the second round during March 23–April 1 and May 7–13, 2015. During the first 2 rounds of sampling, all available workers at the market and public slaughterhouse were requested to provide a serum sample as part of a public health investigation. We conducted a third round of serum sampling during September 22–October 5, 2016, and March 20–23, 2017. The third round of sampling included workers at the market, public slaughterhouse, and the newly opened commercial slaughterhouse. All available workers were requested to provide serum samples, although participation was voluntary. Some, but not all, workers were repeatedly sampled, when feasible, during multiple rounds.

We administered an epidemiologic survey to all workers only during the third round of serum sampling in 2016 and 2017. No surveys were administered in 2014 or




Figure 1. Diagram of study site indicating market and slaughterhouse settings, Abu Dhabi, United Arab Emirates.

2015. The survey consisted of questions covering worker demographics; occupational history; contact with various animal species; travel history; medical history; consumption of raw camel milk, raw camel meat, and camel urine; specific tasks performed with camels; types of personal protective equipment (PPE) worn; and handwashing practices (Appendix 1, <https://wwwnc.cdc.gov/EID/article/25/5/18-1728-App1.pdf>). Separate lists of questions covering specific camel tasks performed were asked of market and slaughterhouse workers because of the different nature of camel tasks among occupational groups. Interviews were conducted in Arabic by staff from the Abu Dhabi Department of Health.

Laboratory Testing

Human serum samples were tested for MERS-CoV antibodies at the US Centers for Disease Control and Prevention (CDC) by using indirect ELISAs for nucleocapsid (N) and spike (S) proteins, followed by a confirmatory microneutralization test, as previously described (26). Samples were initially tested by using both N and S ELISAs as screening assays with serum diluted to 1:400. All serum samples with optical densities above assay cutoff were diluted serially, 4-fold, from 1:100 to 1:6,400, and used for endpoint titer determinations. Serum samples that were positive by N or S ELISA with titers at 1:400, 1:1600, or 1:6,400, plus 10% of samples negative by N or S ELISA at these titers, were tested by using microneutralization with live MERS-CoV performed in a Biosafety Level 3 laboratory, as previously described (26). In addition, we conducted confirmatory microneutralization tests on seronegative samples from any persons who showed a change in seropositivity status over time to confirm changes in seropositivity status. Samples were considered positive if positive on N and S ELISA or if positive on microneutralization. Specimens near the limits of detection but not consistently above or below these limits were considered indeterminate. For the epidemiologic analysis, persons with an indeterminate result were considered seronegative.

Data Analyses

We used Epi Info 7 (<https://www.cdc.gov/epiinfo>) for data entry and R version 3.3.1 (<https://cran.r-project.org/bin/windows/base/old/3.3.1>) for data analysis. We performed comparisons between prevalence of work practices by setting (market vs. slaughterhouse) by using the Pearson χ^2 square test. We used univariable logistic regression to estimate odds ratios, 95% CIs, and p values (Wald test) for all associations between potential risk factors and seropositivity. We assessed associations between demographics, occupational history, contact with various animal species, consumption of camel products, travel history, and medical history with seropositivity for all workers. We separately tested associations between specific interactions with camels, types of PPE worn, and handwashing practices with seropositivity for stratified subgroups of market and slaughterhouse workers because of the different nature of work setting and standard practices between these 2 populations. We then performed additional exploratory data description by occupation on the basis of results of univariable analyses.

We developed 3 multivariable logistic models to identify associations between risk factors and seropositivity. First, we constructed a model of risk factors common to all workers and then constructed occupationally stratified models (i.e., separate models for market workers and slaughterhouse workers) to model specific interactions with camels, PPE use, and handwashing practices. We combined or eliminated highly correlated variables, which were determined by condition indices and variance decomposition proportions. We reduced categorical variables to binary options if small group size was observed. We performed initial variable selection by using least absolute shrinkage and selection operator (LASSO) and then tested person-variable significance by using the likelihood ratio test with a cutoff of $p < 0.05$ within an ordinary logistic regression model. We then included age and number of years worked at current setting as potential confounders in all 3 final models. We excluded persons with missing data at the LASSO stage but included them for the final logistic regression model.

For the stratified market worker and slaughterhouse models, we also included variables significant in the all workers model but not directly relating to camel interactions (e.g., reported underlying conditions) in the final occupationally stratified models. We did not include significant variables directly relating to camel exposures in the

all workers model in the stratified models because more specific camel risk practices were assessed in the stratified models. For market and slaughterhouse models, we tested interactions between significant risk practices and select PPE use and handwashing practices for a protective effect.

Results

Serum Sample Results

We sampled 100 workers in round 1 (2014), 151 workers in round 2 (2015), and 235 workers in round 3 (2016 and 2017); overall MERS-CoV seroprevalence was 6% for round 1, 19% for round 2, and 17% for round 3. Twenty-one persons had specimens taken at rounds 1 and 2, twenty-three at rounds 2 and 3, thirteen at rounds 1 and 3, and twenty-two at all 3 rounds (Figure 2). Of 70 persons who were seronegative at their first sample, only 1 (1.4%, 95% CI 0.1%–8.8%) seroconverted: a 30-year-old man who was a cleaner at the public slaughterhouse tested negative at round 1 and positive at round 2. Of 8 persons who were seropositive at their first sample, 1 (13%) was later found to be seronegative: a 28-year-old man who was an administrative supervisor at the market was resampled between rounds 1 and 3. This person did not report handling camels or their waste and did not perform any tasks directly relating to camels. One additional person who had a positive serologic result at their first and second samples and an indeterminate result at their third sample was not subsequently evaluated for change in seropositive status. Because some study participants might have had different medical record numbers across the 3 sampling rounds, we could not determine all potential seroconversions or losses of seropositivity, although we also performed matching by name and age. We compiled serologic results for all participants who ever tested positive (Appendix 2, <https://wwwnc.cdc.gov/EID/article/25/5/18-1728-App2.pdf>).

Epidemiologic Survey Results

In total, 235 persons both completed the epidemiologic survey and were sampled during round 3. One additional person completed the epidemiologic survey but refused serum sampling and was not included in any analyses. All 235 workers were men, and their median age was 35 years (range 19–64 years). The median number of years worked at the current settings was 6 (range 0.2–15 years). We observed no significant effect of age ($p = 0.26$) or years worked ($p = 0.18$) on seropositivity on univariable analysis.




Figure 2. Number of workers sampled during >1 round of sampling (N = 79), Abu Dhabi, United Arab Emirates. Black stars indicate when serum samples were taken; gray shading indicates follow-up periods.

Worker occupations were categorized into animal handlers (n = 16), camel salesmen (n = 37), other animal salesmen (n = 41), animal or waste transporters (n = 27), butchers (n = 65), cleaners (n = 26), veterinarians (n = 9), and other (e.g., supervisor, cashier, and tourist guide) (n = 14). Salesmen only worked in the market, and butchers only worked in the slaughterhouses. The remaining occupations were found in both settings, but each person could only work at a slaughterhouse or the market. None of the workers reported working at any other job outside of the market or slaughterhouses, and the only animals reported present at home were poultry and stray cats.

Overall, 64 (44%) of 145 market workers had daily contact with camels or their waste, compared with 47 (52%) of 90 slaughterhouse workers (p = 0.28). Certain PPE use and handwashing were more frequently reported by slaughterhouse workers than market workers. Among slaughterhouse workers, 99% reported wearing a dust mask (equivalent to a surgical mask), compared with 21% of market workers (p<0.01). Only 37% of slaughterhouse workers reported

taking their work clothes home, compared with 97% of market workers (p<0.01). Eighty-one percent of slaughterhouse workers reported washing their hands before and after each animal-related task, compared with 21% of market workers (p<0.01). Ninety-three percent of slaughterhouse workers reported washing their hands at the beginning and end of the day, compared with only 56% of market workers (p<0.01).

Univariable Analyses

Rates of seropositivity were higher among market workers (29 [20%] of 145) than among slaughterhouse workers (11 [12%] of 90), although this difference was not statistically significant on univariable analysis (p = 0.17). By occupation, camel salesmen and animal or waste transporters had significantly higher odds of seropositivity than the reference group of other salesmen (Table 1).

Univariable analyses showed that several characteristics were associated with seropositivity among all workers (Table 1), including handling camels or their waste daily. Not all seropositive workers reported handling camels or

Table 1. Characteristics of 235 market and slaughterhouse workers, by MERS-CoV serostatus, Abu Dhabi, United Arab Emirates*

Characteristic	Total no. participants	No. (%) participants		OR (95% CI)	p value
		Seronegative, n = 195	Seropositive, n = 40		
Work >50 h/wk	132	103 (78.0)	29 (22.0)	2.35 (1.14–5.17)	0.025
Worked another job in previous year	30	25 (83.3)	5 (16.7)	0.97 (0.31–2.53)	0.956
Occupation					
Other salesman	41	40 (97.6)	1 (2.4)	Ref	Ref
Animal handler	16	15 (93.8)	1 (6.3)	2.67 (0.1–70.37)	0.498
Butcher	65	56 (86.2)	9 (13.8)	6.43 (1.14–120.92)	0.083
Camel salesman	37	19 (51.4)	18 (48.6)	37.89 (7.03–707.16)	0.001
Cleaner	26	22 (84.6)	4 (15.4)	7.27 (1–147.06)	0.084
Animal or waste transporter	27	21 (77.8)	6 (22.2)	11.43 (1.79–223.48)	0.029
Veterinarian	9	8 (88.9)	1 (11.1)	5.00 (0.18–135.74)	0.272
Other	14	14 (100.0)	0	NA	0.990
Nationality					
Afghani	27	15 (55.6)	12 (44.4)	Ref	Ref
Bangladeshi	38	30 (78.9)	8 (21.1)	0.33 (0.11–0.97)	0.048
Pakistani	97	92 (94.8)	5 (5.2)	0.07 (0.02–0.21)	<0.001
Sudanese	38	25 (65.8)	13 (34.2)	0.65 (0.23–1.79)	0.404
Other	35	33 (94.3)	2 (5.7)	0.08 (0.01–0.32)	0.002
Contact with					
Camels or waste daily	111	78 (70.3)	33 (29.7)	7.07 (3.14–18.15)	<0.001
Cattle or waste daily	58	50 (86.2)	8 (13.8)	0.73 (0.29–1.61)	0.452
Goats or waste daily	88	79 (89.8)	9 (10.2)	0.43 (0.18–0.91)	0.036
Sheep or waste daily	89	80 (89.9)	9 (10.1)	0.42 (0.18–0.89)	0.031
Drank raw camel milk	25	18 (72.0)	7 (28.0)	2.09 (0.76–5.21)	0.129
Ate raw camel meat†	2	1 (50.0)	1 (50.0)	4.95 (0.19–126.94)	0.262
Travel					
To Saudi Arabia	4	4 (100.0)	0	NA	0.990
Within UAE	67	55 (82.1)	12 (17.9)	1.09 (0.5–2.25)	0.819
Underlying conditions					
Asthma	1	0	1 (100.0)	NA	0.985
Diabetes	9	5 (55.6)	4 (44.4)	4.22 (1–16.71)	0.038
Hypertension	12	9 (75.0)	3 (25.0)	1.68 (0.36–5.93)	0.455
Sought care for respiratory illness	45	36 (80.0)	9 (20.0)	1.28 (0.54–2.84)	0.555
Contact with anyone with respiratory illness	2	1 (50.0)	1 (50.0)	1.38 (0.09–7.84)	0.731
Had chest radiograph	13	10 (76.9)	3 (23.1)	1.5 (0.32–5.18)	0.552

*MERS-CoV, Middle East respiratory syndrome coronavirus; NA, not applicable; OR, odds ratio; Ref, referent; UAE, United Arab Emirates.

†Total seronegative was 194 because answer from 1 person was missing. No workers ever contacted the following species: dogs, cats, bats, rodents, birds, and other animals; 1 worker reported rarely contacting horses.

their waste; 7 workers initially claimed they never handled camels or their waste, although 3 of these later reported that they contacted either camel equipment, viscera, or waste within the slaughterhouse. For the subgroup of market workers, univariable analyses revealed multiple camel exposures to be associated with seropositivity and 2 handwashing practices that were inversely associated with seropositivity (Table 2). For the subgroup of slaughterhouse workers, no individual risk factors were associated with seropositivity (Table 3).

Because camel salesmen had the highest odds of MERS-CoV seropositivity, we summarized their frequency of specific camel exposures separately (Figure 3). Direct observation of camel salesmen in the market showed that most of their time was spent in the camel pens, including while they ate and rested, and direct handling of the animals occurred frequently (data not shown).

Multivariable Analyses

For the multivariable model evaluating risk factors associated with seropositivity in all workers, the following variables remained in the final logistic regression model: handling camels or their waste daily (adjusted odds ratio [aOR]

4.2, 95% CI 1.7–11.8), working as a camel salesman (aOR 4.0, 95% CI 1.6–10.1), and self-reported diabetes (aOR 6.2, 95% CI 1.2–30.3). All 3 factors significantly increased odds of seropositivity.

For market workers, multivariable analysis resulted in a final model in which the following variables were each independently associated with seropositivity: handling live camels (aOR 12.2, 95% CI 3.2–62.9), administering medications to camels (aOR 3.4, 95% CI 1.1–11.2), and self-reported diabetes (aOR 20.9, 95% CI 1.6–341.3). Cleaning equipment was also significantly associated with seropositivity (aOR 3.3, 95% CI 1.1–10.3); substituted for administering medication to camels, this factor produced a model with a near-identical fit along with the other risk factors. Given that administering medications to camels was highly correlated with cleaning equipment, the statistical significance of both factors was lost if both factors were included in the model because of collinearity ($\rho = 0.65$). None of the select PPE and handwashing practices evaluated as interactions with risk practices showed a significant protective effect. No individual risk factors were significantly associated with slaughterhouse workers by multivariable analysis.

Table 2. Comparison of practices among 145 MERS-CoV seronegative and seropositive market workers, Abu Dhabi, United Arab Emirates*

Characteristic	Total no. participants	No. (%) participants		OR (95% CI)	p value
		Seronegative, n = 116	Seropositive, n = 29		
Handle live camels	66	40 (60.6)	26 (39.4)	16.47 (5.38–72.06)	<0.001
Feed camels	51	31 (60.8)	20 (39.2)	6.09 (2.57–15.44)	<0.001
Clean camels	39	22 (56.4)	17 (43.6)	6.05 (2.56–14.83)	<0.001
Clean camel housing	37	19 (51.4)	18 (48.6)	8.35 (3.47–21.11)	<0.001
Handle camel waste	33	16 (48.5)	17 (51.5)	8.85 (3.63–22.56)	<0.001
Clean equipment	33	16 (48.5)	17 (51.5)	8.85 (3.63–22.56)	<0.001
Milk camels	18	12 (66.7)	6 (33.3)	2.26 (0.72–6.49)	0.138
Assist with camel birthing	27	18 (66.7)	9 (33.3)	2.45 (0.94–6.16)	0.060
Give medications to camels	35	18 (51.4)	17 (48.6)	7.71 (3.2–19.34)	<0.001
Contact with ill camel	37	20 (54.1)	17 (45.9)	6.8 (2.85–16.83)	<0.001
Wear dust mask and gloves	25	23 (92.0)	2 (8.0)	0.3 (0.05–1.1)	0.117
Wear respirator	0	0 (NA)	0 (NA)	NA	NA
Wear coveralls	15	13 (86.7)	2 (13.3)	0.59 (0.09–2.3)	0.500
Wear boots	3	3 (100.0)	0 (0.0)	NA	0.991
Who washes your clothes?					
Self	84	69 (82.1)	15 (17.9)	Ref	Ref
Household member	5	4 (80.0)	1 (20.0)	1.15 (0.06–8.49)	0.904
Other worker	56	43 (76.8)	13 (23.2)	1.39 (0.6–3.21)	0.439
Take work clothes home	141	113 (80.1)	28 (19.9)	0.74 (0.09–15.34)	0.801
Wash work clothes at home	89	73 (82.0)	16 (18.0)	0.72 (0.32–1.67)	0.444
Wash work clothes at workplace	6	3 (50.0)	3 (50.0)	4.35 (0.77–24.67)	0.082
Wash work clothes at laundry	63	50 (79.4)	13 (20.6)	1.07 (0.47–2.43)	0.867
Wash hands before and after each animal-related task	40	30 (75.0)	10 (25.0)	1.51 (0.61–3.56)	0.355
Wash hands at mealtimes	145	116 (80.0)	29 (20.0)	NA	NA
Wash hands at bathroom times	145	116 (80.0)	29 (20.0)	NA	NA
Wash hands at prayer times	145	116 (80.0)	29 (20.0)	NA	NA
Wash hands at beginning and end of day	81	70 (86.4)	11 (13.6)	0.4 (0.17–0.92)	0.033
Wash hands at toilets	143	114 (79.7)	29 (20.3)	NA	0.989
Wash hands at restaurant	64	50 (78.1)	14 (21.9)	1.23 (0.54–2.8)	0.616
Wash hands at mosque	69	55 (79.7)	14 (20.3)	1.04 (0.45–2.35)	0.934
Wash hands at barn	4	1 (25.0)	3 (75.0)	13.27 (1.63–274.19)	0.028

*MERS-CoV, Middle East respiratory syndrome coronavirus; NA, not applicable; OR, odds ratio.

Discussion

Our study investigated risk factors for MERS-CoV seropositivity in animal market and slaughterhouse workers at a site previously associated with zoonotic transmission of MERS-CoV. Given the large number of camels present, including many young camels, and the mixing of camels from multiple sources, this site probably facilitates MERS-CoV transmission among camels. Our results demonstrated a relatively high MERS-CoV seroprevalence in workers at this site, ranging from 6% to 19% at each round across all occupations. Because we did not record occupation and other risk factors during the first 2 sampling rounds, we were unable to further assess reasons for the different seropositivity rates between sampling rounds.

We found particularly high seroprevalence in specific occupational groups, namely camel salesmen (49%) and animal or waste transporters (22%). Previous studies of workers with occupational exposure to camels have reported either lower seropositivity rates (e.g., 6.8% of 294 workers with occupational camel contact seropositive in Qatar [21] and 2.3% of 87 camel shepherds seropositive in Saudi Arabia [20]) or comparable seropositivity (e.g., 53% of camel workers positive in Saudi Arabia [22]). Our rates of seropositivity

might underestimate actual exposure to MERS-CoV. Previous studies have demonstrated that examining MERS-CoV-specific T cells from MERS patients is more sensitive than examining serum antibodies alone (27). To examine T-cell responses, peripheral blood mononuclear cells must be collected, which was beyond the scope of our study.

On multivariable analysis, we found that contact with camels or their waste, working as a camel salesman, and self-reported diabetes were all independently associated with seropositivity in all workers. Because of small stratum size, belonging to other occupational groups could not be meaningfully explored as risk factors. Diabetes has previously been shown to be a commonly reported underlying condition in MERS cases (28), has been associated with risk for infection in a case-control study (19), and has been associated with increased risk for death in MERS patients (29). We found an association between diabetes and MERS-CoV seropositivity in a cohort with occupational exposure to camels. Although persons with diabetes might be at increased risk for MERS-CoV infection, the association between diabetes, MERS-CoV infection, and the resulting antibody response is still not fully understood.

Table 3. Comparison of practices among 90 MERS-CoV seronegative and seropositive slaughterhouse workers, Abu Dhabi, United Arab Emirates*

Characteristic	Total no. participants	No. (%) participants		OR (95% CI)	p value
		Seronegative, n = 79	Seropositive, n = 11		
Handle live camels	56	49 (87.5)	7 (12.5)	1.07 (0.3–4.37)	0.918
Perform antemortem exam of camels	6	5 (83.3)	1 (16.7)	1.48 (0.07–10.5)	0.732
Remove hide from camels†	52	46 (88.5)	6 (11.5)	0.83 (0.23–3.12)	0.780
Remove or handle viscera of camels‡	55	48 (87.3)	7 (12.7)	1.51 (0.39–7.4)	0.573
Clean equipment	68	58 (85.3)	10 (14.7)	3.62 (0.63–68.47)	0.233
Handle camel waste	57	50 (87.7)	7 (12.3)	1.01 (0.28–4.15)	0.982
Prepare cuts of camel meat	52	45 (86.5)	7 (13.5)	1.32 (0.37–5.39)	0.675
Conduct postmortem exam of camels	3	2 (66.7)	1 (33.3)	3.85 (0.17–43.92)	0.288
Slaughter camels	41	36 (87.8)	5 (12.2)	1 (0.27–3.57)	0.994
Contact with ill camel	3	2 (66.7)	1 (33.3)	3.85 (0.17–43.92)	0.288
Wear dust mask and gloves	88	77 (87.5)	11 (12.5)	NA	NA
Wear respirator	0	0	0	NA	NA
Wear coveralls	85	74 (87.1)	11 (12.9)	NA	NA
Wear boots	85	74 (87.1)	11 (12.9)	NA	NA
Who washes your clothes					
Self	30	28 (93.3)	2 (6.7)	Ref	Ref
Household member	2	0	2 (100.0)	NA	NA
Other worker	58	51 (87.9)	7 (12.1)	1.92 (0.43–13.49)	0.434
Take work clothes home	33	29 (87.9)	4 (12.1)	0.99 (0.24–3.55)	0.982
Wash work clothes at home	31	27 (87.1)	4 (12.9)	1.1 (0.27–3.98)	0.886
Wash work clothes at workplace	58	51 (87.9)	7 (12.1)	0.96 (0.27–3.93)	0.952
Wash work clothes at laundry	2	2 (100.0)	0	NA	NA
Wash hands before and after each animal-related task	73	64 (87.7)	9 (12.3)	1.05 (0.24–7.39)	0.949
Wash hands at mealtimes	89	78 (87.6)	11 (12.4)	NA	NA
Wash hands at bathroom times	90	79 (87.8)	11 (12.2)	NA	NA
Wash hands at prayer times	88	78 (88.6)	10 (11.4)	0.13 (0–3.41)	0.158
Wash hands at beginning and end of day	84	75 (89.3)	9 (10.7)	0.24 (0.04–1.9)	0.127
Wash hands at toilets	86	76 (88.4)	10 (11.6)	0.39 (0.05–8.4)	0.440
Wash hands at restaurant	0	0	0	NA	NA
Wash hands at mosque	3	3 (100.0)	0	NA	NA
Wash hands at basin	47	43 (91.5)	4 (8.5)	0.48 (0.12–1.71)	0.268

*Exam, examination; MERS-CoV, Middle East respiratory syndrome coronavirus, NA, not applicable; OR, odds ratio.

†Total seronegative was 78 because answer from 1 person was missing.

‡Total seropositive was 10 because answer from 1 person was missing.




Figure 3. Frequency of tasks performed by camel salesmen (N = 37) in market, Abu Dhabi, United Arab Emirates.

However, because persons with diabetes are considered at high risk for developing severe disease from MERS-CoV infection, WHO recommends these persons take precautions when visiting farms or markets where camels are present, including avoiding contact with camels (3).

Among market workers, handling live camels and either administering medications to camels or cleaning equipment were practices associated with significantly increased risk for MERS-CoV seropositivity. Given that administering medications to camels was highly correlated with cleaning equipment, neither factor was statistically significant if both were included in the model. The biological importance of these associations might therefore be difficult to interpret, because either or both risk factors could be statistically associated with MERS-CoV seropositivity and have an undefined strength of association. Practices potentially associated with camel calves, such as milking or assisting with camel birth, were not associated with MERS-CoV seropositivity despite a higher prevalence of viral RNA in camels <1 year of age compared with other ages (30) and a previously reported association between milking camels frequently and seropositivity (31). However, these practices were not commonly reported by market workers in our study, limiting the power to detect an association with seropositivity.

No specific work practices were found to be associated with seropositivity among slaughterhouse workers. Compared with market workers, slaughterhouse workers had less exposure to live camels and a higher self-reported prevalence of potentially protective practices such as PPE use and frequent handwashing. Although our multivariable analysis did not show a significant association between PPE use (e.g., wearing a dust mask and gloves) or handwashing practices and seropositivity, the small sample size might have restricted the power to detect interactions between PPE and camel exposures. Because camel-to-human transmission of MERS-CoV is not fully understood, WHO recommends broad preventive measures for slaughterhouse and market workers, including

wearing facial protection when feasible, washing hands before and after each animal-related task, and washing soiled work clothes and shoes at the work place to avoid exposing family members to soiled work clothing (3). Where feasible, increased use of such measures could be encouraged, particularly in market workers, to decrease risk for infection.

Because only a single human MERS case has been reported in connection with the study site, our reported rates of seroprevalence suggest unrecognized transmission (and potentially unrecognized illness) at this site. However, because the length of time MERS-CoV antibodies persist is unknown (32), the time and place these infections might have occurred is unknown; transmission potential also exists in the United Arab Emirates outside of markets and slaughterhouses. Whether infections were symptomatic is also unknown. Participants were asked whether they had seen a healthcare provider for respiratory illness in the previous 12 months, but such reported illness was not associated with seropositivity, and multiple pathogens other than MERS-CoV could be responsible for any reported respiratory illness. Despite these limitations, MERS-CoV was detected in camels at the market during our study period (25), and an interim seroconversion was noted in 1 worker, suggesting active zoonotic transmission. Taken collectively, our findings suggest an underestimated prevalence of human MERS-CoV infection in settings where the virus is circulating among camels, probably resulting from camel-to-human transmission.

Our study had additional limitations, including the overall sample size and limited number of subjects within specific substrata. Concentration of camel interactions within particular occupational groups limited our ability to differentiate risk among specific camel interactions, despite our use of multivariable analysis. Furthermore, because most persons reported interactions either daily or never, determining whether increased risk was associated with increased frequency of individual tasks was not possible. Also, some MERS-CoV infections might not result in detectable

antibodies, particularly when the infections are asymptomatic or mild (32). Persistence of detectable MERS-CoV antibodies after infection is not well-defined, limiting the ability of serologic testing to define previous infection. Finally, because of incomplete linkage of study participants by medical record numbers across the 3 sampling periods, not all potential seroconversions or losses of seropositivity could be determined.

In summary, our study found significantly increased odds of MERS-CoV seropositivity in persons with exposure to camels, in particular among those who handle live camels. Odds of seropositivity were also significantly higher for camel salesmen, suggesting that preventive measures such as PPE use could focus on specific occupational groups, in addition to individual work practices. Determining groups at highest risk for zoonotic MERS-CoV infection could also inform future vaccine trials in geographic regions where MERS-CoV is known to circulate.

Acknowledgments

We thank Yassir Eltahir for critical input during the conceptualization and implementation of this study.

CDC and the Abu Dhabi Department of Health in the United Arab Emirates funded this study.

This investigation was considered a public health response by the Abu Dhabi Department of Health and CDC.

About the Author

Dr. Khudair is a senior officer in the Communicable Disease Control and Management Section of the Abu Dhabi Department of Health. His research interests include MERS-CoV and tuberculosis. Dr. Killerby is an epidemiologist in the Division of Viral Diseases, National Center for Immunization and Respiratory Diseases, CDC. Her research interests include respiratory viruses such as MERS-CoV, human coronaviruses, and adenoviruses.

References

- Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus ADME, Fouchier RAM. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med*. 2012;367:1814–20. <http://dx.doi.org/10.1056/NEJMoa1211721>
- World Health Organization. Middle East respiratory syndrome coronavirus (MERS-CoV) [cited 2019 Apr 3]. <http://www.who.int/emergencies/mers-cov>
- World Health Organization. WHO MERS-CoV global summary and assessment of risk, 21 July 2017 [cited 2018 Aug 7]. <https://www.who.int/emergencies/mers-cov/risk-assessment-july-2017.pdf>
- Mohd HA, Al-Tawfiq JA, Memish ZA. Middle East respiratory syndrome coronavirus (MERS-CoV) origin and animal reservoir. *Virology*. 2016;13:87. <http://dx.doi.org/10.1186/s12985-016-0544-0>
- Omrani AS, Al-Tawfiq JA, Memish ZA. Middle East respiratory syndrome coronavirus (MERS-CoV): animal to human interaction. *Pathog Glob Health*. 2015;109:354–62. <http://dx.doi.org/10.1080/20477724.2015.1122852>
- Miguel E, Chevalier V, Ayelet G, Ben Bencheikh MN, Boussini H, Chu DK, et al. Risk factors for MERS coronavirus infection in dromedary camels in Burkina Faso, Ethiopia, and Morocco, 2015. *Euro Surveill*. 2017;22:30498. <http://dx.doi.org/10.2807/1560-7917.ES.2017.22.13.30498>
- Chu DKW, Hui KPY, Perera RAPM, Miguel E, Niemeyer D, Zhao J, et al. MERS coronaviruses from camels in Africa exhibit region-dependent genetic diversity. *Proc Natl Acad Sci U S A*. 2018;115:3144–9. <http://dx.doi.org/10.1073/pnas.1718769115>
- Alagaili AN, Briese T, Mishra N, Kapoor V, Sameroff SC, Burbelo PD, et al. Middle East respiratory syndrome coronavirus infection in dromedary camels in Saudi Arabia. *MBio*. 2014;5:e00884–14. <http://dx.doi.org/10.1128/mBio.01002-14>
- Yusof MF, Eltahir YM, Serhan WS, Hashem FM, Elsayed EA, Marzoug BA, et al. Prevalence of Middle East respiratory syndrome coronavirus (MERS-CoV) in dromedary camels in Abu Dhabi Emirate, United Arab Emirates. *Virus Genes*. 2015;50:509–13. <http://dx.doi.org/10.1007/s11262-015-1174-0>
- Haagmans BL, Al Dhahiry SHS, Reusken CBEM, Raj VS, Galiano M, Myers R, et al. Middle East respiratory syndrome coronavirus in dromedary camels: an outbreak investigation. *Lancet Infect Dis*. 2014;14:140–5. [http://dx.doi.org/10.1016/S1473-3099\(13\)70690-X](http://dx.doi.org/10.1016/S1473-3099(13)70690-X)
- Chu DK, Oladipo JO, Perera RA, Kuranga SA, Chan SM, Poon LL, et al. Middle East respiratory syndrome coronavirus (MERS-CoV) in dromedary camels in Nigeria, 2015. *Euro Surveill*. 2015;20:30086. <http://dx.doi.org/10.2807/1560-7917.ES.2015.20.49.30086>
- Meyer B, Müller MA, Corman VM, Reusken CB, Ritz D, Godeke GJ, et al. Antibodies against MERS coronavirus in dromedary camels, United Arab Emirates, 2003 and 2013. *Emerg Infect Dis*. 2014;20:552–9. <http://dx.doi.org/10.3201/eid2004.131746>
- Reusken CB, Messadi L, Feyisa A, Ularanu H, Godeke GJ, Danmarwa A, et al. Geographic distribution of MERS coronavirus among dromedary camels, Africa. *Emerg Infect Dis*. 2014;20:1370–4. <http://dx.doi.org/10.3201/eid2008.140590>
- Hemida MG, Chu DK, Poon LL, Perera RA, Alhammedi MA, Ng HY, et al. MERS coronavirus in dromedary camel herd, Saudi Arabia. *Emerg Infect Dis*. 2014;20:1231–4. <http://dx.doi.org/10.3201/eid2007.140571>
- Reusken CB, Farag EA, Jonges M, Godeke GJ, El-Sayed AM, Pas SD, et al. Middle East respiratory syndrome coronavirus (MERS-CoV) RNA and neutralising antibodies in milk collected according to local customs from dromedary camels, Qatar, April 2014. *Euro Surveill*. 2014;19:20829. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.23.20829>
- Memish ZA, Cotten M, Meyer B, Watson SJ, Alshahfi AJ, Al Rabeeah AA, et al. Human infection with MERS coronavirus after exposure to infected camels, Saudi Arabia, 2013. *Emerg Infect Dis*. 2014;20:1012–5. <http://dx.doi.org/10.3201/eid2006.140402>
- Farag EA, Reusken CB, Haagmans BL, Mohran KA, Stalin Raj V, Pas SD, et al. High proportion of MERS-CoV shedding dromedaries at slaughterhouse with a potential epidemiological link to human cases, Qatar 2014. *Infect Ecol Epidemiol*. 2015;5:28305. <http://dx.doi.org/10.3402/iee.v5.28305>
- Paden C, Yusof M, Al Hammadi Z, Queen K, Tao Y, Eltahir Y, et al. Zoonotic origin and transmission of Middle East respiratory syndrome coronavirus in the UAE. *Zoonoses Public Health*. 2017.
- Alraddadi BM, Watson JT, Almarashi A, Abedi GR, Turkistani A, Sadran M, et al. Risk factors for primary Middle East Respiratory syndrome coronavirus illness in humans, Saudi Arabia, 2014. *Emerg Infect Dis*. 2016;22:49–55. <http://dx.doi.org/10.3201/eid2201.151340>

20. Müller MA, Meyer B, Corman VM, Al-Masri M, Turkestani A, Ritz D, et al. Presence of Middle East respiratory syndrome coronavirus antibodies in Saudi Arabia: a nationwide, cross-sectional, serological study. *Lancet Infect Dis*. 2015;15:559–64. [http://dx.doi.org/10.1016/S1473-3099\(15\)70090-3](http://dx.doi.org/10.1016/S1473-3099(15)70090-3)
21. Reusken CB, Farag EA, Haagmans BL, Mohran KA, Godeke GJ V, Raj S, et al. Occupational exposure to dromedaries and risk for MERS-CoV infection, Qatar, 2013–2014. *Emerg Infect Dis*. 2015;21:1422–5. <http://dx.doi.org/10.3201/eid2108.150481>
22. Alshukairi AN, Zheng J, Zhao J, Nehdi A, Baharoon SA, Layqah L, et al. High prevalence of MERS-CoV infection in camel workers in Saudi Arabia. *MBio*. 2018;9:e01985–18. <http://dx.doi.org/10.1128/mBio.01985-18>
23. Inovio Pharmaceuticals I. Inovio's MERS vaccine generates high levels of antibodies and induces broad-based T Cell responses in phase 1 study [cited 2018 Aug 24]. <http://ir.inovio.com/news-and-media/news/press-release-details/2018/Inovios-MERS-Vaccine-Generates-High-Levels-of-Antibodies-and-Induces-Broad-based-T-Cell-Responses-in-Phase-1-Study/default.aspx>
24. Sample I. Man in Germany dies of complications stemming from Mers virus. 2015 Jun 16 [cited 2018 Aug 7]. <https://www.theguardian.com/science/2015/jun/16/mers-virus-man-dies-germany>
25. Yusof MF, Queen K, Eltahir YM, Paden CR, Al Hammadi ZMAH, Tao Y, et al. Diversity of Middle East respiratory syndrome coronaviruses in 109 dromedary camels based on full-genome sequencing, Abu Dhabi, United Arab Emirates. *Emerg Microbes Infect*. 2017;6:e101. <http://dx.doi.org/10.1038/emi.2017.89>
26. Trivedi S, Miao C, Al-Abdallat MM, Haddadin A, Alqasrawi S, Iblan I, et al. Inclusion of MERS-spike protein ELISA in algorithm to determine serologic evidence of MERS-CoV infection. *J Med Virol*. 2018;90:367–71. <http://dx.doi.org/10.1002/jmv.24948>
27. Zhao J, Alshukairi AN, Baharoon SA, Ahmed WA, Bokhari AA, Nehdi AM, et al. Recovery from the Middle East respiratory syndrome is associated with antibody and T-cell responses. *Sci Immunol*. 2017;2:eaa5393. <http://dx.doi.org/10.1126/sciimmunol.aan5393>
28. Assiri A, Al-Tawfiq JA, Al-Rabeeah AA, Al-Rabiah FA, Al-Hajjar S, Al-Barrak A, et al. Epidemiological, demographic, and clinical characteristics of 47 cases of Middle East respiratory syndrome coronavirus disease from Saudi Arabia: a descriptive study. *Lancet Infect Dis*. 2013;13:752–61. [http://dx.doi.org/10.1016/S1473-3099\(13\)70204-4](http://dx.doi.org/10.1016/S1473-3099(13)70204-4)
29. Choi WS, Kang CI, Kim Y, Choi JP, Joh JS, Shin HS, et al.; Korean Society of Infectious Diseases. Clinical presentation and outcomes of Middle East respiratory syndrome in the Republic of Korea. *Infect Chemother*. 2016;48:118–26. <http://dx.doi.org/10.3947/ic.2016.48.2.118>
30. Wernery U, Corman VM, Wong EY, Tsang AK, Muth D, Lau SK, et al. Acute Middle East respiratory syndrome coronavirus infection in livestock dromedaries, Dubai, 2014. *Emerg Infect Dis*. 2015;21:1019–22. <http://dx.doi.org/10.3201/eid2106.150038>
31. Sikkema RS, Farag EABA, Himatt S, Ibrahim AK, Al-Romaihi H, Al-Marri SA, et al. Risk factors for primary Middle East respiratory syndrome coronavirus infection in camel workers in Qatar during 2013–2014: a case-control study. *J Infect Dis*. 2017;215:1702–5. <http://dx.doi.org/10.1093/infdis/jix174>
32. Choe PG, Perera RAPM, Park WB, Song K-H, Bang JH, Kim ES, et al. MERS-CoV antibody responses 1 year after symptom onset, South Korea, 2015. *Emerg Infect Dis*. 2017;23:1079–84. <http://dx.doi.org/10.3201/eid2307.170310>

Address for correspondence: Marie E. Killerby, Centers for Disease Control and Prevention, 1600 Clifton Road NE, Mailstop H24-5, Atlanta, GA 30329-4027, USA; email: lxo9@cdc.gov

EID Podcast: Unraveling the Mysteries of Middle East Respiratory Syndrome Coronavirus

Middle East respiratory syndrome coronavirus (MERS-CoV) is a novel CoV known to cause severe acute respiratory illness in humans; approximately 40% of confirmed cases have been fatal. Human-to-human transmission and multiple outbreaks of respiratory illness have been attributed to MERS-CoV, and severe respiratory illness caused by this virus continues to be identified. As of February 23, 2014, the World Health Organization has reported 182 laboratory-confirmed cases of MERS-CoV infection, including 79 deaths, indicating an ongoing risk for transmission to humans in the Arabian Peninsula.



Visit our website to listen:
[http://www2c.cdc.gov/podcasts/
player.asp?f=8631627](http://www2c.cdc.gov/podcasts/player.asp?f=8631627)

EMERGING INFECTIOUS DISEASES

Risk Factors for MERS-CoV Seropositivity among Animal Market and Slaughterhouse Workers, Abu Dhabi, United Arab Emirates, 2014–2017

Appendix 1

Market and Slaughterhouse Worker Surveys

Market Worker Survey

Form Serial Number: ER/BW/_____/16
Interview Date: ____/____/____ Day Month Year
Interviewer initials: ____

Name of the animal market _____	
Location of the animal market (region and city) _____	
<u>Subject Information</u>	
FIRST NAME _____ LAST NAME _____	
AGE (years)	<i>GENDER</i> <input type="checkbox"/> <i>Male</i> <input type="checkbox"/> <i>Female</i>
D.O.B: dd/mm/yyyy	<i>NATIONALITY</i> _____

CONSENTED

YES **NO**

BLOOD SPECIMEN COLLECTED

YES **NO**

Specimen collection date:

____/____/____
Day Month Year

Investigator Notes:

Sample Label:

(Place sticker here)

Home phone _____ Mobile phone 1 _____

Mobile phone 2 _____

CURRENT HOME ADDRESS

Street _____

City _____ Area: _____

INTERNAL USE:

Data Entered:
Initials:

Data Entry Checked:
Initials:

OCCUPATION

What is your job title at the animal market? _____

Do you have animals at your home? **YES** **NO**

If YES, specify type and number of animals at home: _____

How long have you worked at this animal market? _____years _____months

If less than one year, have you worked at any other animal markets prior to this one?

YES (specify location, number, or name of the animal market) _____

NO

How many hours per week do you typically work at the animal market? _____hours/week

How many hours per week do you typically work with animals outside the animal market?
_____ hours/week

Do you currently hold other jobs aside from working at the animal market? **YES** **NO**

If YES, Please specify location and job title of other job(s): _____

How often do you perform each of the following?

- | | | | |
|--|---------------------------------------|---|---------------------------------------|
| Handle camels | <input type="checkbox"/> NEVER | <input type="checkbox"/> SOMETIMES | <input type="checkbox"/> DAILY |
| Feed camels | <input type="checkbox"/> NEVER | <input type="checkbox"/> SOMETIMES | <input type="checkbox"/> DAILY |
| Clean camels | <input type="checkbox"/> NEVER | <input type="checkbox"/> SOMETIMES | <input type="checkbox"/> DAILY |
| Clean camel housing | <input type="checkbox"/> NEVER | <input type="checkbox"/> SOMETIMES | <input type="checkbox"/> DAILY |
| Handle or dispose camel waste | <input type="checkbox"/> NEVER | <input type="checkbox"/> SOMETIMES | <input type="checkbox"/> DAILY |
| Clean equipment | <input type="checkbox"/> NEVER | <input type="checkbox"/> SOMETIMES | <input type="checkbox"/> DAILY |
| Milk camels | <input type="checkbox"/> NEVER | <input type="checkbox"/> SOMETIMES | <input type="checkbox"/> DAILY |
| Slaughter camels | <input type="checkbox"/> NEVER | <input type="checkbox"/> SOMETIMES | <input type="checkbox"/> DAILY |
| Assist with camel parturition (birthing) | <input type="checkbox"/> NEVER | <input type="checkbox"/> SOMETIMES | <input type="checkbox"/> DAILY |
| Give medications/vaccines to camels | <input type="checkbox"/> NEVER | <input type="checkbox"/> SOMETIMES | <input type="checkbox"/> DAILY |

Other tasks (specify and indicate frequency):

ANIMAL EXPOSURES

In the past 12 months, have you had direct physical contact with any animals or their waste?

- Camels Never Rarely Monthly Weekly DAILY
- Goats Never Rarely Monthly Weekly DAILY
- Sheep Never Rarely Monthly Weekly DAILY
- Cattle Never Rarely Monthly Weekly DAILY
- Horses Never Rarely Monthly Weekly DAILY
- Dogs Never Rarely Monthly Weekly DAILY
- Cats Never Rarely Monthly Weekly DAILY
- Bats Never Rarely Monthly Weekly DAILY
- Rodents Never Rarely Monthly Weekly DAILY
- Birds Never Rarely Monthly Weekly DAILY

Other animals (specify and indicate frequency):

Were any of the animals you had contact with ill? YES NO Do not recall

If YES, please describe the following:

Which animals were ill? _____

What were their symptoms? _____

Did you develop any illness after this contact with the ill animal? YES NO

If YES, what were your symptoms? _____

AND how long after this contact did your symptoms begin?

- <1 Week 1-2 Weeks 3-4 Weeks >1 Month

In the past 12 months, have you eaten any raw camel milk or milk products?

- YES NO Do not recall

In the past 12 months, have you drunk camel urine?

- YES NO Do not recall

In the past 12 months, have you eaten uncooked camel meat?

- YES NO Do not recall

PPE and HYGIENE

Do you usually wear personal protective equipment (PPE) when handling animals or their waste?

YES NO

If YES, check all that apply:

Gloves Dust Mask Respirator Coveralls Boots or boot covers

Eye protection (goggles, safety glasses) Other (specify):

Do you take your work clothes home with you? YES NO

Where are your work clothes washed? Home Work place Other (specify)

Who washes your work clothes? Self Household member Other worker

At which of the following times do you usually wash your hands at the animal market (check all that apply)?

Before and after each animal-related task

At meal times

At bathroom times

At prayer times

At the beginning and end of the day

Where do you wash your hands at the animal market? _____

TRAVEL HISTORY

During the last 12 months, did you visit: YES NO do not recall

If yes, check all that apply:

Bahrain Palestinian Territories Lebanon

Iraq Qatar Yemen

Iran Saudi Arabia Oman

Israel Syria Jordan

Kuwait South Korea Other

(list) _____

Did you travel within United Arab Emirates? YES NO

If YES, where? _____

ILLNESS HISTORY

During the last 12 months, have you seen a doctor for a respiratory illness? (for symptoms including any of the following: cough, fever, runny nose, shortness of breath, rapid or shallow breathing, sore throat, vomiting after cough, and wheezing):

YES **NO** **Do not recall**

If YES...

Where did you seek care for this illness (name/location of medical institution)?

Can you describe your symptoms during this time?

What medications did you take during this time?

Was a chest x-ray performed?

YES **NO** **Do not recall**

If YES,

Health insurance card number _____

Medical record number _____

Do you have any underlying medical conditions (e.g. immunocompromised, heart disease, pulmonary disease (COPD, emphysema, cancer, diabetes, hypertension, smoking, etc.)?)

YES **NO**

If YES, please describe _____

HUMAN EXPOSURES

Did you have contact with anyone hospitalized for respiratory illness during the last 12 months?

YES **NO** **Do not recall**

ADDITIONAL NOTES:

Slaughterhouse Worker Survey

Form Serial Number: ER/BW/_____/16
Interview Date: ____/____/____
Day Month Year
Interviewer initials: _____

Name of the slaughter house _____

Location of the slaughter house (region and city) _____

Subject Information

FIRST NAME _____ LAST NAME _____

AGE (years)

D.O.B: dd/mm/yyyy

GENDER

Male

Female

NATIONALITY _____

CONSENTED

YES

NO

BLOOD SPECIMEN COLLECTED

YES

NO

Specimen collection date:

____/____/____
Day Month Year

Investigator Notes:

Sample Label:

(Place sticker here)

Home phone _____ Mobile phone 1 _____

Mobile phone 2 _____

CURRENT HOME ADDRESS

Street _____

City _____ Area: _____

INTERNAL USE:

**Data Entered:
Initials:**

**Data Entry Checked:
Initials:**

ANIMAL EXPOSURES

In the past 12 months, have you had direct physical contact with any animals or their waste?

- Camels Never Rarely Monthly Weekly DAILY
- Goats Never Rarely Monthly Weekly DAILY
- Sheep Never Rarely Monthly Weekly DAILY
- Cattle Never Rarely Monthly Weekly DAILY
- Horses Never Rarely Monthly Weekly DAILY
- Dogs Never Rarely Monthly Weekly DAILY
- Cats Never Rarely Monthly Weekly DAILY
- Bats Never Rarely Monthly Weekly DAILY
- Rodents Never Rarely Monthly Weekly DAILY
- Birds Never Rarely Monthly Weekly DAILY

Other animals (specify and indicate frequency):

Were any of the animals you had contact with ill? YES NO Do not recall

If YES, please describe the following:

Which animals were ill? _____

What were their symptoms? _____

Did you develop any illness after this contact with the ill animal? YES NO

If YES, what were your symptoms? _____

AND how long after this contact did your symptoms begin?

<1 Week 1-2 Weeks 3-4 Weeks >1 Month

In the past 12 months, have you eaten any raw camel milk or milk products?

YES NO Do not recall

In the past 12 months, have you drunk camel urine?

YES NO Do not recall

In the past 12 months, have you eaten uncooked camel meat?

YES NO Do not recall

PPE and HYGIENE

Do you usually wear personal protective equipment (PPE) when handling animals or their waste?

YES NO

If YES, check all that apply:

Gloves Dust Mask Respirator Coveralls Boots or boot covers

Eye protection (goggles, safety glasses) Other (specify):

Do you take your work clothes home with you? YES NO

Where are your work clothes washed? Home Work place Other (specify)

Who washes your work clothes? Self Household member Other worker

At which of the following times do you usually wash your hands at the slaughter house (check all that apply)?

Before and after each animal-related task

At meal times

At bathroom times

At prayer times

At the beginning and end of the day

Where do you wash your hands at the slaughter house? _____

TRAVEL HISTORY

During the last 12 months, did you visit: YES NO do not recall

If yes, check all that apply:

Bahrain Palestinian Territories Lebanon

Iraq Qatar Yemen

Iran Saudi Arabia Oman

Israel Syria Jordan

Kuwait South Korea Other

(list) _____

Did you travel within United Arab Emirates? YES NO

If YES, where? _____

ILLNESS HISTORY

During the last 12 months, have you seen a doctor for a respiratory illness? (for symptoms including any of the following: cough, fever, runny nose, shortness of breath, rapid or shallow breathing, sore throat, vomiting after cough, and wheezing):

YES **NO** **Do not recall**

If YES...

Where did you seek care for this illness (name/location of medical institution)?

Can you describe your symptoms during this time?

What medications did you take during this time?

Was a chest x-ray performed?

YES **NO** **Do not recall**

If YES,

Health insurance card number _____

Medical record number _____

Do you have any underlying medical conditions (e.g. immunocompromised, heart disease, pulmonary disease (COPD, emphysema, cancer, diabetes, hypertension, smoking, etc.)?)

YES **NO**

If YES, please describe _____

HUMAN EXPOSURES

Did you have contact with anyone hospitalized for respiratory illness during the last 12 months?

YES **NO** **Do not recall**

ADDITIONAL NOTES:

Risk Factors for MERS-CoV Seropositivity among Animal Market and Slaughterhouse Workers, Abu Dhabi, United Arab Emirates, 2014–2017

Appendix 2

Appendix 2 Table. Results of N ELISA, S ELISA, and microneutralization tests for all participants who ever had a seropositive test result*

Participant ID	Sample number	Year of collection	Sampling round	OD N ELISA at 1:400 cutoff 0.29	OD S ELISA at 1:400 cutoff 0.2	MNT (cutoff 20)	Interpretation†
1	1st	2014	1	1.57	0.32	40	Positive
2	1st	2014	1	1.54	0.26	20	Positive
3	1st	2014	1	0.64	0.33	80	Positive
4	1st	2014	1	0.79	0.04	20	Positive
5	1st	2014	1	1.12	0.19	20	Positive
6	1st	2014	1	1.11	0.31	80	Positive
7	1st	2015	2	0.21	0.18	40	Positive
8	1st	2014	1	0.22	0.01	<20	Negative
5	2nd	2015	2	0.35	0.14	20	Positive
6	2nd	2015	2	0.40	0.24	40	Positive
3	2nd	2015	2	0.22	0.34	40	Positive
8	1st	2015	2	1.16	0.37	40	Positive
8	1st	2015	2	0.45	0.20	20	Positive
9	1st	2015	2	0.78	1.37	80	Positive
10	1st	2015	2	0.15	0.48	80	Positive
11	1st	2015	2	0.15	0.33	40	Positive
12	1st	2015	2	0.83	0.28	20	Positive
13	1st	2015	2	0.86	0.20	40	Positive
14	1st	2015	2	0.03	0.68	20	Positive
15	1st	2015	2	0.48	0.26	40	Positive
16	1st	2015	2	0.76	0.37	40	Positive
17	1st	2015	2	0.38	0.86	20	Positive
18	1st	2015	2	0.40	0.28	160	Positive
19	1st	2015	2	0.48	0.94	80	Positive
20	1st	2015	2	0.56	0.56	80	Positive
21	1st	2015	2	0.58	0.54	40	Positive
22	1st	2015	2	0.34	0.54	40	Positive
23	1st	2015	2	0.13	0.32	40	Positive
24	1st	2015	2	0.44	0.16	20	Positive
25	1st	2015	2	0.69	0.35	20	Positive
26	1st	2015	2	0.72	0.45	40	Positive
27	1st	2015	2	0.03	0.92	320	Positive
28	1st	2015	2	0.65	0.22	20	Positive
29	1st	2015	2	0.49	0.67	40	Positive
30	1st	2015	2	0.64	0.28	80	Positive
2	2nd	2015	2	1.33	0.62	<20	Positive
5	3rd	2016	3	0.26	0.09	<20	Indeterminate‡
7	2nd	2016	3	0.09	0.10	20	Positive
31	1st	2016	3	1.01	0.36	160	Positive
32	1st	2016	3	0.13	0.80	20	Positive
33	1st	2016	3	0.29	0.87	40	Positive
34	1st	2016	3	0.07	0.64	40	Positive
35	1st	2016	3	0.43	1.14	40	Positive
36	1st	2016	3	0.04	0.34	20	Positive
37	1st	2016	3	0.02	0.88	80	Positive
38	1st	2016	3	0.19	0.85	40	Positive
39	1st	2016	3	0.39	0.93	20	Positive
40	1st	2016	3	0.05	1.39	20	Positive
2	3rd	2016	3	0.36	0.94	20	Positive

Participant ID	Sample number	Year of collection	Sampling round	OD N ELISA at 1:400 cutoff 0.29	OD S ELISA at 1:400 cutoff 0.2	MNT (cutoff 20)	Interpretation†
25	2nd	2016	3	0.00	0.04	80	Positive
21	2nd	2016	3	0.25	0.59	20	Positive
41	1st	2016	3	0.52	0.48	<20	Positive
4	2nd	2016	3	0.01	0.02	<20	Negative
42	1st	2016	3	0.80	0.76	<20	Positive
43	1st	2017	3	1.33	0.74	<20	Positive
44	1st	2017	3	0.55	0.33	<20	Positive
45	1st	2017	3	0.02	0.34	20	Positive
46	1st	2017	3	0.46	0.98	80	Positive
16	2nd	2017	3	0.65	0.79	160	Positive
47	1st	2017	3	0.53	0.60	80	Positive
48	1st	2017	3	1.60	0.70	640	Positive
49	1st	2017	3	0.24	0.53	20	Positive
50	1st	2017	3	0.39	0.46	20	Positive
51	1st	2017	3	0.13	0.27	20	Positive
52	1st	2017	3	0.42	0.46	20	Positive
53	1st	2017	3	0.38	0.46	40	Positive
54	1st	2017	3	0.08	0.43	40	Positive
55	1st	2017	3	0.22	0.28	20	Positive
56	1st	2017	3	0.49	0.72	40	Positive
57	1st	2017	3	0.40	0.75	20	Positive
58	1st	2017	3	0.58	0.87	160	Positive
59	1st	2017	3	0.53	0.33	40	Positive
60	1st	2017	3	0.24	0.27	40	Positive
61	1st	2017	3	0.66	1.02	80	Positive
62	1st	2017	3	0.47	0.79	160	Positive
63	1st	2017	3	0.20	0.51	20	Positive
64	1st	2017	3	0.08	0.60	20	Positive
65	1st	2017	3	0.12	0.34	20	Positive

*Cell shading indicates ELISA results above the cutoff point.

†Samples were considered positive if positive on N and S ELISA, or if positive on microneutralization. Specimens near the limits of detection but not consistently above or below these limits were considered indeterminate. For the epidemiologic analysis, individuals with an indeterminate result were considered seronegative.

‡Participant 5 was considered negative for epidemiologic analysis during sampling round 3, but as this indeterminate second sample result was reported after a positive result one year prior, participant 5 was not considered to have lost seropositive status.