Our report has limitations. Our sample size of asymptomatic cases is small, and follow-up was short. Recall bias of exposure history is another limitation; in the absence of clear symptom onset, asymptomatic persons might be less likely to accurately recall exposures than persons with symptoms. Finally, that the study took a place during the post-peak period of the epidemic in Wuhan, so contacts could have been seropositive already; those tested were seronegative, but most contacts did not have serologic testing.

In conclusion, as the population returns to the workplace, asymptomatic SARS-CoV-2-infected persons could be among workers. Although we did not detect transmission among 41 contacts of persons who were SARS-CoV-2-positive, such transmission cannot be excluded. Therefore, continued testing, self-quarantine, and mask-wearing should be encouraged to reduce the risk for additional outbreaks.

Acknowledgments

We thank all our colleagues for helping us during the current study.

This study was supported by Zhejiang University special scientific research fund for COVID-19 prevention and control, the Huazhong University of Science and Technology (HUST) COVID-19 Rapid Response Call (grant no. 2020kfyXGYJ019).

About the Author

Dr. Han is a clinician in the Department of Radiology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China. Her research interests including image diagnosis of pneumonia and lung cancer.

References

- Qiu J. Covert coronavirus infections could be seeding new outbreaks. Nature. 2020 Mar 20 [Epub ahead of print]. https://doi.org/10.1038/d41586-020-00822-x
- Pan A, Liu L, Wang C, Guo H, Hao X, Wang Q, et al. Association of public health interventions with the epidemiology of the COVID-19 outbreak in Wuhan, China. JAMA. 2020;323:1–9. https://doi.org/10.1001/ jama.2020.6130
- Wuhan Municipal Health Commission. CN-HEALTHCARE: Hubei health, 2020 April 19 [in Chinese] [cited 2020 May 25]. https://www.cn-healthcare.com/articlewm/20200419/ content-1105403.html
- Furukawa NW, Brooks JT, Sobel J. Evidence supporting transmission of severe acute respiratory syndrome coronavirus 2 while presymptomatic or asymptomatic. Emerg Infect Dis. 2020 May 4 [Epub ahead of print]. https://doi.org10.3201/eid2607.201595
- 5. Arons MM, Hatfield KM, Reddy SC, Kimball A, James A, Jacobs JR, et al. Presymptomatic SARS-CoV-2 infections and

transmission in a skilled nursing facility. N Engl J Med. 2020;382:2081–90. https://doi.org/10.1056/ NEJMoa2008457

- Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. N Engl J Med. 2020;382:1177–9. https://doi.org/10.1056/NEJMc2001737
- Kim SE, Jeong HS, Yu Y, Shin SU, Kim S, Oh TH, et al. Viral kinetics of SARS-CoV-2 in asymptomatic carriers and presymptomatic patients. Int J Infect Dis. 2020;95:441–3. https://doi.org/10.1016/j.ijid.2020.04.083
- He D, Zhao S, Lin Q, Zhuang Z, Cao P, Wang MH, et al. The relative transmissibility of asymptomatic COVID-19 infections among close contacts. Int J Infect Dis. 2020;94: 145–7. https://doi.org/10.1016/j.ijid.2020.04.034
- Zhou R, Li F, Chen F, Liu H, Zheng J, Lei C, et al. Viral dynamics in asymptomatic patients with COVID-19. Int J Infect Dis. 2020;96:288–290. https://doi.org/10.1016/ j.ijid.2020.05.03

Address for correspondence: Heshui Shui, Department of Radiology, Union Hospital, Tongji Medical College, Wuhan 430030, China; email: heshuishi@hust.edu.cn

Effects of Proactive Social Distancing on COVID-19 Outbreaks in 58 Cities, China

Zhanwei Du, Xiaoke Xu, Lin Wang, Spencer J. Fox, Benjamin J. Cowling, Alison P. Galvani, and Lauren Ancel Meyers

Author affiliations: The University of Texas at Austin, Austin, Texas, USA (Z. Du, S.J. Fox, L.A. Meyers); Dalian Minzu University, Dalian, China (X. Xu); University of Cambridge, Cambridge, UK (L. Wang); The University of Hong Kong, Hong Kong, China (B.J. Cowling); Center for Infectious Disease Modeling and Analysis, Yale School of Public Health, New Haven, Connecticut, USA (A.P. Galvani); Santa Fe Institute, Santa Fe, New Mexico, USA (L.A. Meyers)

DOI: https://doi.org/10.3201/eid2609.201932

Cities across China implemented stringent social distancing measures in early 2020 to curb coronavirus disease outbreaks. We estimated the speed with which these measures contained transmission in cities. A 1-day delay in implementing social distancing resulted in a containment delay of 2.41 (95% CI 0.97–3.86) days. On December 31, 2019, a cluster of atypical pneumonia in Wuhan, China, was reported to the regional office of the World Health Organization (WHO). Its etiology was later identified as the novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Coronavirus disease (COVID-19) spread rapidly across China and internationally (1); as of April 9, 2020, a total of 1,436,198 confirmed cases and 85,522 deaths had been reported in 209 countries (2). In the absence of pharmaceutical prophylactic options, the primary means of COVID-19 control are social distancing interventions, including school closures, work restrictions, shelter-in-place measures, and travel bans.

In late January, reported COVID-19 cases rose steeply in Hubei Province, and imported cases sparked outbreaks in many other cities throughout China. By February 14, 2020, the government had limited the movement of >500 million persons across 80 cities, many of which rapidly enacted multiple social distancing orders to slow the local spread of the virus, including restricting nonessential services and public transit (3–6). Given the substantial economic and societal costs of such measures (7), estimates of their effectiveness can serve as critical evidence for intervention policy decisions worldwide (8).

Using case data from online reports published by the Chinese Center for Disease Control and health commissions (Appendix Table 4, https://wwwnc.cdc. gov/EID/article/26/9/20-1932-App1.pdf), we estimated the time elapsed between the first reported case in a city and successful containment of the outbreak (χ). Technically, we consider an outbreak contained when the 95% CI of the instantaneous reproduction number (R_t) drops below 1. We analyzed the speed of COVID-19 containment for 58 cities in mainland China outside of Huebei Province that had \geq 20 confirmed cases by February 14, 2020 (Figure; Appendix Tables 2, 3). Collectively, these cities deployed 7 different types of interventions over the course of their epidemics (9): bans on entertainment and public gatherings;



Figure. Coronavirus disease (COVID-19) introductions, transmission, and containment for 2 provincial capitals, China, before February 15, 2020. A) Estimated daily incidence of COVID-19 cases and the implementation of local social distancing measures in Xi'an. B) Estimated daily incidence of COVID-19 cases and the implementation of local social distancing measures in Nanjing. C, D) Estimated daily time-varying reproduction numbers (R_i). Green line indicates the median and gray shading 95% CI for R_i . We calculated the number of days from the first reported imported case until the upper 95% CI drops below 1 (χ) for (C) Xi'an and (D) Nanjing. E) The distribution of χ across 58 cities in mainland China. Mean duration of outbreaks is 21 days (SD ± 7). Based on an area under the curve comparison between gamma, log-normal, and Weibull distributions fitted via maximum-likelihood to the data, we found that the χ values are roughly Weibull distributed with scale 22.94 (95% CI 21.12–24.91) and shape 3.28 (95% CI 2.68–4.02), indicated by black line. F) The distribution of time between the first locally reported case and the first social distancing measure resembles a Weibull distribution with scale 14.24 (95% CI 13.01–15.60) and shape 2.98 (95% CI 2.44–3.65).

broad restrictions on public service including healthcare, schooling, shopping, and restaurants; initiation of a level 1 response entailing systematic testing and isolation of confirmed cases; suspension of intracity public transport; suspension of travel between cities; reporting of confirmed cases; recruitment of governmental staff and volunteers to enforce quarantine and social distancing. The mean $(\pm SD)$ time between the first confirmed case and the implementation of the first social distancing measure was 13 (± 4.7) days. By the time these measures were enacted, the median cumulative reported cases in a city was 40, but the range was 9-248 across the 58 cities. The mean time until successful containment was 21 (+7) days after the first reported case and 8 (+ 6.8) days following the initiation of interventions. During the period of containment, the reproduction number (R,) declined by an average of 54.3% (+17.6%) (Appendix Figure 2).

Using a combination of linear regression and best-subsets model selection (10), we found that the timing of the first intervention and the initiation of level 1 response significantly predicted the speed of containment across the 36 cities that deployed all 7 interventions ($R^2 = 0.27$; p<0.001) (Appendix Figure 1). A delay of 1 day in implementing the first intervention is expected to prolong an outbreak by 2.41 (95% CI 0.96–3.86) days. In contrast, the timing of the level 1 response was inversely related to the speed of containment. Level 1 responses were initiated by the central government across mainland China over the course of 1 week, starting with the hardest hit areas in and near Hubei Province on the first day and working outwards toward more distant cities. Thus, the day of level 1 initiation within this 1-week period is a likely indicator for the initial severity of an outbreak and the corresponding difficulty of containment.

We have estimated the value of proactive social distancing interventions in terms of a reduction in days until successful containment. However, because most cities implemented multiple measures quickly and simultaneously, we are unable to disentangle the efficacies of individual modes of social distancing. We note that our estimates of R, may be biased by the limited case report data available before February 14, 2020; we lack information about testing rates and priorities in China before February 14. As public health agencies around the globe struggle to determine when to implement potentially costly social distancing measures, these estimates highlight the potential long-term benefits of early and decisive action.

Financial support was provided by US National Institutes of Health (grant no. U01 GM087719), the National Natural Science Foundation of China (grant no. 61773091), and Tito's Handmade Vodka.

About the Author

Dr. Du is a postdoctoral researcher in the Department of Integrative Biology at the University of Texas at Austin. He develops mathematical models to elucidate the transmission dynamics, surveillance, and control of infectious diseases.

References

- Chen S, Yang J, Yang W, Wang C, Bärnighausen T. COVID-19 control in China during mass population movements at New Year. Lancet. 2020;395:764–6. https://doi.org/10.1016/S0140-6736(20)30421-9
- World Health Organization. Coronavirus disease 2019 ([COVID-19)]: situation report 80. 2020 [cited 2020 Apr 9]. https://www.who.int/docs/default-source/coronaviruse/ situation-reports/20200409-sitrep-80-covid-19.pdf
- Chan JF-W, Yuan S, Kok K-H, To KK-W, Chu H, Yang J, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. Lancet. 2020;395:514– 23. https://doi.org/10.1016/S0140-6736(20)30154-9
- Kraemer MUG, Yang C-H, Gutierrez B, Wu C-H, Klein B, Pigott DM, et al. The effect of human mobility and control measures on the COVID-19 epidemic in China. Science. 2020 Mar 25 [cited 2020 Mar 26]. https://science.sciencemag.org/ content/early/2020/03/25/science.abb4218
- Chinazzi M, Davis JT, Ajelli M, Gioannini C, Litvinova M, Merler S, et al. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. Science. 2020;368:395–400. https://doi.org/10.1126/science.aba9757
- Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72,314 cases from the Chinese Center for Disease Control and Prevention. JAMA. 2020;323:1239–42. https://doi.org/10.1001/jama.2020.2648
- Ayittey FK, Ayittey MK, Chiwero NB, Kamasah JS, Dzuvor C. Economic impacts of Wuhan 2019-nCoV on China and the world. J Med Virol. 2020;92:473–5. https://doi.org/10.1002/ jmv.25706
- Leung K, Wu JT, Liu D, Leung GM. First-wave COVID-19 transmissibility and severity in China outside Hubei after control measures, and second-wave scenario planning: a modelling impact assessment. Lancet. 2020;395:1382–93. https://doi.org/10.1016/S0140-6736(20)30746-7
- Tian H, Liu Y, Li Y, Wu C-H, Chen B, Kraemer MUG, et al. An investigation of transmission control measures during the first 50 days of the COVID-19 epidemic in China. Science. 2020;368:638–42. https://doi.org/10.1126/science.abb6105
- Yang H. The case for being automatic: introducing the automatic linear modeling (LINEAR) procedure in SPSS statistics. Multiple Linear Regression Viewpoints. 2013;39:27–37.

Address for correspondence: Lauren Ancel Meyers, Department of Integrative Biology, The University of Texas at Austin, 1 University Station C0990, Austin, TX 78712, USA; email: laurenmeyers@austin.utexas.edu

Acknowledgments

We thank Simon Cauchemez for helpful discussions.

Article DOI: https://doi.org/10.3201/eid2609.201932

Effects of Proactive Social Distancing on COVID-19 Outbreaks in 58 Cities, China

Appendix

Supplemental Methods

We collected data from online reports published by China Center for Disease Control and Prevention and health commissions (Appendix Table 4). The data comprised 8,410 confirmed cases before February 15, 2020, across 271 cities in mainland China with known dates of symptom onset and classified as imported or locally infected. In any case, with missing symptom onset timings, we allocate a random value post the official announcement following the gamma distribution of the period from the onset of symptoms (*1*). We also aggregated data on the timing of 7 different classes of social distancing measures in 58 cities outside Hubei province (Appendix Table 1; Appendix 2).

Estimation of Rt

We estimate the time-varying reproduction numbers for the outbreaks in 58 cities using the R package EpiEstim (2) based on the method of Ref (3). Given time series data for incident imported and locally-infected cases and the distribution of serial intervals, the algorithm produces the time series of R_t with medians and 95% CI. We assume the serial interval follows the gamma distribution (mean 5.11 days, SD 2.68 days (1) and the length of sliding time window for an estimated 7 days (4).

Regression Analysis

To assess the impact of intervention type and timing on the speed of containment, we applied a simple regression and variable selection method (Appendix Table 2). Specifically, we fit a linear regression model predicting the time until containment (i.e., days between symptom onset for the first reported case and the estimated 95% CI upper bound of R_t dropping below 1) as given by

$$\chi = \sum_{i=1}^{8} \beta_i T_i + \epsilon$$

where β_i is the coefficient of the *i*th variable T_i and ϵ is the intercept.

We analyze data from a subset consisting of 36 of 58 cities in mainland China that implemented all 8 interventions and contained the local outbreak by February 14, 2020. We used the best-subsets regression to identify the best fit model for all models containing k, in which k range is 1–8, and chose our final model based on the Akaike information criterion corrected for small sample sizes (5).

| | | Candidate model |
|-------------------------------|---|-----------------|
| Symbol | Definition | predictor |
| t ₀ | Day of symptom onset for the first reported case | No |
| <i>t</i> e | Day on which the earliest social distancing measure(s) was enacted | No |
| t _{control_95} | Day that the upper 95% CI bound of R_t drops below one (without rebounding) | No |
| X | Containment period: the number of days between symptom onset of the first reported case (t_0) and the upper 95% CI of R_t dropping below 1 ($t_{control_{95}}$) | No |
| T _{SD} † | Days between first reported case (t_0) and implementation of the first social distancing measure implemented (t_e) | Yes |
| $\mathcal{T}_{entertainment}$ | Days between the first reported case (t_0) and ban on entertainment and public gatherings (e.g., bar, café, cinema) | Yes |
| T _{service} | Days between the first reported case (t_0) and restrictions on public services, including hospitals, schools, stores, and restaurants | Yes |
| T _{level-1} ‡ | Days between the first reported case (t_0) and the initiation of urban level-1 response for systematic testing and isolation of confirmed cases | Yes |
| $\mathcal{T}_{intra_trans}$ | Days between the first reported case (t_0) and suspension of intracity public transport (bus and subway) | Yes |
| T _{inter_trans} | Days between the first reported case (t_0) and suspension of inbound and outbound travel (i.e., intercity rail, highway, and air travel) | Yes |
| T _{report} | Days between the first reported case (t_0) and online posting of confirmed case reports | Yes |
| T _{assist} | Days between the first reported case (t ₀) and the recruitment of governmental staff and volunteers to assist with guarantine and social distancing | Yes |

Appendix Table 1. Variable definitions for analysis of the mitigating effect of social distancing on coronavirus spread in cities in China, 2020

Appendix Table 2. Results of linear regression relating immediacy of interventions to speed of containment, China*

| | <u> </u> | | | | | |
|---|----------------------|---------|--|--|--|--|
| Predictors | Coefficient | p value | | | | |
| Intercept | 14.37 (9.02, 19.72) | 0.000 | | | | |
| T _{SD} † | 2.41 (0.97, 3.86) | 0.002 | | | | |
| T _{level-1} | -1.87 (-3.14, -0.60) | 0.005 | | | | |
| *Speed of containment is defined as days until upper 95% CI of $R_t < 1$. We included in the regression results for 36 | | | | | | |
| of 58 cities that implemented all 8 intervention measures and had sufficient data for prediction available (λ) by | | | | | | |
| February 14, 2020. We selected the parameters in the table from among all of the candidates in Appendix Table 1 | | | | | | |
| using the best-subsets method (6) that identifies the most informative combinations of predictors with respect to the | | | | | | |
| Hurvich and Tsai's Information Criterion (7). The fitted model has $R^2 = 0.27$, p<0.001. | | | | | | |
| \dagger Days between first reported case (t_0) and implementation of the first social distancing measure implemented (t_e). | | | | | | |
| $\pm D$ and the provided the second second the second secon | | | | | | |

the provided first reported case (t_0) and implementation of the first social distancing measure implemented (t_e). ‡Days between the first reported case (t_0) and the initiation of urban level-1 response for systematic testing and isolation of confirmed cases.

Appendix Table 3. Distribution fits for the time between the first reported case and containment (λ) and time between the first reported case and the implementation of the first social distancing measure*

| | | | | Akaike's Information |
|-----------------|--------------|---------------------------------------|------------------------|----------------------|
| Time | Distribution | Shape (95% CI) | Scale (95% CI) | Criterion |
| χ | Gamma | 7.63 (5.343,10.895) | 2.694 (1.864,3.893) | 394.164 |
| | Lognormal | 2.956 (2.854,3.058) | 0.389 (0.329,0.476) | 398.975 |
| | Weibull | 3.282 (2.682,4.016) | 22.937 (21.119,24.911) | 390.329 |
| T _{SD} | Gamma | 6.121 (4.294,8.725) | 2.079 (1.437,3.008) | 349.986 |
| | Lognormal | 2.46 (2.342,2.577) | 0.448 (0.379,0.549) | 357.869 |
| | Weibull | 2.981 (2.436,3.649) | 14.244 (13.008,15.597) | 344.642 |
| *D 1 | | · · · · · · · · · · · · · · · · · · · | | |

*Data were taken from 58 cities in mainland China before February 15, 2020. Timing is calculated in terms of days from symptom onset of the first reported case in the city. Containment is defined by the first day that the estimated upper 95% CI for Rt permanently drops below 1.

| | | | | | | | | | | | T0 since |
|----------------------|------------------|---------|---------------|------------|------------|----------|----------|--------|----------|---------|----------|
| City name | City name | | | . . | | Intra_ | Inter_ | | | | Jan. 1, |
| (English) | (Chinese) | Chi | Entertainment | Service | Level-1 | trans | trans | Report | Assist | SD | 2020 |
| Sanya | <u> </u> | 11 | 17 | 21 | 17 | 18 | 21 | 14 | 25 | 14 | 8 |
| Zhongshan | 中山 | 8 | 13 | 13 | 15 | 16 | 18 | 23 | 28 | 13 | 11 |
| Linyi | 临沂 | 20 | 17 | 17 | 16 | ~~ | 18 | 15 | 24 | 15 | 8 |
| Jiujiang | 九江 | 37 | 23 | 23 | 20 | 23 | 26 | 20 | 29 | 20 | 3 |
| Xinyang | 信阳 | 24 | 21 | 21 | 23 | 23 | 30 | 23 | 31 | 21 | 3 |
| North Sea | 北海 | 16 | 14 | | 16 | 23 | 23 | 12 | 24 | 12 | 9 |
| Nanjing | 南京 | 20 | 11 | 11 | 12 | | 18 | 10 | 30 | 10 | 12 |
| Nanning | 南宁 | 30 | | | 10 | | | 11 | 23 | 10 | 15 |
| Nanchang | 南昌 | 25 | 15 | 15 | 11 | | 15 | 19 | 29 | 11 | 12 |
| Nanyang | 南阳 | 22 | 20 | 20 | 22 | 22 | 22 | 30 | 30 | 20 | 4 |
| Hefei | 合肥 | 20 | 14 | 25 | 13 | 15 | 17 | 18 | 24 | 13 | 10 |
| Harbin | 哈 尔滨 | 21 | 21 | 14 | | 22 | | 22 | | 14 | 13 |
| Shangqiu | 商丘 | 35 | 22 | 22 | | 26 | 26 | | | 22 | 2 |
| Tianjin | 天津 | 28 | | 18 | 15 | 23 | 18 | 13 | 24 | 13 | 9 |
| Weihai | 威海 | 15 | 17 | 17 | 9 | 15 | 12 | 9 | 30 | 9 | 15 |
| Ningbo | 宁波 | 20 | 6 | 6 | 6 | 10 | 10 | 12 | 27 | 6 | 17 |
| Anging | 安 庆 | 18 | 15 | 22 | 13 | 16 | 17 | 16 | 28 | 13 | 10 |
| Yichun | 宜春 | 25 | 9 | 9 | 9 | 11 | 15 | 9 | 29 | 9 | 16 |
| Suzhou | 宿州 | 30 | 13 | 19 | 12 | 16 | 15 | 15 | 31 | 12 | 12 |
| Yuevang | 岳阳 | 32 | 20 | 23 | 18 | 23 | 18 | 24 | 26 | 18 | 6 |
| Pingdingshan | 亚而山 | 27 | 14 | 14 | 15 | | | 20 | | 14 | 11 |
| Kaifeng | モ封 | 10 | 3 | 3 | 9 | | | 5 | 33 | 3 | 21 |
| Zhangijakou | | 24 | 6 | 6 | 7 | 14 | 9 | 11 | 26 | 6 | 18 |
| Xuzhou | 永 谷 州 | 28 | 17 | 17 | 17 | 20 | 20 | 19 | 30 | 17 | 7 |
| Huizbou | 声 州 | 20 | 16 | 16 | 18 | 20 | 20 | 14 | 28 | 1/ | 0 |
| Chongdu | 志州 | 20 | 10 | 20 | 20 | | 20 | 14 | 20 | 14 | 9 11 |
| Euzbou | 火和 | 20 | 20 | 20 | 20 | 17 | 17 | 15 | 20 | 15 | 10 |
| Fuziliou Vievu | 北川 | 24 | 10 | 10 | 10 | 17 | 17 | 10 | 29 | 10 | 10 |
| Musi | 初ホ | 33 | 27 | 21 | 20 | 10 | 20 | 29 | 29 | 20 | 3 |
| VVUXI | 无物 | 22 | 11 | 9 | 9 | 13 | 11 | 0 | 30 | 0 | 10 |
| Hangzhou | 机州 | 19 | 16 | 16 | 12 | 21 | 16 | 15 | 27 | 12 | 11 |
| Talan | 泰安 | 14 | 3 | 3 | 3 | 6 | 6 | | 25 | 3 | 21 |
| l alznou | 泰州 | 24 | 13 | 13 | 12 | 17 | 17 | 14 | 30 | 12 | 12 |
| Jinan | 济南 | 23 | 15 | 15 | 15 | | 17 | 15 | 23 | 15 | 9 |
| Jining | 济宁 | 24 | 10 | 10 | 8 | 14 | 10 | 8 | 25 | 8 | 16 |
| Haikou | 海口 | 24 | 20 | 20 | 16 | _ | 20 | 13 | 25 | 13 | 9 |
| Huaian | 淮安 | 21 | 12 | 9 | 5 | 7 | 8 | 7 | 30 | 5 | 19 |
| Shenzhen | 深圳 | 15 | | | 23 | 28 | 28 | 28 | 28 | 23 | 1 |
| Wenzhou | 温州 | 22 | 20 | 20 | 19 | 26 | 23 | 17 | 27 | 17 | 4 |
| Weifang | 潍坊 | 20 | 6 | 6 | 6 | 7 | 10 | 9 | 30 | 6 | 18 |
| Zhuhai | 珠海 | 16 | 13 | 14 | 19 | 25 | 19 | 13 | 28 | 13 | 10 |
| Yiyang | 益阳 | 21 | 21 | 23 | 21 | 25 | 23 | 22 | 29 | 21 | 3 |
| Yancheng | 盐城 | 20 | 18 | 19 | 15 | | 20 | 17 | 30 | 15 | 9 |
| Shijiazhuang | 石家庄 | 18 | 17 | 17 | 14 | 17 | 17 | 12 | 26 | 12 | 10 |
| Shaoxing | 绍兴 | 7 | 10 | 10 | 9 | 9 | 13 | 11 | 26 | 9 | 14 |
| Wuhu | 芜湖 | 8 | 11 | | 10 | 17 | 13 | 12 | 26 | 10 | 14 |
| Suzhou | 苏州 | 16 | 12 | 12 | 11 | 15 | 16 | 10 | 30 | 10 | 13 |
| Pingxiang | 萍乡 | 25 | 16 | 16 | 16 | 17 | 18 | 16 | 29 | 16 | 10 |
| Quzhou | 衢州 | 15 | 15 | 15 | 14 | 21 | 18 | 14 | 23 | 14 | 9 |
| Xi'an | 西安 | 19 | 14 | 17 | 15 | 18 | 16 | 13 | 28 | 13 | 10 |
| Xuchang | 许昌 | 21 | 4 | 6 | 7 | 7 | 7 | 7 | 27 | 4 | 20 |
| Ganzhou | 赣州 | 29 | 20 | 20 | 19 | 35 | 24 | 26 | 29 | 19 | 6 |
| Zhenazhou | 郑州 | 11 | 22 | 22 | 23 | 23 | 23 | 17 | 31 | 17 | 4 |
| Chenzhou | 林小山 | 11 | 13 | 13 | 12 | 16 | 16 | 11 | 30 | 11 | 12 |
| Jinhua | 全化 | 21 | 12 | 12 | 11 | 15 | 15 | 11 | 27 | 11 | 12 |
| Changehun | 亚 干 长寿 | 13 | 13 | 13 | 13 | 16 | 14 | 11 | 26 | 11 | 12 |
| Fuvana | 以甘 自 归 | 10 | 15 | 15 | 15 | 22 | 17 | 16 | 20 | 1/ | 10 |
| i uyany Ma'anshan | 中中 正蛇正 | 10 | 10 | 20 | ۱ <u>ی</u> | 20 12 | 12 | 10 | 23 10 | 0 | 15 |
| Thumodion | 与 牧 山 山 市 山 正 | 0 | 10 | 20 10 | 9 1 / | 13 | 1Z 04 | 11 | 42 04 | 9 10 | 10 |
| Znumadian | 牡勻店 | Э | ١Z | 12 | 14 | | 21 | 14 | 31 | 12 | 12 |

Appendix Table 4. Data used in the analysis, which is also available at Github and can be downloaded from https://github.com/MeyersLabUTexas/Proactive-social-distancing-in-Chinese-cities

References

- Zhang J, Litvinova M, Wang W, Wang Y, Deng X, Chen X, et al. Evolving epidemiology and transmission dynamics of coronavirus disease 2019 outside Hubei province, China: a descriptive and modelling study. Lancet Infect Dis. 2020 Apr 2 [Epub ahead of print]. PubMed https://doi.org/10.1016/S1473-3099(20)30230-9
- 2. Cori A. EpiEstim: Estimate time varying reproduction numbers from epidemic curves. 2019 [cited 2020 Jun 1]. https://cran.r-project.org/web/packages/EpiEstim/index.html
- 3. Thompson RN, Stockwin JE, van Gaalen RD, Polonsky JA, Kamvar ZN, Demarsh PA, et al. Improved inference of time-varying reproduction numbers during infectious disease outbreaks. Epidemics. 2019;29:100356. <u>PubMed</u> <u>https://doi.org/10.1016/j.epidem.2019.100356</u>
- 4. Cowling BJ, Ali ST, Ng TWY, Tsang TK, Li JCM, Fong MW, et al. Impact assessment of nonpharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study. Lancet Public Health. 2020;5:e279–88. <u>PubMed</u> <u>https://doi.org/10.1016/S2468-2667(20)30090-6</u>
- 5. Hurvich CM, Tsai C-L. Regression and time series model selection in small samples. Biometrika. 1989;76:297–307.
- 6. Yang H. The case for being automatic: introducing the automatic linear modeling (LINEAR) procedure in SPSS statistics. Multiple Linear Regression Viewpoints. 2013;39:27–37.
- 7. IBM Knowledge Center. Information Criteria [cited 2020 Mar 30]. https://www.ibm.com/support/knowledgecenter/SSLVMB_23.0.0/spss/tutorials/mixed_diet_i nfo_02.html



Appendix Figure 1. Predicted versus observed speed of containment (λ) for 36 cities in mainland China based on the fitted regression model (Appendix Table 1). The data have a Pearson correlation coefficient of 0.52 (p<0.001).



Appendix Figure 2. The number of days between the first reported case and containment (λ) for 58 cities in mainland China that achieved containment before February 15, 2020. Outbreaks are considered contained when the estimated upper 95% CI bound of R_t drops below 1 without rebounding. Orange indicates the 48 cities that achieved containment within 4 weeks; blue indicates the 10 cities that did not.