

## SARS-CoV-2 Superspread in Fitness Center, Hong Kong, China, March 2021

Daniel K.W. Chu,<sup>1</sup> Haogao Gu,<sup>1</sup> Lydia D.J. Chang, Sammi S.Y. Cheuk, Shreya Gurung, Pavithra Krishnan, Daisy Y.M. Ng, Gigi Y.Z. Liu, Carrie K.C. Wan, Dominic N.C. Tsang, Malik Peiris, Leo L.M. Poon

Author affiliations: The University of Hong Kong, Hong Kong, China (D.K.W. Chu, H. Gu, L.D.J. Chang, S.S.Y. Cheuk, S. Gurung, P. Krishnan, D.Y.M. Ng, G.Y.Z. Liu, C.K.C. Wan, M. Peiris, L.L.M. Poon); The Government of Hong Kong Special Administrative Region, Hong Kong (D.N.C. Tsang)

DOI: <https://doi.org/10.3201/eid2708.210833>

To investigate a superspreading event at a fitness center in Hong Kong, China, we used genomic sequencing to analyze 102 reverse transcription PCR–confirmed cases of severe acute respiratory syndrome coronavirus 2 infection. Our finding highlights the risk for virus transmission in confined spaces with poor ventilation and limited public health interventions.

Hong Kong, China, is at the end of a fourth wave of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. The virus causing this wave was introduced in September 2020 (GISAID clade GH) (1) and has continued to evolve in Hong Kong. As of April 30, 2021, a total of 11,771 SARS-CoV-2 cases had been laboratory confirmed; more than half (56%) were detected during the fourth wave. We describe a superspreading event that occurred in a 3,000-ft<sup>2</sup> fitness center in March 2021 (Appendix Figure 1, <https://wwwnc.cdc.gov/EID/article/27/8/21-0833-App1.pdf>).

On March 10, 2021, an asymptomatic 27-year-old male fitness trainer (patient FC1) received a positive reverse transcription PCR (RT-PCR) test result as part of a voluntary coronavirus disease (COVID-19) screening program. This program provided services to persons for community or private purposes (e.g., for work or travel). The fitness trainer had previously received a negative COVID-19 test result on February 17, 2021. He taught small group classes in the fitness center every day from February 28 through March 8, except March 4, 2021.

His positive test result triggered a local health authority to conduct epidemiologic investigation and contact tracing. The fitness center was immediately

closed to the public. The local government also issued a compulsory testing notice to those who had visited this center from February 25 through March 10. About 300 visitors were tested and 101 cases were confirmed (7 staff members and 94 customers; case-patients FC2–FC102) (Appendix Table 1). All case-patients had recently visited this center; >80% of cases were detected within 3 days of the first case (Appendix Figure 2). Another 53 SARS-CoV-2–positive persons were subsequently identified; they had had close contact with the 102 case-patients but no epidemiologic link to the fitness center.

Of the 102 case-patients, all were hospitalized according to local standard practice, recovered uneventfully, and were discharged. None had received COVID-19 vaccination before this outbreak. A total of 46 case-patients were asymptomatic at the time of testing. The percentage of asymptomatic case-patients in this cluster (45%) is higher than that of all persons with confirmed cases in Hong Kong (30%;  $p<0.005$ ). It is not known whether the general physical well-being of case-patients in this cluster affected their clinical status. Their ages, on average, were lower than that of all persons with confirmed cases in Hong Kong (38 vs. 44 years;  $p<0.005$ ).


Among the 56 symptomatic case-patients, signs and symptoms started to develop for 36 of them during March 9–11; the earliest onset date was March 6 (case-patient FC46). Assuming the average incubation period of COVID-19 to be  $\approx 5$  days (2), the superspreading event might have occurred around March 5. Because SARS-CoV-2 can be transmitted by asymptomatic and presymptomatic persons (3), our data did not enable us to identify the index case-patient of this cluster.

To exclude unrelated transmission chains in this fitness center, we used next-generation sequencing to study respiratory samples from 59 of the case-patients (1,4). We used 5 epidemiologically unrelated local case-patients, including 4 detected in the same period, as controls. All virus sequences from the fitness center outbreak genetically clustered together and were genetically distinct from the controls (Figure), demonstrating that this superspreading event was caused by a single virus introduction.

Many case-patients, including FC1 and FC46, were located at the root of this phylogenetic branch. There are a few minor clades in this phylogenetic branch, suggesting that the initial introduction triggered multiple independent transmission chains thereafter in this setting.

SARS-CoV-2 transmission in fitness centers/gyms has been reported (5–8). SARS-CoV-2 can be

<sup>1</sup>These first authors contributed equally to this article.



**Figure.** Phylogenetic tree of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) viruses detected in a fitness club in Hong Kong, China, in March 2021. Viruses from clades L, S, V, G, GH, GR, and O (others) are also included in the analysis. Near full-length genomes of studied samples were deduced by a previously described Illumina (<https://www.illumina.com>) sequencing protocol (sequence coverage >100) (1,4). Human SARS-CoV-2 WIV04 is selected to be the root of this phylogenetic tree. The tree was constructed by using the neighbor-joining method. Only bootstrap values >80 are shown. EPI ISL accession nos. for sequences retrieved in GISAID (<https://www.gisaid.org>) are provided. Scale bar indicates estimated genetic distance.

transmitted by close contact, droplets, or fomites (9). Uncontrolled physical activities in a fitness center might produce any or all of these transmission modes (e.g., increased physical contact, increased levels of exhaled respiratory droplets in a confined space because of vigorous breathing, and shared communal space and equipment). Although in this study we were unable to identify the predominant transmission mode accounting for this superspreading event, a recent report indicates that physical activities in a fitness center can create a pronounced level of saliva aerosol (10). An air change rate of 2.2/hour in a fitness center is insufficient to dilute the amount of saliva aerosol generated from physical activities (10). Of note, mask wearing during exercise was not compulsory by law at the time of this outbreak. Many case-patients in our study reported not wearing a mask while training at that time (e.g., weight training, high-intensity circuit training, and boxing). A follow-up investigation revealed that this center has air conditioning units but lacks a fresh air and exhaust duct system. This finding suggests that poor ventilation might have played a major role in this outbreak.

After this outbreak, new recommendations were issued to prevent superspreading events at fitness centers in Hong Kong. For staff in these settings, RT-PCR testing for SARS-CoV-2 every 2 weeks is compulsory, and staff are prioritized to receive COVID-19 vaccination. For all persons in fitness settings, mask wearing at all times is now mandatory, except when showering or eating. Recommendations for air ventilation in all fitness centers are under investigation.

### Acknowledgments

We gratefully acknowledge the staff from the originating laboratories responsible for obtaining the specimens and from the submitting laboratories where the genome data were generated and shared via GISAID (Appendix Table 2). We acknowledge the technical support provided by colleagues from the Centre for PanorOmic Sciences of the University of Hong Kong.

Virus sequences reported in this study are available in GISAID (<http://platform.gisaid.org>; accession nos. EPI\_ISL\_1824501 to EPI\_ISL\_1824564). The epidemiologic data for the 102 case-patients can be accessed in a public database (<https://data.gov.hk/en-data/dataset/hk-dh-chpsebctdr-novel-infectious-agent>).

This work is supported by grants from the National Institute of Allergy and Infectious Diseases (U01AI151810), and the Health and Medical Research

Fund (COVID190205). We also acknowledge the Centre for Health Protection of the Department of Health for providing epidemiologic data for the study.

### About the Author

Dr. Chu is a research assistant professor at The University of Hong Kong, China. His research interests focus on diagnostic virology, molecular diagnostics, and virus evolution.

### References

1. Chu DKW, Hui KPY, Gu H, Ko RLW, Krishnan P, Ng DYM, et al. Introduction of ORF3a-Q57H SARS-CoV-2 variant causing fourth epidemic wave of COVID-19, Hong Kong, China. *Emerg Infect Dis.* 2021;27:1492–5. <https://doi.org/10.3201/eid2705.210015>
2. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med.* 2020;382:1199–207. <https://doi.org/10.1056/NEJMoa2001316>
3. Wu P, Liu F, Chang Z, Lin Y, Ren M, Zheng C, et al. Assessing asymptomatic, pre-symptomatic and symptomatic transmission risk of SARS-CoV-2. *Clin Infect Dis.* 2021;ciab271. <https://doi.org/10.1093/cid/ciab271>
4. Choi EM, Chu DKW, Cheng PKC, Tsang DNC, Peiris M, Bausch DG, et al. In-flight transmission of SARS-CoV-2. *Emerg Infect Dis.* 2020;26:2713–6. <https://doi.org/10.3201/eid2611.203254>
5. Groves LM, Usagawa L, Elm J, Low E, Manuzak A, Quint J, et al. Community transmission of SARS-CoV-2 at three fitness facilities – Hawaii, June–July 2020. *MMWR Morb Mortal Wkly Rep.* 2021;70:316–20. <https://doi.org/10.15585/mmwr.mm7009e1>
6. Bae S, Kim H, Jung TY, Lim JA, Jo DH, Kang GS, et al. Epidemiological characteristics of COVID-19 outbreak at fitness centers in Cheonan, Korea. *J Korean Med Sci.* 2020;35:e288. <https://doi.org/10.3346/jkms.2020.35.e288>
7. Lendacki FR, Teran RA, Gretsich S, Fricchione MJ, Kerins JL. COVID-19 outbreak among attendees of an exercise facility – Chicago, Illinois, August–September 2020. *MMWR Morb Mortal Wkly Rep.* 2021;70:321–5. <https://doi.org/10.15585/mmwr.mm7009e2>
8. Jang S, Han SH, Rhee JY. Cluster of coronavirus disease associated with fitness dance classes, South Korea. *Emerg Infect Dis.* 2020;26:1917–20. <https://doi.org/10.3201/eid2608.200633>
9. Sia SF, Yan LM, Chin AWH, Fung K, Choy KT, Wong AYL, et al. Pathogenesis and transmission of SARS-CoV-2 in golden hamsters. *Nature.* 2020;583:834–8. <https://doi.org/10.1038/s41586-020-2342-5>
10. Blocken B, van Druenen T, Ricci A, Kang L, van Hooff T, Qin P, et al. Ventilation and air cleaning to limit aerosol particle concentrations in a gym during the COVID-19 pandemic. *Build Environ.* 2021;193:107659. <https://doi.org/10.1016/j.buildenv.2021.107659>

Address for correspondence: Leo Poon, School of Public Health, The University of Hong Kong, Rm 528, Lab Block 21, Sassoon Rd, Pokfulam, Hong Kong, China; email: llmpoon@hku.hk

# SARS-CoV-2 Superspread in Fitness Center, Hong Kong, China, March 2021

## Appendix

**Appendix Table 1.** Epidemiology data of FC1-FC2

Case no.	Report date (2021)	Sex	Age	Onset date (2021)
FC1	Mar 10	Male	27	Asymptomatic
FC2	Mar 11	Male	46	Mar 9
FC3	Mar 11	Male	40	Asymptomatic
FC4	Mar 11	Male	56	Asymptomatic
FC5	Mar 11	Male	30	Mar 8
FC6	Mar 11	Male	44	Mar 8
FC7	Mar 11	Male	47	Mar 9
FC8	Mar 11	Female	23	Mar 8
FC9	Mar 11	Female	30	Mar 9
FC10	Mar 11	Female	30	Mar 9
FC11	Mar 11	Male	34	Asymptomatic
FC12	Mar 11	Male	41	Asymptomatic
FC13	Mar 11	Female	19	Mar 9
FC14	Mar 11	Male	47	Mar 9
FC15	Mar 11	Female	54	Asymptomatic
FC16	Mar 11	Female	31	Asymptomatic
FC17	Mar 11	Male	35	Asymptomatic
FC18	Mar 12	Female	42	Mar 8
FC19	Mar 12	Male	34	Mar 7
FC20	Mar 12	Female	42	Asymptomatic
FC21	Mar 12	Male	39	Asymptomatic
FC22	Mar 12	Male	33	Mar 10
FC23	Mar 12	Female	26	Mar 10
FC24	Mar 12	Male	36	Mar 9
FC25	Mar 12	Male	34	Mar 10
FC26	Mar 12	Male	39	Mar 10
FC27	Mar 12	Male	53	Mar 10
FC28	Mar 12	Female	33	Asymptomatic
FC29	Mar 12	Male	42	Mar 11
FC30	Mar 12	Male	37	Asymptomatic
FC31	Mar 12	Male	46	Mar 10
FC32	Mar 12	Female	38	Mar 9
FC33	Mar 12	Male	41	Asymptomatic
FC34	Mar 12	Female	22	Asymptomatic
FC35	Mar 12	Male	36	Asymptomatic
FC36	Mar 12	Male	20	Asymptomatic
FC37	Mar 12	Female	35	Mar 10
FC38	Mar 12	Female	33	Mar 10
FC39	Mar 12	Female	24	Mar 10
FC40	Mar 12	Female	37	Mar 11
FC41	Mar 12	Male	36	Mar 11
FC42	Mar 12	Female	38	Asymptomatic
FC43	Mar 12	Male	40	Mar 9
FC44	Mar 12	Female	36	Mar 9
FC45	Mar 12	Male	44	Mar 8
FC46	Mar 12	Female	32	Mar 6
FC47	Mar 12	Male	34	Asymptomatic
FC48	Mar 12	Female	37	Mar 10
FC49	Mar 12	Male	41	Mar 9
FC50	Mar 12	Male	27	Asymptomatic
FC51	Mar 12	Male	45	Asymptomatic
FC52	Mar 12	Male	31	Mar 11
FC53	Mar 12	Female	44	Asymptomatic
FC54	Mar 12	Female	25	Asymptomatic


Case no.	Report date (2021)	Sex	Age	Onset date (2021)
FC55	Mar 12	Female	38	Asymptomatic
FC56	Mar 12	Male	41	Mar 11
FC57	Mar 12	Male	36	Asymptomatic
FC58	Mar 13	Female	37	Mar 11
FC59	Mar 13	Female	37	Asymptomatic
FC60	Mar 13	Male	61	Asymptomatic
FC61	Mar 13	Female	28	Mar 11
FC62	Mar 13	Male	48	Asymptomatic
FC63	Mar 13	Female	34	Mar 11
FC64	Mar 13	Male	36	Mar 11
FC65	Mar 13	Female	31	Mar 11
FC66	Mar 13	Male	35	Mar 10
FC67	Mar 13	Female	36	Mar 11
FC68	Mar 13	Male	40	Asymptomatic
FC69	Mar 13	Male	49	Mar 12
FC70	Mar 13	Male	33	Asymptomatic
FC71	Mar 13	Male	38	Asymptomatic
FC72	Mar 13	Male	37	Asymptomatic
FC73	Mar 13	Male	48	Mar 11
FC74	Mar 13	Male	62	Asymptomatic
FC75	Mar 13	Male	39	Asymptomatic
FC76	Mar 13	Male	36	Asymptomatic
FC77	Mar 13	Male	33	Mar 12
FC78	Mar 13	Female	36	Mar 12
FC79	Mar 13	Female	48	Asymptomatic
FC80	Mar 13	Male	26	Mar 9
FC81	Mar 13	Female	36	Asymptomatic
FC82	Mar 13	Female	39	Mar 10
FC83	Mar 14	Male	47	Mar 13
FC84	Mar 14	Female	37	Asymptomatic
FC85	Mar 14	Male	31	Asymptomatic
FC86	Mar 14	Male	46	Asymptomatic
FC87	Mar 14	Female	34	Mar 13
FC88	Mar 14	Female	37	Mar 13
FC89	Mar 15	Female	43	Asymptomatic
FC90	Mar 15	Male	47	Mar 13
FC91	Mar 15	Female	45	Mar 13
FC92	Mar 15	Female	36	Mar 14
FC93	Mar 16	Male	42	Asymptomatic
FC94	Mar 17	Female	34	Asymptomatic
FC95	Mar 18	Male	42	Mar 16
FC96	Mar 19	Male	46	Mar 17
FC97	Mar 20	Male	44	Asymptomatic
FC98	Mar 21	Female	37	Asymptomatic
FC99	Mar 22	Male	39	Mar 20
FC100	Mar 22	Female	28	Asymptomatic
FC101	Mar 22	Female	39	Mar 20
FC102	Mar 23	Male	33	Asymptomatic

**Appendix Table 2.** Studied sequences from GISAID


Virus name	Accession No.	Collected	Originating laboratory	Submitting laboratory	Submitted by
Australia/VIC1158/2020	EPI_ISL_430602	2020 Apr 9	Victorian Infectious Diseases Reference Laboratory (VIDRL)	Microbiological Diagnostic Unit Public Health Laboratory and Victorian Infectious Diseases Reference Laboratory, The Peter Doherty Institute for Infection and Immunity	Caly L. et al
Australia/VIC1787/2020	EPI_ISL_456649	2020 May 27	Victorian Infectious Diseases Reference Laboratory (VIDRL)	Microbiological Diagnostic Unit Public Health Laboratory and Victorian Infectious Diseases Reference Laboratory, Doherty Institute	Caly L. et al
Australia/VIC98/2020	EPI_ISL_419810	2020 Mar 16	Victorian Infectious Diseases Reference Laboratory (VIDRL)	Victorian Infectious Diseases Reference Laboratory and Microbiological Diagnostic Unit Public Health Laboratory, Doherty Institute	Caly L. et al
England/20132080404/2020	EPI_ISL_423805	2020 Mar 24	Respiratory Virus Unit, Microbiology Services Colindale, Public Health England	Respiratory Virus Unit, Microbiology Services Colindale, Public Health England	Monica Galiano et al
England/CAMB-77909/2020	EPI_ISL_439431	2020 Mar 31	Department of Pathology, University of Cambridge	Wellcome Sanger Institute for the COVID-19 Genomics UK (COG-UK) consortium	Luke W Meredith et al
England/NORW-EC85D/2020	EPI_ISL_457566	2020 May 13	Quadram Institute Bioscience	COVID-19 Genomics UK (COG-UK) Consortium	Dave J. Baker et al
England/SHEF-C03E5/2020	EPI_ISL_420221	2020 Mar 29	Virology Department, Sheffield Teaching Hospitals NHS Foundation Trust	Department of Infection, Immunity and Cardiovascular Disease, The Florey Institute, The Medical School, University of Sheffield	Thushan de Silva et al
France/HDF-3668/2020	EPI_ISL_443316	2020 Mar 25	CH Compiègne Laboratoire de Biologie	National Reference Center for Viruses of Respiratory Infections, Institut Pasteur, Paris	Mélanie Albert et al
HongKong/VB20175856/2020	EPI_ISL_539823	2020 Aug 14	Communicable Disease Branch	Hong Kong Department of Health	Alan K.L. Tsang et al
HongKong/VM20066524/2020	EPI_ISL_539814	2020 Jul 21	Tuen Mun Hospital	Hong Kong Department of Health	Alan K.L. Tsang et al
Iceland/99/2020	EPI_ISL_417768	2020 Mar 10	The National University Hospital of Iceland	deCODE genetics	Daniel F Gudbjartsson et al
India/OR-RMRC164/2020	EPI_ISL_455764	2020 May 7	REGIONAL VRDL, ICMR-RMRC BBSR	Immunogenomics lab, Institute of Life Sciences, Bhubaneswar	Sunil Raghav et al
Japan/Hu_DP_Kng_19-031/2020	EPI_ISL_420889	2020 Feb 14	Takayuki Hishiki Kanagawa Prefectural Institute of Public Health	Takayuki Hishiki Kanagawa Prefectural Institute of Public Health	Hishiki et al
NewZealand/20VR3045/2020	EPI_ISL_456402	2020 Apr 25	Wellington SCL	Institute of Environmental Science and Research (ESR)	Matt Storey et al



Virus name	Accession No.	Collected	Originating laboratory	Submitting laboratory	Submitted by
Portugal/PT0163/2020	EPI_ISL_453879	2020 Mar 28	unknown	Instituto Nacional de Saude (INSA)	Borges et al et al
Spain/MD-ISCI-201738/2020	EPI_ISL_419237	2020 Mar 7	Fundacion Jimenez Diaz	Instituto de Salud Carlos III	Iglesias-Caballero et al
USA/CA-CZB-1237/2020	EPI_ISL_454657	2020 May 12	County of Santa Clara Public Health Department	Chan-Zuckerberg Biohub	CZB Cliahub Consortium et al
Wales/PHWC-3180E/2020	EPI_ISL_446492	2020 Apr 11	Wales Specialist Virology Centre	Public Health Wales Microbiology Cardiff	Catherine Moore et al
Wales/PHWC-32B6E/2020	EPI_ISL_446656	2020 Apr 14	Wales Specialist Virology Centre	Public Health Wales Microbiology Cardiff	Catherine Moore et al
Wuhan/HB-WH1-131/2020	EPI_ISL_454910	2020 Mar 2	Wuhan Chain Medical Labs (CMLabs)	State Key Laboratory of Biotherapy of Sichuan University	Baowen Du et al
Wuhan/WIV04/2019	EPI_ISL_402124	2019 Dec 30	Wuhan Jinyintan Hospital	Wuhan Institute of Virology, Chinese Academy of Sciences	Peng Zhou et al



**Appendix Figure 1.** Coronavirus disease (COVID-19) pandemic waves in Hong Kong, China. Number of cases with different epidemiologic links are shown. Arrow indicates the outbreak that occurred in the fitness center in Hong Kong in March 2021. This Figure is modified from the one available from the School of Public Health, The University of Hong Kong (<https://covid19.sph.hku.hk/>).



**Appendix Figure 2.** COVID-19 cases epidemiologically linked to a fitness center. The number of daily detected cases, daily detected asymptomatic cases, and reported onset dates in the studied period are shown. The first reported case was detected on 10 Mar 2021.