

Highly Pathogenic Avian Influenza A(H5N1) Virus–Induced Mass Death of Wild Birds, Caspian Sea, Russia, 2022

Ivan Sobolev, Alimurad Gadzhiev, Kirill Sharshov, Olesia Ohlopkova, Kristina Stolbunova, Artem Fadeev, Nikita Dubovitskiy, Alexandra Glushchenko, Victor Irza, Maxim Perkovsky, Kirill Litvinov, Natalia Meshcheriakova, Guy Petherbridge, Alexander Shestopalov

In May 2022, we observed a substantial die-off of wild migratory waterbirds on Maliy Zhemchuzhniy Island in the Caspian Sea, Russia. The deaths were caused by highly pathogenic avian influenza A(H5N1) clade 2.3.4.4.b virus. Continued surveillance of influenza viruses in wild bird populations is needed to predict virus spread over long distances.

The coastal and estuarine wetlands of the northern Caspian Sea, which borders southeast Russia, provide support for millions of waterfowl and shorebirds during nesting, molting, migration, and wintering periods (1,2); >300 species of birds are found in this region. The area is crossed by several migration flyways (Figure 1), of which the Black Sea/Mediterranean Flyway is the main migratory route (3). The region plays a critical role in the reproduction of colonial nesting birds, such as pelicans, cormorants, herons, gulls, and terns.

Maliy Zhemchuzhniy Island is located in the northern part of the Caspian Sea (Figure 1). Monitoring data on waterbirds has shown the high ecologic

importance of this area, not only during the nesting period but also during bird migration. The island has had >150 species of birds registered since 2016. A breeding colony of Caspian gulls is located on the island, along with colonies of Great black-headed gulls and Caspian terns, which are all listed in the Red Data Book of Russia. We investigated mass deaths of wild migratory waterbirds on Maliy Zhemchuzhniy Island that occurred in May 2022. The study was approved by the Committee on Biomedical Ethics at the Federal Research Center of Fundamental and Translational Medicine in Novosibirsk, Russia (protocol nos. 2013-23, 2019-3, and 2021-10).

The Study

On April 28, 2022, near the end of the egg incubation period, we had counted a total of 26,769 Great black-headed gull nests, 7,340 Caspian gull nests, and 5,267 Caspian tern nests on Maliy Zhemchuzhniy Island. In May, 1 week later, we detected mass deaths of waterbirds on the island comprising 25,157 Great black-headed gulls, 3,507 Caspian gulls, 5,641 Caspian terns, and 14 Dalmatian pelicans (Appendix 1 Figure 1, <https://wwwnc.cdc.gov/EID/article/29/12/23-0330-App1.pdf>). Nearly all gull and tern chicks died during the nesting period. The mass death event began during hatching of Great black-headed gulls. We only found the corpses of chicks (with down but without feathers) that were similar in age. We assume that not all of the chicks actually died from disease; death of adult birds likely led to the deaths of chicks in their nests. The Caspian terns were still incubating eggs at that time; consequently, the death of adult terns led to the death of egg clutches in their nests. We

Author affiliations: Federal Research Center of Fundamental and Translational Medicine, Novosibirsk, Russia (I. Sobolev, K. Sharshov, O. Ohlopkova, K. Stolbunova, N. Dubovitskiy, A. Glushchenko, A. Shestopalov); Dagestan State University, Makhachkala, Russia (A. Gadzhiev); Smorodintsev Research Institute of Influenza, Saint Petersburg, Russia (A. Fadeev); Federal Governmental State-Financed Institution Federal Center for Animal Health, Vladimir, Russia (V. Irza); Astrakhan State Biosphere Nature Reserve, Astrakhan, Russia (M. Perkovsky, K. Litvinov, N. Meshcheriakova); Caspian Center for Nature Conservation, Makhachkala (G. Petherbridge)

DOI: <https://doi.org/10.3201/eid2912.230330>




Figure 1. Major bird migration flyways (A) and sampling area of birds (B) in study of highly pathogenic avian influenza A(H5N1) virus–induced mass death of wild birds, Caspian Sea, Russia, 2022. Map of migration routes was provided online by the East Asian–Australasian Flyway Partnership (<https://www.eaaflyway.net/the-flyway>). Yellow shading in panel A indicates the location of the Caspian Sea; red rectangles in both panels indicate sampling location of dead birds on Maliy Zhemchuzhnyi Island.

did not observe live chicks on the island during the remaining 2022 nesting season.

In May 2022, we collected 10 samples from deceased Caspian terns on the island. All samples tested positive for H5Nx avian influenza virus (AIV) by real-time PCR. We characterized 5 isolates by using complete genome sequencing, phylogenetic analysis, and intravenous pathogenicity index testing (Table). We identified all 5 isolates as highly pathogenic avian influenza (HPAI) viruses, according to the amino acid sequence of the hemagglutinin (HA) polybasic proteolytic cleavage site (PLREKRRKR/G) (Appendix 1 Figure 2) and according to intravenous pathogenicity index values of 2.92–2.93 (reference values were for chickens) (Appendix). We determined the HPAI neuraminidase (NA) subtype was N1 by using sequence analysis.

The A/goose/Guangdong/1/96 (Gs/GD) strain, which was isolated in 1996 from a domestic goose, is considered the ancestor of AIV carrying the highly pathogenic H5 HA subtype (4,5). The Gs/GD lineage of HPAI H5N1 viruses evolved into several sublineages that subsequently reassorted with low pathogenicity avian influenza (LPAI) viruses, leading to the formation of H5Nx reassortant variants (6,7). Viruses of the Gs/GD lineage ini-

tially circulated in southern China. However, during 2004–2005, they began spreading throughout Asia, Europe, the Middle East, and Africa (8). An increased number of available sequences led to the creation of the H5 virus classification and identification of 10 clades (4,8–13). Subsequently, clade 2 was split into 5 subclades, each containing numerous genetic subgroups. Clade 2.3.4.4 is dominant and has been divided into several genetic subgroups, including 2.3.4.4.b, which, in turn, diverged into 2 sublineage branches, B1 and B2 (14). According to phylogenetic analysis of HA segments, we found the Caspian Sea strains (the AIV strains isolated from dead birds in the Caspian Sea region) belonged to HPAI H5N1 clade 2.3.4.4.b (Figure 2). Furthermore, we found the HPAI H5N1 virus isolates from Maliy Zhemchuzhnyi Island belonged to the B2 sublineage because isoleucine was present at position 548 in HA (14).

The polymerase basic (PB) 1, polymerase acidic (PA), HA, nucleoprotein (NP), NA, and matrix (M) protein gene segments of the Caspian Sea strains were phylogenetically related to H5N1 viruses previously identified in wild birds in Israel in January 2022. However, all 8 gene segments were similar to those of strains from Romania isolated in February

Table. Sequenced viruses isolated from Caspian terns on Maliy Zhemchuzhnyi Island in study of highly pathogenic avian influenza A(H5N1) virus–induced mass death of wild birds, Caspian Sea, Russia, 2022*

Isolate	Location	Sample type	Collection date	IVPI	GISAID no.
A/Caspian_tern/Astrakhan/30/2022	45°02' N, 48°19' E	Intestine	2022 May 15	2.92	EPI_ISL_16020401
A/Caspian_tern/Astrakhan/32/2022	45°02' N, 48°19' E	Intestine	2022 May 15	2.92	EPI_ISL_16020402
A/Caspian_tern/Astrakhan/34/2022	45°02' N, 48°19' E	Intestine	2022 May 15	2.93	EPI_ISL_16020403
A/Caspian_tern/Astrakhan/36/2022	45°02' N, 48°19' E	Liver	2022 May 15	2.93	EPI_ISL_16020404
A/Caspian_tern/Astrakhan/38/2022	45°02' N, 48°19' E	Liver	2022 May 15	2.92	EPI_ISL_16020405

*Nucleotide sequences were deposited in the GISAID database (<https://www.gisaid.org>). IVPI, intravenous pathogenicity index.

LPAI viruses during 2020–2021; NS segments likely emerged from LPAI viruses detected during 2021–2022. NS sequences closely related to those of strains isolated in the Caspian Sea regions and Romania were found in LPAI viruses predominantly circulating in Asia during 2019–2021 (Appendix 1 Figure 9). HPAI viruses with such NS sequences have been identified only in Romania and the Caspian Sea.

Gene segments of HPAI H5N1 viruses from the Caspian Sea were closely related to virus segments found in different parts of Eurasia. Specifically, PB1, PA, HA, NA, and M protein gene segments were predominantly related to those in Europe, whereas related NP and NS segments were more prevalent in Asia. In addition, the PA segment from the Caspian Sea strains was also identified in Africa, and PB2 was related to PB2 of viruses detected in the Far East (Japan, Korea, and China), Siberia (Novosibirsk region), Bangladesh, and Europe (Italy, Slovenia, and Czech Republic). Three major flyways pass through the Caspian Sea region: the Black Sea/Mediterranean Flyway, the West Asian–East African Flyway, and the Central Asian Flyway. However, we found that gene segments of HPAI viruses from the Caspian Sea were related to variants identified in the Far East, indicating widespread distribution and exchange of influenza virus genes well beyond the major flyways. Therefore, continued surveillance and monitoring of AIVs (primarily HPAI viruses) in wild bird populations will be needed worldwide to track and predict the spread of these viruses over long distances.

Acknowledgments

We thank those persons who provided sequence information to the GISAID's EpiFlu database (Appendix 2 Table, <https://wwwnc.cdc.gov/EID/article/29/12/23-0330-App2.xlsx>).

The study was supported by state assignment from the Ministry of Science and Higher Education of the Russian Federation (no. 122120600015-2 for sample collection) and Russian Science Foundation (project no. 23-64-00005 for laboratory diagnostics, virology experiments, and analysis).

About the Author

Dr. Sobolev is a researcher at the Research Institute of Virology, Federal Research Center of Fundamental and Translational Medicine, Siberian Branch, Russian Academy of Sciences, Russia. His primary research interest is the molecular diagnosis and epidemiology of avian influenza viruses.

References

- Alekseev AY, Murashkina TA, Dzhamalutdinov DM, Abdullaev SS, Akhmedrabadanov KA, Sharshov KA. Analysis of bird migrations of the aquatic and near-water complex on the territory of the Republic of Dagestan and justification for the choice of key monitoring points for influenza type A [Russian]. *South of Russia: Ecology, Development*. 2019;14:137–49. <https://doi.org/10.18470/1992-1098-2019-1-137-149>
- Petherbridge G, Gadzhiev AA, Shestapalov AM, Alekseev AY, Sharsov KA, Daudova MG. An early warning system for highly pathogenic viruses borne by waterbird species and related dynamics of climate change in the Caspian Sea region: outlines of a concept. *South of Russia: Ecology, Development*. 2022;17:223–64. <https://doi.org/10.18470/1992-1098-2022-2-233-263>
- Veen J, Yurlov AK, Delany SN, Mihantiev AI, Selivanova MA, Boere GC. An atlas of movements of southwest Siberian waterbirds. Wageningen (the Netherlands): Wetlands International; 2005.
- Dhingra MS, Artois J, Robinson TP, Linard C, Chaiban C, Xenarios I, et al. Global mapping of highly pathogenic avian influenza H5N1 and H5Nx clade 2.3.4.4 viruses with spatial cross-validation. *Elife*. 2016;5:e19571. <https://doi.org/10.7554/eLife.19571>
- Duan L, Campitelli L, Fan XH, Leung YHC, Vijaykrishna D, Zhang JX, et al. Characterization of low-pathogenic H5 subtype influenza viruses from Eurasia: implications for the origin of highly pathogenic H5N1 viruses. *J Virol*. 2007;81:7529–39. <https://doi.org/10.1128/JVI.00327-07>
- Shao W, Li X, Goraya MU, Wang S, Chen JL. Evolution of influenza A virus by mutation and re-assortment. *Int J Mol Sci*. 2017;18:1650. <https://doi.org/10.3390/ijms18081650>
- Hill NJ, Hussein IIM, Davis KR, Ma EJ, Spivey TJ, Ramey AM, et al. Reassortment of influenza A viruses in wild birds in Alaska before H5 clade 2.3.4.4 outbreaks. *Emerg Infect Dis*. 2017;23:654–7. <https://doi.org/10.3201/eid2304.161668>
- Lee DH, Bertran K, Kwon JH, Swayne DE. Evolution, global spread, and pathogenicity of highly pathogenic avian influenza H5Nx clade 2.3.4.4. *J Vet Sci*. 2017;18:269–80. <https://doi.org/10.4142/jvs.2017.18.S1.269>
- WHO/OIE/FAO H5N1 Evolution Working Group. Toward a unified nomenclature system for highly pathogenic avian influenza virus (H5N1). *Emerg Infect Dis*. 2008;14:e1. PubMed <https://doi.org/10.3201/eid1407.071681>
- WHO/OIE/FAO H5N1 Evolution Working Group. Continuing progress towards a unified nomenclature for the highly pathogenic H5N1 avian influenza viruses: divergence of clade 2.2 viruses. *Influenza Other Respir Viruses*. 2009;3:59–62. <https://doi.org/10.1111/j.1750-2659.2009.00078.x>
- WHO/OIE/FAO H5N1 Evolution Working Group. Continued evolution of highly pathogenic avian influenza A (H5N1): updated nomenclature. *Influenza Other Respir Viruses*. 2012;6:1–5. <https://doi.org/10.1111/j.1750-2659.2011.00298.x>
- World Health Organization/World Organisation for Animal Health/Food and Agriculture Organization (WHO/OIE/FAO) H5N1 Evolution Working Group. Revised and updated nomenclature for highly pathogenic avian influenza A (H5N1) viruses. *Influenza Other Respir Viruses*. 2014;8:384–8. <https://doi.org/10.1111/irv.12230>
- Claes F, Morzaria SP, Donis RO. Emergence and dissemination of clade 2.3.4.4 H5Nx influenza viruses—how is the Asian HPAI H5 lineage maintained. *Curr Opin*

- Virol. 2016;16:158–63. <https://doi.org/10.1016/j.coviro.2016.02.005>
14. Pohlmann A, King J, Fusaro A, Zecchin B, Banyard AC, Brown IH, et al. Has epizootic become enzootic? Evidence for a fundamental change in the infection dynamics of highly pathogenic avian influenza in Europe, 2021. *mBio*. 2022;13:e0060922. <https://doi.org/10.1128/mbio.00609-22>
15. Sobolev I, Sharshov K, Dubovitskiy N, Kurskaya O, Alekseev A, Leonov S, et al. Highly pathogenic avian


influenza A(H5N8) virus clade 2.3.4.4b, western Siberia, Russia, 2020. *Emerg Infect Dis*. 2021;27:2224–7. <https://doi.org/10.3201/eid2708.204969>

Address for correspondence: Ivan Sobolev, Research Institute of Virology, Federal Research Center of Fundamental and Translational Medicine, 2 Timakov St, Novosibirsk 630117, Russia; email: sobolev_i@hotmail.com

October 2023

Congenital Syphilis

- Serotype Distribution and Disease Severity in Adults Hospitalized with *Streptococcus pneumoniae* Infection, Bristol and Bath, UK, 2006–2022
- Spike in Congenital Syphilis, Mississippi, USA, 2016–2022
- Carbapenem-Resistant *Klebsiella pneumoniae* in Large Public Acute-Care Healthcare System, New York, New York, USA, 2016–2022
- Posttransfusion Sepsis Attributable to Bacterial Contamination in Platelet Collection Set Manufacturing Facility, United States
- Effects of COVID-19 on Maternal and Neonatal Outcomes and Access to Antenatal and Postnatal Care, Malawi
- Emergence of SARS-CoV-2 Delta Variant and Effect of Nonpharmaceutical Interventions, British Columbia, Canada
- Community Outbreak of *Pseudomonas aeruginosa* Infections Associated with Contaminated Piercing Aftercare Solution, Australia, 2021
- Characteristics of and Deaths among 333 Persons with Tuberculosis and COVID-19 in Cross-Sectional Sample from 25 Jurisdictions, United States
- Cycle Threshold Values as Indication of Increasing SARS-CoV-2 New Variants, England, 2020–2022
- Comprehensive Case–Control Study of Protective and Risk Factors for Buruli Ulcer, Southeastern Australia
- *Treponema pallidum* Detection at Asymptomatic Oral, Anal, and Vaginal Sites in Adults Reporting Sexual Contact with Persons with Syphilis



- *Listeria monocytogenes* Transmission from Donated Blood to Platelet Transfusion Recipient, Italy
- Imported Toxicogenic *Corynebacterium Diphtheriae* in Refugees with Polymicrobial Skin Infections, Germany, 2022
- Expansion of Invasive Group A *Streptococcus* M1UK Lineage in Active Bacterial Core Surveillance, United States, 2019–2021
- Estimate of COVID-19 Deaths, China, December 2022–February 2023
- Mpox in Children and Adolescents during Multicountry Outbreak, 2022–2023
- Outbreak of Sexually Transmitted Nongroupable *Neisseria meningitidis*-Associated Urethritis, Vietnam
- *Pseudomonas aeruginosa* High-Risk Sequence Type 463 Co-Producing KPC-2 and AFM-1 Carbapenemases, China, 2020–2022
- Seafood-Associated Outbreak of *ctx*-Negative *Vibrio mimicus* Causing Cholera-Like Illness, Florida, USA
- Influenza A(H5N1) Virus Infections in 2 Free-Ranging Black Bears (*Ursus americanus*), Quebec, Canada
- Limited Outbreak of Highly Pathogenic Influenza A(H5N1) in Herring Gull Colony, Canada, 2022
- Antimicrobial Resistance in Slaughterhouses, Kenya
- Human-to-Human Transmission of Andes Virus Modeled in Syrian Hamsters
- *Candida auris* Clinical Isolates Associated with Outbreak in Neonatal Unit of Tertiary Academic Hospital, South Africa
- Sporadic Shiga Toxin–Producing *Escherichia coli*–Associated Pediatric Hemolytic Uremic Syndrome, France, 2012–2021
- Stability of Monkeypox Virus in Body Fluids and Wastewater
- Ancestral Origin and Dissemination Dynamics of Reemerging Toxicogenic *Vibrio cholerae*, Haiti
- Managing Risk for Congenital Syphilis, Perth, Western Australia, Australia
- Estimated Costs of 4-Month Pulmonary Tuberculosis Treatment Regimen, United States
- Human Tularemia Epididymo-Orchitis Caused by *Francisella tularensis* Subspecies *holartica*, Austria

**EMERGING
INFECTIOUS DISEASES**

To revisit the October 2023 issue, go to:
<https://wwwnc.cdc.gov/eid/articles/issue/29/10/table-of-contents>

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

Highly Pathogenic Avian Influenza A(H5N1) Virus–Induced Mass Death of Wild Birds, Caspian Sea, Russia, 2022

Appendix 1

Additional Materials and Methods

Sampling Area

The Caspian Sea region is a crucial area for avian influenza virus (AIV) surveillance in Eurasia, because it is situated at the convergence of 3 major bird migratory routes: the Central Asian Flyway, East Africa/West Asia Flyway, and Black Sea/Mediterranean Flyway (Figure 1, main text). Thus, the main migratory pathways of birds from Europe, Africa, and Asia intersect here (1–3). Because of the abundance of saltwater bays, lagoons, river deltas, and lake systems, the western coast of the Caspian Sea provides a highly favorable route for the seasonal migrations of numerous bird species in Eurasia (4,5). The ecologic characteristics of those unique habitats enable wild waterfowl and shorebirds to utilize numerous wetlands in the Caspian Sea region for both wintering and nesting. Moreover, this region also provides potential stopover sites for birds that migrate further into Europe or Africa.

Reconstructive analysis of the flight routes of wild birds in general, and gulls (Laridae) in particular, was performed as previously described (5) according to data from the Bird Ringing Center of Russia (<https://sev-in.ru/en/bird-ringing-and-scientificinformation-cente>) (Appendix 1 Figures 10). As a result, it has been shown that flight routes passing through the Caspian Sea region cover not only the territory of Eurasia but also parts of Africa. They also pass through countries in Europe and the Middle East. Those results enables us to assume that the influenza virus spreads through avian spring migration to the Caspian Sea, either directly from Israel and the Levant or through Europe (likely via Romania, as indicated by phylogenetic analysis). In

addition, analysis of flight routes and migration corridors (notable for their diverse destinations) indicates the significance of the Caspian region in the spread of avian influenza virus and early detection of new variants.

Maliy Zhemchuzhniy Island is located in the northern part of the Caspian Sea near the maritime boundary of the Republic of Kalmykia and the Astrakhan region (latitude 45°02' N, longitude 48°19' E). This flat, low-lying island, composed of sand and clam shells, was formed during a regression of the Caspian Sea in the 1930s on the site of an underwater shoal. It is constantly exposed to wind and waves, which cause changes in its configuration, and areas are either completely or partially lacking terrestrial vegetation. During 2016–2022, the size of the island remained relatively constant. In 2021, the island was estimated to be 26.33 ha and had a length of ≈ 2 km and maximum width of ≈ 0.3 km.

Samples

In May 2022, we collected 10 samples from Caspian terns on the island during a mass die-off of wild birds. In all of those samples, the matrix protein gene segment of AIV A and the hemagglutinin gene segment of the H5Nx subtype were detected by using multiplex real-time PCR (AmpliSens Influenza virus A H5N1-FRT PCR kit; InterLabService Ltd, <https://en.interlabservice.ru>). All analyzed viruses were isolated from 10-day-old chicken embryonated eggs by using chicken embryo inoculation. All viruses caused the death of chicken embryos within 2 days. Isolates were shown to be H5 positive by using real-time PCR.

Genome Sequencing and Phylogenetic Analysis

Next-generation sequencing of complete genomes was performed by using the Illumina MiSeq platform (<https://www.illumina.com>) and associated reagent kits according to the manufacturer's methodology. RNA was extracted by using the QIAamp Viral RNA Mini Kit (<https://www.qiagen.com>). Whole-genome amplification was performed by using a modified protocol (6). DNA libraries were prepared by using a Nextera DNA Flex Library Prep kit (Illumina) and sequenced by using MiSeq Reagent Kit v3 (600-cycle) (Illumina). Consensus sequences were generated by using Bowtie software (<https://www.bowtiepro.com>).

Nucleotide sequences were deposited in the GISAID database under accession nos. EPI_ISL_16020401–405. Multiple alignments were performed by using MUSCLE (<https://www.ebi.ac.uk/Tools/msa/muscle>); editing, which included translating the nucleotide

sequences into amino acid sequences, was performed by using BioEdit (<https://bioedit.software.informer.com>) and UGENE (<http://www.ugene.net>) software. Initial maximum-likelihood phylogenies for each of the gene segments were generated with RAxML (7) by using the general time-reversible nucleotide substitution model. Final dendrograms were generated and visualized by using MEGA5 (8). Bootstrap support values were generated by using 1,000 rapid bootstrap replicates.

Intravenous Pathogenicity Index

For determining intravenous pathogenicity index (IVPI) values for 9 viruses, 0.1 mL of a 1:10 dilution of infectious allantoic fluid was intravenously inoculated into ten 6-week-old specific pathogen-free chickens. The IVPI was calculated according to the OIE standard protocol (https://www.woah.org/fileadmin/Home/eng/Health_standards/tahm/3.03.04_AI.pdf). Virus isolates with an IVPI value >1.2 were considered highly pathogenic avian influenza A viruses. The challenge study and all experiments with live viruses were conducted in a Biosafety Level 3 facility.


References

1. Danilenko EA, Soldatov MS. Bird migration in the Caspian region [in Russian], 2015. <http://de.geogr.msu.ru/casp>
2. Iverson SA, Gavrillov A, Katzner TE, Takekawa JY, Miller TA, Hagemeyer W, et al. Migratory movements of waterfowl in Central Asia and avian influenza emergence: sporadic transmission of H5N1 from east to west. *IBIS*. 2011;153:279–92. <https://doi.org/10.1111/j.1474-919X.2010.01095.x>
3. Gulyaeva M, Sobolev I, Sharshov K, Kurskaya O, Alekseev A, Shestopalova L, et al. Characterization of avian-like influenza A(H4N6) virus isolated from Caspian seal in 2012. *Virology*. 2018;33:449–52. [PubMed <https://doi.org/10.1007/s12250-018-0053-y>](https://doi.org/10.1007/s12250-018-0053-y)
4. Vilkov EV. The genesis and evolution of the Caspian Sea lagoons as avifauna refuges at the transboundary scale. *Open J Marine Sci*. 2016;6:115–24. <https://doi.org/10.4236/ojms.2016.61011>
5. Vilkov EV. Changes in the strategy of migration of gulls (Laridae) along the western coast of the Caspian Sea as a result of environmental changes in space and time. *Contemp Probl Ecol*. 2016;9:233–53. <https://doi.org/10.1134/S199542551603015X>

6. Zhou B, Donnelly ME, Scholes DT, St George K, Hatta M, Kawaoka Y, et al. Single-reaction genomic amplification accelerates sequencing and vaccine production for classical and swine origin human influenza A viruses. *J Virol.* 2009;83:10309–13. [PubMed https://doi.org/10.1128/JVI.01109-09](https://doi.org/10.1128/JVI.01109-09)
7. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics.* 2014;30:1312–3. [PubMed https://doi.org/10.1093/bioinformatics/btu033](https://doi.org/10.1093/bioinformatics/btu033)
8. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol.* 2011;28:2731–9. [PubMed https://doi.org/10.1093/molbev/msr121](https://doi.org/10.1093/molbev/msr121)



Appendix 1 Figure 1. Dead waterbirds found on Maliy Zhemchuzhniy Island in the northern Caspian Sea, Russia, May 2022.




Appendix 1 Figure 3. Phylogenetic analysis of neuraminidase gene segments of avian influenza viruses. Trees were constructed by using the maximum-likelihood method. Closed circles indicate the highly pathogenic avian influenza H5N1 virus strains isolated from dead Caspian terns on Maliy Zhemchuzhnyi Island, Caspian Sea, Russia in May 2022. Red box indicates virus strains from Israel and Romania closely related to the Caspian Sea strains. Sequences were obtained from the GISAID EpiFlu database (<https://www.gisaid.org>). Scale bar indicates nucleotide substitutions per site. HPAI, high pathogenicity avian influenza; LPAI, low pathogenicity avian influenza.



Appendix 1 Figure 4. Phylogenetic analysis of polymerase basic 2 protein gene segments of avian influenza viruses. Trees were constructed by using the maximum-likelihood method. Closed circles indicate the highly pathogenic avian influenza H5N1 virus strains isolated from dead Caspian terns on Maliy Zhemchuzhniy Island, Caspian Sea, Russia in May 2022. Red box indicates virus strains from

Romania closely related to the Caspian Sea strains. Red asterisks indicate low pathogenicity viruses found within highly pathogenic virus groups that were related to the Caspian Sea strains (presumably, highly pathogenic viruses appeared because of reassortment with low pathogenicity viruses). Closed triangles indicate Egyptian-like virus strains from Russia isolated in 2020. Sequences were obtained from the GISAID EpiFlu database (<https://www.gisaid.org>). Scale bar indicates nucleotide substitutions per site. HPAI, high pathogenicity avian influenza; LPAI, low pathogenicity avian influenza.




Appendix 1 Figure 5. Phylogenetic analysis of polymerase basic 1 protein gene segments of avian influenza viruses. Trees were constructed by using the maximum-likelihood method. Closed circles indicate the highly pathogenic avian influenza H5N1 virus strains isolated from dead Caspian terns on Malyy Zhemchuzhniy Island, Caspian Sea, Russia in May 2022. Red box indicates virus strains from Israel

and Romania closely related to the Caspian Sea strains. Closed triangles indicate Egyptian-like virus strains from Russia isolated in 2020. Sequences were obtained from the GISAID EpiFlu database (<https://www.gisaid.org>). Scale bar indicates nucleotide substitutions per site. HPAI, high pathogenicity avian influenza; LPAI, low pathogenicity avian influenza.




Appendix 1 Figure 6. Phylogenetic analysis of polymerase acidic protein gene segments of avian influenza viruses. Trees were constructed by using the maximum-likelihood method. Red asterisks

indicate low pathogenicity viruses found within highly pathogenic virus groups that were related to the Caspian Sea strains (presumably, highly pathogenic viruses appeared because of reassortment with low pathogenicity viruses). Red box indicates virus strains from Romania closely related to the Caspian Sea strains. Closed circles indicate the highly pathogenic avian influenza H5N1 virus strains isolated from dead Caspian terns on Maliy Zhemchuzhniy Island, Caspian Sea, Russia in May 2022. Closed triangles indicate Egyptian-like virus strains from Russia isolated in 2020. Sequences were obtained from the GISAID EpiFlu database (<https://www.gisaid.org>). Scale bar indicates nucleotide substitutions per site. HPAI, high pathogenicity avian influenza; LPAI, low pathogenicity avian influenza.




Appendix 1 Figure 7. Phylogenetic analysis of nucleoprotein gene segments of avian influenza viruses. Trees were constructed by using the maximum-likelihood method. Closed circles indicate the highly pathogenic avian influenza H5N1 virus strains isolated from dead Caspian terns on Maliy Zhemchuzhnyi Island, Caspian Sea, Russia in May 2022. Red box indicates virus strains from Israel and Romania closely related to the Caspian Sea strains. Red asterisks indicate low pathogenicity viruses found within highly pathogenic virus groups that were related to the Caspian Sea strains (presumably, highly pathogenic viruses appeared because of reassortment with low pathogenicity viruses). Closed triangles

indicate Egyptian-like virus strains from Russia isolated in 2020. Sequences were obtained from the GISAID EpiFlu database (<https://www.gisaid.org>). Scale bar indicates nucleotide substitutions per site. HPAI, high pathogenicity avian influenza; LPAI, low pathogenicity avian influenza.



Appendix 1 Figure 8. Phylogenetic analysis of matrix protein gene segments of avian influenza viruses. Trees were constructed by using the maximum-likelihood method. Closed circles indicate the highly pathogenic avian influenza H5N1 virus strains isolated from dead Caspian terns on Maliy Zhemchuzhnyi Island, Caspian Sea, Russia in May 2022. Red box indicates virus strains from Israel and Romania closely related to the Caspian Sea strains. Closed triangles indicate Egyptian-like virus strains from Russia isolated in 2020. Sequences were obtained from the GISAID EpiFlu database (<https://www.gisaid.org>). Scale bar indicates nucleotide substitutions per site. HPAI, high pathogenicity avian influenza; LPAI, low pathogenicity avian influenza.



Appendix 1 Figure 9. Phylogenetic analysis of nonstructural protein gene segments of avian influenza viruses. Trees were constructed by using the maximum-likelihood method. Closed circles indicate the highly pathogenic avian influenza H5N1 virus strains isolated from dead Caspian terns on Maliy Zhemchuzhniy Island, Caspian Sea, Russia in May 2022. Red box indicates virus strains from Israel and Romania closely related to the Caspian Sea strains. Closed triangles indicate Egyptian-like virus strains from Russia isolated in 2020. Sequences were obtained from the GISAID EpiFlu database (<https://www.gisaid.org>). Scale bar indicates nucleotide substitutions per site. HPAI, high pathogenicity avian influenza; LPAI, low pathogenicity avian influenza.



Appendix 1 Figure 10. Reconstructive analysis of flight routes according to data from the Russian Bird Ringing Center. Numbers represent different bird species: 1, *Anser anser*; 2, *Cygnus olor*; 3, *Anas platyrhynchos*; 4, *Anas crecca*; 5, *Anas strepera*; 6, *Anas penelope*; 7, *Anas acuta*; 8, *Anas querquedula*; 9, *Anas clypeata*; 10, *Netta rufina*; 11, *Aythya ferina*; 12, *Bucephala clangula*; 13, *Pandion haliaetus*; 14, *Arenaria interpres*; 15, *Himantopus himantopus*; 16, *Calidris ferruginea*; 17, *Calidris alba*; 18, *Larus ichthyaetus*; 19, *Larus minutus*; 20, *Larus fuscus*; 21, *Larus cachinnans*; 22, *Larus canus*; 23, *Larus genei*. Image and descriptions are from (4), licensed under CC BY 4.0 (<https://creativecommons.org/licenses/by/4.0>).