

Replication of Novel Zoonotic-Like Influenza A(H3N8) Virus in Ex Vivo Human Bronchus and Lung

Kenrie P.Y. Hui, John C.W. Ho, Ka-Chun Ng, Samuel M.S. Cheng, Ko-Yung Sit, Timmy W.K. Au, Leo L.M. Poon, John M. Nicholls, Malik Peiris, Michael C.W. Chan

Human infection with avian influenza A(H3N8) virus is uncommon but can lead to acute respiratory distress syndrome. In explant cultures of the human bronchus and lung, novel H3N8 virus showed limited replication efficiency in bronchial and lung tissue but had a higher replication than avian H3N8 virus in lung tissue.

Avian influenza viruses (AIVs) with reassortments between AIVs from domestic poultry and wild birds sporadically cross species barriers, leading to human infections. Viruses with internal genes of H9N2, hemagglutinin, and neuraminidase acquired from wild birds constitute the zoonotic H5N1, H7N9, and H10N8 viruses (1–3) and can lead to severe influenza.

In 2022, two human infections with novel influenza A(H3N8) viruses were reported in Henan and Hunan Province, China (4,5). The first case was identified in a 4-year-old boy with acute respiratory distress syndrome, and the second case occurred in a 5-year-old boy with mild disease. Phylogenetic analysis revealed that the novel H3N8 viruses were triple reassortments containing the Eurasian avian H3 gene of wild-bird origin, the North American avian N8 gene derived from the wild bird AIV, and G57 genotype H9N2 internal genes from AIVs found in poultry in China (6,7). H3N8 viruses that are genetically similar to the zoonotic H3N8 viruses reported in China (4,5) have been isolated in poultry markets in Hong Kong, China (8). Those novel avian H3N8 viruses are

antigenically distant from contemporary human influenza A(H3N2) viruses, and little cross-reactive immunity to these chicken H3N8 viruses exists in the human population (8). We assessed the replication of the novel influenza A(H3N8) virus in human ex vivo bronchus and lung tissues (Appendix, <https://wwwnc.cdc.gov/EID/article/29/6/22-1680-App1.pdf>).

The Study

The viruses used in this study were H9N2/Y280, pH1N1, avH3N8/MP16, novel H3N8, and H5N1/483 (Appendix Table 1). The novel H3N8 virus was isolated from chickens and is genetically closely related to the virus causing zoonotic human disease in China (A/Henan/4-10CNIC/2022/H3N8) (8). Their hemagglutinin genes share a 99.1% similarity, and the neuraminidase genes share a 98.7% similarity. The avH3N8 virus was isolated from wild bird droppings in Mai Po, Hong Kong, and is genetically unrelated to the virus causing zoonotic disease in China. The novel H3N8 virus failed to propagate in Madin-Darby canine kidney (MDCK) cells but could be propagated in eggs and titrated in chicken embryo fibroblasts (DF-1), whereas the other strains could be propagated and titrated in MDCK cells. We performed titration in cells that support the replication of the influenza A viruses rather than in all DF-1 cells, because pH1N1 virus did not replicate in DF-1 cells. We investigated the virus replication kinetics by measuring viral matrix protein segment RNA in culture supernatants using real-time quantitative reverse transcription PCR and 50% tissue culture infectious dose (TCID₅₀) assay for infectious virus titers (Figure 1).

In bronchial tissues, pH1N1 virus had a higher level of viral RNA than did avH3N8, novel H3N8, and H5N1, whereas levels of viral RNA of H9N2 virus were higher than those of avH3N8 and H5N1 virus (Figure 1, panels A, C). The viral RNA levels of avH3N8, novel H3N8, and H5N1 virus were similar.

Author affiliations: The University of Hong Kong School of Public Health, Hong Kong, China (K.P.Y. Hui, J.C.W. Ho, K.-C. Ng, S.M.S. Cheng, L.L.M. Poon, M. Peiris, M.C.W. Chan); Centre for Immunology and Infection, Hong Kong (K.P.Y. Hui, J.C.W. Ho, L.L.M. Poon, M. Peiris, M.C.W. Chan); The University of Hong Kong Division of Cardiothoracic Surgery, Hong Kong (K.-Y. Sit, T.W.K. Au); The University of Hong Kong School of Clinical Medicine, Hong Kong (J.M. Nicholls)

DOI: <https://doi.org/10.3201/eid2906.221680>




Figure 1. Comparative replication competence of zoonotic-like influenza A(H3N8) viruses isolated from chicken and other human and avian viruses in ex vivo cultures of human bronchus and lung tissue. Viral M segment RNA copies (A, B) and viral titers (E, F) in culture supernatants were collected at 1, 24, and 48 hours postinfection with H9N2/Y280, pH1N1, avH3N8/MP16, novel H3N8, or H5N1/483 viruses and measured by quantitative reverse transcription PCR (A, B) and TCID₅₀ (E, F). C, D) Viral load from panels A and B by virus strain. G, H) Viral titers from panels E and F by virus strain. Data are geometric mean \pm SD. Statistical analysis was performed using 2-way (A, B, E, F) or 1-way (C, D, G, H) analysis of variance followed by Tukey posttest; $p < 0.05$ was considered to be statistically significant. Detailed information on viruses used in study is provided in the Appendix (<https://wwwnc.cdc.gov/EID/article/29/6/22-1680-App1.pdf>). AUC, area under the curve; M, matrix; TCID₅₀, 50% tissue culture infectious dose.

The viral RNA level of H5N1 virus was the highest among all the tested strains in human lung tissues, followed by H9N2 (Figure 1, panels B, D). Viral RNA levels of novel H3N8 and pH1N1 viruses were higher than those of avH3N8 virus. Measurement of viral RNA using quantitative reverse transcription PCR is sensitive, but it cannot distinguish defective viral particles from infectious ones. Therefore, we performed the TCID₅₀ assay to monitor the infectious viral titers.

As expected, replication of pH1N1 virus was the highest among all tested strains in human bronchial tissues, in both viral titers and area under the curve values (Figure 1, panels E, G). The titers of H5N1 virus were similar to those of H9N2 virus, whereas H5N1 virus had higher replication competence than did avH3N8 and novel H3N8 viruses. We observed a discrepant trend between viral RNA copies and infectious titers for H9N2. Viral RNA levels of H9N2 were similar to those of pH1N1 (Figure 1, panels A,

C), but the infectious titers of H9N2 virus were significantly lower than that of pH1N1 in bronchus (Figure 1, panels E, G).

In lung tissue, H5N1 virus had the highest replication of all strains tested (Figure 1, panels F, H). Similar to pH1N1 virus, H9N2 had higher titers than the 2 H3N8 viruses in lung tissues. avH3N8 had the lowest titer measured by TCID₅₀. The novel H3N8 virus replicated poorly in mammalian MDCK cells but replicated efficiently in DF-1 avian cells (Appendix Figure). Those findings imply that the novel H3N8 virus has not yet adapted to mammal hosts, which was confirmed by limited replication in human bronchial and lung tissues (Figure 1, panels E, F).

We fixed infected tissues at 48 hours postinfection and stained them for influenza A nucleoprotein immunohistochemistry (Appendix). Consistent with TCID₅₀ findings, bronchial tissues infected with pH1N1 had the most extensive distribution of viral




Figure 2. Tissue tropism of influenza A viruses in ex vivo cultures of human bronchus and lung tissue. Immunohistochemical staining of influenza A nucleoprotein in ex vivo cultures of human bronchial tissues (A) and lung tissues (B) at 48 hours postinfection with H9N2/Y280, pH1N1, H3N8/MP16, novel H3N8, and H5N1/483 viruses. Positive cells are indicated by red-brown color. Images are representative of 3 individual donors. Scale bar indicates 100 μm. Detailed information on viruses used in study is provided in the Appendix (<https://wwwnc.cdc.gov/EID/article/29/6/22-1680-App1.pdf>).

Table. Source of gene segments of novel and avian influenza A(H3N8) viruses*

| Gene segment | Novel H3N8 | avH3N8/MP16 |
|--------------------|------------|-------------|
| Polymerase basic 2 | H9N2 | H3N8 |
| Polymerase basic 1 | H9N2 | H6N1 |
| Polymerase acidic | H9N2 | H6N2 |
| Hemagglutinin | H3N8 | H3N8 |
| Nucleoprotein | H9N2 | H3N8 |
| Neuraminidase | H3N8 | H3N8 |
| Matrix | H9N2 | H1N1 |
| Nonstructural | H9N2 | H7N1 |

*Detailed information on viruses used in study is provided in the Appendix (<https://wwwnc.cdc.gov/EID/article/29/6/22-1680-App1.pdf>).

antigen, whereas we observed moderate levels of viral antigen staining in tissues infected with novel H3N8, H9N2, and H5N1 virus. No viral antigen staining was observed in tissues infected with avH3N8 (Figure 2, panel A). In the lung, H5N1 virus infection demonstrated the most extensive viral antigen staining, followed by infection with pH1N1, novel H3N8, and avH3N8 virus, which demonstrated the least extensive staining (Figure 2, panel B).

The discrepancy between the viral load in RNA copies and TCID₅₀ titers of H9N2 and avH3N8 infection suggests that infection with those viruses might produce high levels of defective particles that cannot be detected by TCID₅₀ assay. Immunohistochemistry staining of viral antigen serves as alternative evidence of virus replication in human tissues. The staining correlates more with TCID₅₀ results than with viral RNA analysis for all the viruses.

Amino acid comparisons of the novel H3N8 and avH3N8 viruses demonstrated that they shared the same stalk length in the NA gene but did not have the G228S mutation that enhances binding to mammalian receptors (Appendix Table 2). The internal genes of the novel H3N8 virus were reassorted from H9N2 virus, whereas the internal genes of the avH3N8 came from H3N8, H6N1, H6N2, H3N8, H1N1, and H7N1 (Table). Neither virus had the E627K mutation in polymerase basic 2 that confers mammal adaptation, virulence, and transmissibility. The novel H3N8 virus had the A588V mutation in polymerase basic 2 that promotes mammal adaptation, but avian H3N8 virus did not have this mutation. This difference might contribute to higher replication of the novel H3N8 virus in human lung tissue. The S31N mutation found in the matrix protein 2 of the novel H3N8 virus provided adamantane resistance.

Conclusions

Although zoonotic H3N8 viruses have a dual receptor-binding affinity of α -2,3 and α -2,6 receptors (7), our findings show that this factor does not confer an advantage for replication in human bronchial tissue.

Our findings demonstrated inefficient replication of the novel H3N8 virus in human bronchial tissues, which implies limited efficiency to transmit among humans. This finding is in line with a recent serologic surveillance study in which no poultry workers were positive for antibodies for the novel H3N8 virus (7), and only 2 human cases have been documented since April 2022 (4,5). The moderate replication ability of the novel H3N8 virus in human lung tissue suggests that the virus causes less severe disease than H5N1 virus.

In summary, our findings suggest that the zoonotic-like avian H3N8 virus has limited efficiency for human-to-human transmission and, at present, is unlikely to cause severe disease in humans. However, the limited cross-reactive immunity against this novel H3N8 virus in the human population (8) and the emergence of novel H3N8 viruses by continuous reassortment between AIVs in wild birds and poultry demonstrate that the zoonotic and pandemic potential of avian H3N8 viruses should be closely monitored.

Acknowledgments

We thank Jenny Chan, Rachel Ching, and Kevin Fung for technical support.

This work was supported by grants from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and Human Services (contract no. 75N93021C00016) and the Theme-Based Research Scheme (Ref: T11-712/19-N) under University Grants Committee of Hong Kong Special Administrative Region.

About the Author

Dr. Hui is an assistant professor in the School of Public Health at the University of Hong Kong. Her research interests include risk assessment, understanding the pathogenesis of emerging respiratory viruses, and the development of therapeutic options for severe influenza diseases and coronavirus infections.

References

- Guan Y, Shortridge KF, Krauss S, Webster RG. Molecular characterization of H9N2 influenza viruses: were they the donors of the "internal" genes of H5N1 viruses in Hong Kong? *Proc Natl Acad Sci U S A*. 1999;96:9363-7. <https://doi.org/10.1073/pnas.96.16.9363>
- Lam TT, Zhou B, Wang J, Chai Y, Shen Y, Chen X, et al. Dissemination, divergence and establishment of H7N9 influenza viruses in China. *Nature*. 2015;522:102-5. <https://doi.org/10.1038/nature14348>
- Ma C, Lam TT, Chai Y, Wang J, Fan X, Hong W, et al. Emergence and evolution of H10 subtype influenza viruses in


- poultry in China. *J Virol.* 2015;89:3534–41. <https://doi.org/10.1128/JVI.03167-14>
4. Cheng D, Dong Y, Wen S, Shi C. A child with acute respiratory distress syndrome caused by avian influenza H3N8 virus. *J Infect.* 2022;85:174–211. <https://doi.org/10.1016/j.jinf.2022.05.007>
 5. Tan X, Yan X, Liu Y, Wu Y, Liu JY, Mu M, et al. A case of human infection by H3N8 influenza virus. *Emerg Microbes Infect.* 2022;11:2214–7. <https://doi.org/10.1080/22221751.2022.2117097>
 6. Li Y, Li P, Xi J, Yang J, Wu H, Zhang Y, et al. Wild bird-origin H3N8 avian influenza virus exhibit well adaptation in mammalian host. *J Infect.* 2022;84:579–613. <https://doi.org/10.1016/j.jinf.2021.12.014>
 7. Yang R, Sun H, Gao F, Luo K, Huang Z, Tong Q, et al. Human infection of avian influenza A H3N8 virus and the viral origins: a descriptive study. *Lancet Microbe.* 2022;3:e824–34. [https://doi.org/10.1016/S2666-5247\(22\)00192-6](https://doi.org/10.1016/S2666-5247(22)00192-6)
 8. Sit THC, Sun W, Tse ACN, Brackman CJ, Cheng SMS, Tang AWY, et al. Novel zoonotic avian influenza A(H3N8) virus in chicken, Hong Kong, China. *Emerg Infect Dis.* 2022;28:2009–15. <https://doi.org/10.3201/eid2810.221067>

Address for correspondence: Michael C.W. Chan, School of Public Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pok Fu Lam, Hong Kong, China; email: mchan@hku.hk

March 2023

World TB Day

- Risk for Prison-to-Community Tuberculosis Transmission, Thailand, 2017–2020
- Multicenter Retrospective Study of Vascular Infections and Endocarditis Caused by *Campylobacter* spp., France
- Yellow Fever Vaccine–Associated Viscerotropic Disease among Siblings, São Paulo State, Brazil
- *Bartonella* spp. Infections Identified by Molecular Methods, United States
- COVID-19 Test Allocation Strategy to Mitigate SARS-CoV-2 Infections across School Districts
- Using Discarded Facial Tissues to Monitor and Diagnose Viral Respiratory Infections
- Postacute Sequelae of SARS-CoV-2 in University Setting
- Associations of *Anaplasma phagocytophilum* Bacteria Variants in *Ixodes scapularis* Ticks and Humans, New York, USA
- Prevalence of *Mycobacterium tuberculosis* Complex among Wild Rhesus Macaques and 2 Subspecies of Long-Tailed Macaques, Thailand, 2018–2022
- Increase in Colorado Tick Fever Virus Disease Cases and Effect of COVID-19 Pandemic on Behaviors and Testing Practices, Montana, 2020
- Comparative Effectiveness of COVID-19 Vaccines in Preventing Infections and Disease Progression from SARS-CoV-2 Omicron BA.5 and BA.2, Portugal
- Clonal Dissemination of Antifungal-Resistant *Candida haemulonii*, China



- Correlation of High Seawater Temperature with *Vibrio* and *Shewanella* Infections, Denmark, 2010–2018
- Tuberculosis Preventive Therapy among Persons Living with HIV, Uganda, 2016–2022
- Nosocomial Severe Fever with Thrombocytopenia Syndrome in Companion Animals, Japan, 2022
- *Burkholderia thailandensis* Isolated from the Environment, United States
- *Mycobacterium leprae* in Armadillo Tissues from Museum Collections, United States
- Reemergence of Lymphocytic Choriomeningitis Mammarenavirus, Germany
- *Emergomyces pasteurianus* in Man Returning to the United States from Liberia and Review of the Literature
- New Detection of Locally Acquired Japanese Encephalitis Virus Using Clinical Metagenomics, New South Wales, Australia
- Recurrent Cellulitis Revealing *Helicobacter cinaedi* in Patient on Ibrutinib Therapy, France
- *Inquilinus limosus* Bacteremia in Lung Transplant Recipient after SARS-CoV-2 Infection
- Sustained Mpox Proctitis with Primary Syphilis and HIV Seroconversion, Australia
- SARS-CoV-2 Infection in a Hippopotamus, Hanoi, Vietnam
- Clonal Expansion of Multidrug-Resistant *Streptococcus dysgalactiae* Subspecies *equisimilis* Causing Bacteremia, Japan, 2005–2021
- Extended Viral Shedding of MERS-CoV Clade B Virus in Llamas Compared with African Clade C Strain
- Seroprevalence of Specific SARS-CoV-2 Antibodies during Omicron BA.5 Wave, Portugal, April–June 2022
- SARS-CoV-2 Incubation Period during the Omicron BA.5–Dominant Period in Japan
- Risk Factors for Reinfection with SARS-CoV-2 Omicron Variant among Previously Infected Frontline Workers
- Genomic Analysis of Early Monkeypox Virus Outbreak Strains, Washington, USA

**EMERGING
INFECTIOUS DISEASES**

To revisit the March 2023 issue, go to:
<https://wwwnc.cdc.gov/eid/articles/issue/29/3/table-of-contents>

Article DOI: <https://doi.org/10.3201/eid2906.221680>

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

Replication of Novel Zoonotic-Like Influenza A(H3N8) Virus in Ex Vivo Human Bronchus and Lung

Appendix

Materials and Methods

Viruses

In addition to the zoonotic-like influenza H3N8 (A/Env/Hong_Kong/MKT_AB_13cp/2022, novel H3N8) which is a chicken virus, an avian virus isolated from wild bird droppings in Mai Po, Hong Kong (A/Env/Hong Kong/MP16_265/2016, avH3N8/MP16), a highly pathogenic avian influenza (HPAI) H5N1 virus (A/Hong Kong/483/1997, H5N1/483) isolated from a fatal human infection, a 2009 pandemic influenza virus (A/Hong Kong/415742/2009, pH1N1) isolated from a patient in Hong Kong and a duck H9N2 virus (A/Duck/Hong Kong/Y280/97, H9N2/Y280) were used as controls. The novel H3N8 virus was propagated in eggs while all the other influenza viruses were passaged in Madin-Darby canine kidney (MDCK) cells. Virus stock was aliquoted and titrated to determine the plaque forming unit per mL (pfu/mL) in chicken embryo fibroblast (DF-1) cells for the novel H3N8 virus and MDCK cells for the other influenza viruses. The experiments were carried out in a Bio-safety level 3 (BSL-3) facility at the School of Public Health, LKS Faculty of Medicine, The University of Hong Kong.

Ex vivo cultures and infection of human bronchus and lung

Fresh non-tumor bronchus and lung tissues were obtained from patients undergoing elective surgery in Department of Surgery of Queen Mary Hospital and were removed as part of clinical care but surplus for routine diagnostic requirements as detailed previously (1). Sampling

of tissues is defined by convenience. Fragments of human tissues were infected with each virus at 1×10^6 pfu/mL for 1 h at 37°C. Mock-infected tissue served as negative controls. The explants were washed three times with PBS and placed in culture medium (F-12K nutrient mixture with L-glutamine, and antibiotics) with or without a sterile surgical pathology sponge to establish an air-liquid interface condition in 24-well culture plates in a 37°C incubator with 5% CO₂. Infectious viral titers in culture supernatants were assessed at 1, 24, and 48 hours post-infection (hpi) by TCID₅₀ assay. Bronchus and lung tissues were fixed at 48 hpi in 10% formalin and processed for immunohistochemistry staining. Ethics approval of the use of human tissues was granted by the institutional review board of the University of Hong Kong and the hospital authority (approval number UW 20–862).

Viral titration by TCID₅₀ assay

A confluent 96-well tissue culture plate of DF-1 or MDCK cells was prepared 1 day before the virus titration (TCID₅₀) assay. Cells were washed once with PBS and replenished serum-free MEM medium supplemented with 100 units/ml penicillin and 100 µg/ml streptomycin and 2 µg/ml of tosylsulfonyl phenylalanylchloromethyl ketone treated trypsin for DF-1 and MDCK cells. Serial dilutions of virus supernatant, from 0.5 log to 7 log, were performed before adding the virus dilutions onto the plates in quadruplicate. The plates were observed for cytopathic effect daily. The endpoint of viral dilution leading to CPE in 50% of inoculated wells was estimated using the Karber method (2). AUC was calculated from the viral titers from different time points indicated in the y-axis.

Quantitative reverse transcription PCR assay

The viral RNA in 20 µL culture supernatants was extracted using the QIAamp Viral RNA Mini kit (Qiagen). RNA was reverse-transcribed by using random 6-mer primers with PrimeScript RT reagent Kit (Takara). Viral RNA copy number of target genes was performed using an ABI ViiA 7 real-time PCR system (Applied Biosystems). All procedures were performed according to the manufacturers' instructions as previously described (3,4).

Immunohistochemical staining

The human tissues were fixed with 10% formalin overnight, embedded in paraffin blocks and stained for influenza viral protein as previously described (4). Briefly, the paraffin-embedded tissues were sectioned, deparaffinised, digested with pronase and blocked with an


avidin/biotin blocking kit (Vector Labs, Burlingame, CA, USA). The sections were then stained with influenza A virus nucleoprotein-specific mouse monoclonal antibody HB65 (EVL, Woerden, The Netherlands) and a biotinylated secondary antibody. The bound antibodies were visualized with a Strep-ABC complex and an AEC substrate kit.

Statistical analysis

Experiments with the human *ex vivo* cultures were performed independently with five and six different donors of bronchus and lung, respectively. Results shown in figures are geometric mean (+/-SD). Area-under-the-curve (AUC) was calculated by integrating infectious virus titers at 24–48 hpi in *ex vivo* bronchus, lung tissues. The differences in log₁₀-transformed viral titers and quantitative viral RNA of M segment between viruses and over time were compared using two-way ANOVA followed by a Tukey's multiple-comparison test using GraphPad Prism version 9.0.0. Comparison of AUC between viruses was calculated using one-way ANOVA followed by a Tukey's multiple-comparison test. Differences were considered significant at a *P* value less than 0.05.

References

1. Hui KP, Chan LL, Kuok DI, Mok CK, Yang ZF, Li RF, et al. Tropism and innate host responses of influenza A/H5N6 virus: an analysis of *ex vivo* and *in vitro* cultures of the human respiratory tract. *Eur Respir J*. 2017;49:1601710. [PubMed https://doi.org/10.1183/13993003.01710-2016](https://doi.org/10.1183/13993003.01710-2016)
2. Kärber G. Contribution to the collective treatment of pharmacological experimental series [in German]. *Arch. Exp. Pathol. Pharmacol*. 1931;162:480–3. [PubMed https://doi.org/10.1007/BF01863914](https://doi.org/10.1007/BF01863914)
3. Hui KPY, Ho JCW, Cheung MC, Ng KC, Ching RHH, Lai KL, et al. SARS-CoV-2 Omicron variant replication in human bronchus and lung *ex vivo*. *Nature*. 2022;603:715–20. [PubMed https://doi.org/10.1038/s41586-022-04479-6](https://doi.org/10.1038/s41586-022-04479-6)
4. Hui KP, Chan LL, Kuok DI, Mok CK, Yang ZF, Li RF, et al. Tropism and innate host responses of influenza A/H5N6 virus: an analysis of *ex vivo* and *in vitro* cultures of the human respiratory tract. *Eur Respir J*. 2017;49:1601710. [PubMed https://doi.org/10.1183/13993003.01710-2016](https://doi.org/10.1183/13993003.01710-2016)



Appendix Figure. Novel H3N8 virus replication in DF-1 cells infected at MOI 0.01. TCID₅₀ assay was performed in DF-1 cells. Dotted line represents the detection limit of TCID₅₀ assay.

Appendix Table 1. Influenza A virus strains used

| Virus | Strain name |
|-------------|----------------------------------|
| H9N2/Y280 | A/Duck/Hong Kong/Y280/97 |
| pH1N1 | A/Hong Kong/415742/2009 |
| avH3N8/MP16 | A/Env/Hong Kong/MP16_265/2016 |
| Novel H3N8 | A/Env/Hong_Kong/MKT_AB_13cp/2022 |
| H5N1/483 | A/Hong Kong/483/1997 |

Appendix Table 2. Amino acid comparison between the novel H3N8 (AB_13) and avian H3N8 (MP16)

| | amino acid position | 2 | 6 | 8 | 19 | 45 | 57 | 58 | 70 | 83 | 92 | 102 | 112 | 119 | 124 | 128 | 131 | 137 | 145 | 159 | |
|---|---------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|
| HA1 | | | | | | | | | | | | | | | | | | | | | |
| A/Env/Hong_Kong/MKT_AB_13cp/2022 | | N | K | S | S | N | K | V | M | K | S | V | I | D | S | A | S | G | N | S | |
| A/Env/Hong_Kong/MP16_265/2016_(H3N8)_Anas_acuta | | D | N | N | A | S | R | I | L | T | N | I | V | E | G | T | T | S | S | N | |
| | amino acid position | 160 | 163 | 172 | 214 | 228 | 257 | 260 | 264 | 275 | 312 | | | | | | | | | | |
| | | S | L | Y | V | G | F | I | R | E | N | | | | | | | | | | |
| | | A | V | D | I | G | Y | M | K | D | T | | | | | | | | | | |
| NA | | | | | | | | | | | | | | | | | | | | | |
| | amino acid position | 14 | 20 | 22 | 26 | 30 | 37 | 38 | 39 | 43 | 44 | 49 | 50 | 59 | 61 | 62 | 69 | 71 | 74 | 75 | |
| A/Env/Hong_Kong/MKT_AB_13cp/2022 | | V | L | I | I | T | P | R | D | R | N | V | I | I | K | V | N | I | I | E | |
| A/Env/Hong_Kong/MP16_265/2016_(H3N8)_Anas_acuta | | I | F | V | V | I | G | K | G | G | V | I | V | V | R | I | S | V | M | P | |
| | amino acid position | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 89 | 92 | 125 | 154 | 161 | 191 | 198 | 209 | 235 | 237 | 253 | 254 | |
| | | R | P | E | N | D | H | F | L | A | T | Y | I | I | A | I | E | F | A | F | |
| | | Y | W | K | E | G | T | Y | i | V | I | F | V | V | S | V | D | Y | I | Y | |
| | amino acid position | 257 | 260 | 263 | 265 | 266 | 267 | 283 | 311 | 329 | 355 | 375 | 376 | 377 | 381 | 383 | 388 | 389 | 392 | 397 | |
| | | K | K | E | A | E | I | E | R | S | N | K | V | R | V | N | I | K | V | S | |
| | | S | R | G | R | D | V | D | K | A | T | R | I | K | T | T | V | R | I | L | |
| | amino acid position | 410 | 414 | 415 | 416 | 452 | 469 | | | | | | | | | | | | | | |
| | | V | K | R | N | I | E | | | | | | | | | | | | | | |
| | | I | G | K | D | V | K | | | | | | | | | | | | | | |
| PB2 | | | | | | | | | | | | | | | | | | | | | |
| | amino acid position | 588 | 627 | | | | | | | | | | | | | | | | | | |
| A/Env/Hong_Kong/MKT_AB_13cp/2022 | | V | E | | | | | | | | | | | | | | | | | | |
| A/Env/Hong_Kong/MP16_265/2016_(H3N8)_Anas_acuta | | A | E | | | | | | | | | | | | | | | | | | |
| M2 | | | | | | | | | | | | | | | | | | | | | |
| | amino acid position | 31 | | | | | | | | | | | | | | | | | | | |
| A/Env/Hong_Kong/MKT_AB_13cp/2022 | | N | | | | | | | | | | | | | | | | | | | |
| A/Env/Hong_Kong/MP16_265/2016_(H3N8)_Anas_acuta | | S | | | | | | | | | | | | | | | | | | | |