

- assay. *Microbiol Spectr*. 2021;9:e0046221. <https://doi.org/10.1128/Spectrum.00462-21>
6. Okuma K, Iwakawa K, Turnidge JD, Grubb WB, Bell JM, O'Brien FG, et al. Dissemination of new methicillin-resistant *Staphylococcus aureus* clones in the community. *J Clin Microbiol*. 2002;40:4289-94. <https://doi.org/10.1128/JCM.40.11.4289-4294.2002>
 7. Durand G, Javerliat F, Bes M, Veyrieras JB, Guignon G, Mugnier N, et al. Routine whole-genome sequencing for outbreak investigations of *Staphylococcus aureus* in a national reference center. *Front Microbiol*. 2018;9:511. <https://doi.org/10.3389/fmicb.2018.00511>
 8. Ito T, Ma XX, Takeuchi F, Okuma K, Yuzawa H, Hiramatsu K. Novel type V staphylococcal cassette chromosome mec driven by a novel cassette chromosome recombinase, ccrC. *Antimicrob Agents Chemother*. 2004;48:2637-51. <https://doi.org/10.1128/AAC.48.7.2637-2651.2004>
 9. Herold BC, Immergluck LC, Maranan MC, Lauderdale DS, Gaskin RE, Boyle-Vavra S, et al. Community-acquired methicillin-resistant *Staphylococcus aureus* in children with no identified predisposing risk. *JAMA*. 1998;279:593-8. <https://doi.org/10.1001/jama.279.8.593>
 10. Tomic V, Svetina Sorli P, Trinka D, Sorli J, Widmer AF, Trampuz A. Comprehensive strategy to prevent nosocomial spread of methicillin-resistant *Staphylococcus aureus* in a highly endemic setting. *Arch Intern Med*. 2004;164:2038-43. <https://doi.org/10.1001/archinte.164.18.2038>

Address for correspondence: Stéphane Corvec, Institut de Biologie des Hôpitaux de Nantes, Service de Bactériologie et des contrôles Microbiologiques, CHU de Nantes, 9 quai Moncoussu, 44093 Nantes CEDEX 01, France; email: stephane.corvec@chu-nantes.fr

Inadvertent Platelet Transfusion from Monkeypox Virus–Infected Donor to Recipient, Thailand, 2023

Jiratchaya Puenpa, Duangnapa Intharasongkroh, Sompong Vongpunsawad, Dootchai Chaiwanichsiri, Yong Poovorawan

Author affiliations: Center of Excellence in Clinical Virology, Chulalongkorn University Faculty of Medicine, Bangkok, Thailand (J. Puenpa, S. Vongpunsawad, Y. Poovorawan); National Blood Center, Thai Red Cross Society, Bangkok (D. Intharasongkroh, D. Chaiwanichsiri); FRS(T), The Royal Society of Thailand, Sanam Sueapa, Dusit, Bangkok (Y. Poovorawan)

In Thailand, platelet product from a blood donor was transfused to a recipient who had dengue. Two days later, the donor was confirmed to have monkeypox virus infection. Monkeypox virus DNA was undetectable in recipient specimens up to 2 weeks after transfusion. The recipient remained asymptomatic at 4 weeks of monitoring.

DOI: <http://doi.org/10.3201/eid3003.231539>

Monkeypox virus (MPXV), a double-stranded DNA virus that primarily infects rodents in sub-Saharan Africa, causes mpox disease. MPXV is a member of the genus *Orthopoxvirus* in the family *Poxviridae*. MPXV clade I is endemic to Central Africa and clade II to West Africa. Clade II is further subdivided into IIa and IIb. Strains from the recent global emergence appear to belong to clade IIb (<https://nextstrain.org/mpox/all-clades>).

The potential to unknowingly transmit MPXV from donated blood products exists despite routine stringent screening of bloodborne pathogens at donation centers. Thailand first reported mpox in a 27-year-old male tourist from Africa in Phuket province on July 21, 2022; nonoutbreak sporadic infections have since been identified (1). By May 2023, ~40 infections had been laboratory-confirmed. Infections surged after Pride Festivals, which took place in Bangkok and Pattaya City in June 2023; infections peaked in August and then declined. As of November 4, 2023, the Ministry of Public Health Thailand (MoPH) had identified 582 infections (563 male and 19 female patients; median age 33 years, age range 1–64 years) and 2 deaths. Here, we describe an unintended administration of platelets from an MPXV-infected donor to a dengue-infected recipient and the subsequent follow-up to monitor for potential MPXV transmission.

On July 24, 2023, an apparently healthy 22-year-old man donated whole blood at the National Blood Center (NBC) of the Thai Red Cross in Bangkok (Figure). That afternoon, he experienced fever and malaise. On July 26, itchy skin rash and lesions appeared on his hands, feet, and anus, which prompted him to go to a hospital. His doctor sought consultation with the Department of Disease Control at MoPH, where samples of the skin lesion, oropharyngeal swab, and plasma were tested for MPXV by real-time PCR to detect the F3L gene region (BioPerfectus, <https://www.bioperfectus.com>). MPXV DNA was detected only in the lesion (cycle threshold [Ct] 21.7) and oropharyngeal (Ct 31.5) swab samples.

NBC processes blood donations individually and routinely screened for hepatitis B/C and syphilis. Derived products from donations are primarily leukocyte-poor red cells, leukocyte-depleted pooled plate-




Figure. Timeline of MPXV-infected blood donor (red) and platelet recipient (blue), Thailand, 2023. MPXV, monkeypox virus.

let concentrate, and fresh frozen plasma, prepared in accordance with guidelines of the European Directorate for the Quality of Medicines & Healthcare (2). Specifically, the platelet concentrate is prepared from a pool of 4 donor buffy coats of the same ABO blood group, diluted with either plasma from one of the buffy coat donations or a platelet additive solution, centrifuged to separate the platelets, filtered to deplete leukocyte, and stored for bacterial testing before distribution.

On July 31, the NBC was alerted to the potential of an MPXV-contaminated donation, which prompted recalls of all blood components derived from the 22-year-old donor. That same day, red blood cells and plasma derived from the donor materials were successfully retrieved and destroyed; however, the platelet concentrate had already been administered to an 11-year-old female recipient who had ongoing dengue infection.

To characterize MPXV in the donation, our laboratory received residual donor plasma and red cells that the NBC had, from which we extracted DNA by using the magLEAD 12 gC instrument (Precision System Science, <https://www.pss.co.jp>) according to the manufacturer's instructions. We tested for MPXV DNA by generic real-time PCR to detect the tumor necrosis

factor receptor gene located at the terminal inverted repeat region on the MPXV genome, in accordance with the US Centers for Disease Control and Prevention protocol (3). We confirmed the result using conventional PCR to amplify the DNA helicase and Schlafen protein genes (Appendix, <https://wwwnc.cdc.gov/EID/article/30/3/23-1539-App1.pdf>). We Sanger sequenced amplicons, and deposited nucleotides into GenBank (accession nos. OR790439–40).

Plasma yielded detectable MPXV DNA ($C_t \approx 35$); red blood cells did not. Phylogenetic analysis of the DNA helicase gene sequence suggests that the MPXV strain in the donor belonged to clade IIb (lineage B) and genetically clustered with strains previously identified in Taiwan, Japan, and the United States (88% bootstrap support) (Appendix Figure).

MPXV DNA was undetectable in serum and throat swab samples collected from the platelet recipient on August 1, 3, 7, and 14. No mpox-associated symptoms were evident 4 weeks posttransfusion. Incubation period for mpox is 3–17 days (mean 8.5 days) (4,5).

We posit that there was a low risk for transfusion-transmitted infection for several reasons. First, detection of MPXV DNA in the residual donated plasma does not indicate infectious virus, as was shown

in a viral load study using cell culture as surrogate for infectivity (6). Thus, nucleic acid detection does not prove the presence of viable or infectious virus, as Cohen et al. demonstrated in a smallpox-vaccine study (7). We pooled and extensively prepared platelet products from multiple donors, which may have diluted out any residual virus before transfusion 1 week later. In conclusion, our study shows that a blood donation from a donor with detectable MPXV viral DNA did not appear to transmit the infection to a pooled-platelet recipient.

This work was supported by the Health Systems Research Institute, the National Research Council of Thailand, the Center of Excellence in Clinical Virology, Chulalongkorn University, King Chulalongkorn Memorial Hospital, the MK Restaurant Group and Aunt Thongkam Foundation, and the BJC Big C Foundation. J.P. reports financial support from the Second Century Fund Fellowship of Chulalongkorn University.

About the Author

Dr. Puenpa is a postdoctoral fellow at the Center of Excellence in Clinical Virology in the Faculty of Medicine at Chulalongkorn University. Her primary research interests are molecular epidemiology and evolution of human enteroviruses.

Reference

1. Ministry of Public Health Thailand (MoPH). Thailand's first monkeypox case identified in Phuket tourist [in Thai]. 2022 Jul 22 [cited 2023 Nov 1]. <https://pr.moph.go.th/?url=pr/detail/2/04/176575>
2. European Directorate for the Quality of Medicines & HealthCare. Guide to the preparation, use and quality assurance of blood components. 21st edition [cited 2023 Oct 11]. https://freepub.edqm.eu/publications/AUTOPUB_48/detail
3. Li Y, Zhao H, Wilkins K, Hughes C, Damon IK. Real-time PCR assays for the specific detection of monkeypox virus West African and Congo Basin strain DNA. *J Virol Methods*. 2010;169:223–7. <https://doi.org/10.1016/j.jviromet.2010.07.012>
4. Centers for Disease Control and Prevention. Clinical recognition [cited 2023 Oct 24]. <https://www.cdc.gov/poxvirus/mpox/clinicians/clinical-recognition.html>
5. Miura F, van Ewijk CE, Backer JA, Xiridou M, Franz E, Op de Coul E, et al. Estimated incubation period for monkeypox cases confirmed in the Netherlands, May 2022. *Euro Surveill*. 2022;27:2200448. <https://doi.org/10.2807/1560-7917.ES.2022.27.22.2200448>
6. Lim CK, McKenzie C, Deerain J, Chow EPF, Towns J, Chen MY, et al. Correlation between monkeypox viral load and infectious virus in clinical specimens. *J Clin Virol*. 2023;161:105421. <https://doi.org/10.1016/j.jcv.2023.105421>
7. Cohen JL, Hohman P, Preuss JC, Li L, Fischer SH, Fedorko DP. Detection of vaccinia virus DNA, but not infectious virus, in the blood of smallpox vaccine recipients. *Vaccine*. 2007;25:4571–4. <https://doi.org/10.1016/j.vaccine.2007.03.044>

Address for correspondence: Yong Poovorawan, Center of Excellence in Clinical Virology, Faculty of Medicine, Chulalongkorn University, 1873 Rama 4 Rd, Pathumwan, Bangkok 10330, Thailand; email: yong.p@chula.ac.th

Detection of Invasive *Anopheles stephensi* Mosquitoes through Molecular Surveillance, Ghana

Yaw A. Afrane, Anisa Abdulai, Abdul R. Mohammed, Yaw Akuamoah-Boateng, Christopher M. Owusu-Asenso, Isaac K. Sraku, Stephina A. Yanney, Keziah Malm, Neil F. Lobo.

Author affiliations: University of Ghana, Accra, Ghana (Y.A. Afrane, A. Abdulai, A.R. Mohammed, Y. Akuamoah-Boateng, C.M. Owusu-Asenso, I.K. Sraku, S.A. Yanney); Ghana Health Service, Accra (K. Malm); University of Notre Dame, Notre Dame, Indiana, USA (N.F. Lobo)

DOI: <https://doi.org/10.3201/eid3003.231638>

The invasive *Anopheles stephensi* mosquito has rapidly expanded in range in Africa over the past decade. Consistent with World Health Organization guidelines, routine entomologic surveillance of malaria vectors in Accra, Ghana, now includes morphologic and molecular surveillance of *An. stephensi* mosquitoes. We report detection of *An. stephensi* mosquitoes in Ghana.

Anopheles stephensi is an invasive mosquito species originating from parts of Southeast Asia and the Arabian Peninsula (1). Over the past decade, *An. stephensi* mosquitoes have been expanding in range and have now been documented in several countries in Africa (2). First detected in Djibouti, on the Horn of Africa, in 2012, this vector has been implicated in urban malaria outbreaks (3). They were also detected in Ethiopia in 2016 and 2018 (4,5). *An. stephensi* mosquitoes were subsequently detected in Sudan (2016), Somalia (2019), Nigeria (2020), and Kenya (2023) (2,3,5–7). This invasive vector poses a major threat to current malaria control and elimination efforts. The ability of *An. stephensi* mosquitoes to breed in artificial containers enables them to thrive in urban areas, setting them apart from other major

Inadvertent Platelet Transfusion from Mpox-Infected Donor to Recipient, Thailand, 2023

Appendix

Gene Amplification and DNA Sequencing

The DNA helicase and Schlafen protein gene sequences were amplified using semi-nested RT-PCR with the primers specified in Appendix Table, resulting in amplicon lengths of 315 bp and 314 bp, respectively. The first-round PCR were performed using PerfectTaq MasterMix PCR system, according to the manufacturer's instructions (5 PRIME, Darmstadt, Germany). The amplification conditions for PCR involved 40 cycles with the following parameters: denaturation at 94 °C for 30 seconds, annealing at 50 °C for 30 seconds, and extension at 68 °C for 1 minute 45 seconds. A final extension step at 68 °C for 5 minutes was included. In the second-round PCR reaction, 1 µL of the first-round reaction served as a template, using second-round primers and PerfectTaq MasterMix (5 PRIME, Darmstadt, Germany) in accordance with the manufacturer's instructions. Subsequently, 40 cycles were conducted with the following conditions: denaturation at 94 °C for 30 seconds, annealing at 50 °C for 30 seconds, and extension at 72 °C for 90 seconds, followed by a final extension at 72 °C for 5 minutes. The sequencing and product amplification were performed simultaneously in both the forward and reverse directions at First BASE Laboratories Sdn Bhd (Selangor Darul Ehsan, Malaysia).

Appendix Table. Primers used for conventional real-time PCR assays

Target region	Primers	Sequence (5'-3')	Position	Strand
DNA helicase	F129088	CACTCCAGCACCCGCAGAG	129088-129107	Sense
	F129182	ATGTCACTATTAAAGATGGAGTAT	129182-129206	Sense
	R129496	AGTGAAGAGTGATGTATAGAGG	129474-129496	Antisense
Schlafen protein	F167584	TTACTACTGTAGACGTGCATGG	167584-167606	Sense
	F167644	TGCTTCCGATTCCAATCTGG	167644-167665	Sense
	R167957	AAGACATGCTCCCATAGTCTTC	167935-167957	Antisense



Appendix Figure. Phylogenetic analysis of the DNA helicase gene region (position 123,565–123,898 nt). The phylogenetic tree was constructed using the neighbor-joining method with 1,000 bootstrap replicates and implemented in MEGA version 7 (www.megasoftware.net). Evolutionary distances were computed using the maximum composite likelihood method. Bootstrap values >75 are shown. Scale bar represents substitutions per site.