

2020 NATIONAL HOSPITAL CARE SURVEY (NHCS) TECHNICAL DOCUMENTATION

For Public Use Data Files



**Division of Health Care Statistics
National Center for Health Statistics
February 2024**

Overview Summary

This document provides detailed information and guidance for users of the 2020 National Hospital Care Survey (NHCS) public use data files. As a principal source of information on health care utilization in the United States, NHCS collects inpatient and emergency department (ED) data from a nationally representative sample of U.S. hospitals sourced from administrative and electronic health records data. The 2020 NHCS is conducted by the National Center for Health Statistics (NCHS) and is a member of the National Health Care Surveys – a family of surveys which measure health care utilization across a variety of health care providers and settings.

Section 1 of this document includes information on the scope of the survey, its sources of data, and the confidentiality protections related to the data. Section 2 contains details on the sampling process, data collection procedures, and weighting methodology used to produce national estimates on hospital utilization. Section 3 provides information on the number of sampled hospitals that were eligible to participate in NHCS and submitted data in 2020. Section 4 details the contents of the 2020 NHCS public use data files and the edits used in the creation of the files. Section 5 contains an explanation of the procedures used to accurately produce variance estimates. National Center for Health Statistics presentation standards for proportions, counts, and rates, and their relation to NHCS data are discussed in Section 6, and the data analysis guidelines are provided in Section 7. Section 8 provides variable missingness by both inpatient and ED settings, and Section 9 provides a comparison of frequencies between the NHCS public use data files and NHCS restricted use data file, as well as to the Agency for Healthcare Research and Quality's Healthcare Cost and Utilization Project (HCUP) nationwide samples. Section 10 provides further information on the availability of the NHCS restricted use data files available in NCHS and Federal Research Data Centers. Section 11 provides a list of preferred reporting items for complex sample survey analysis for 2020 NHCS.

Suggested Citation

Technical Documentation: National Center for Health Statistics. Division of Health Care Statistics. *2020 National Hospital Care Survey (NHCS) Public Use Data Files Documentation*, December 2023. Hyattsville, Maryland.

Data File: National Center for Health Statistics. Division of Health Care Statistics. 2020 National Hospital Care Survey (NHCS) public use data files. 2023. Hyattsville, Maryland.

Contact Information

Data users can find the latest information about NHCS on our website, at:

<https://www.cdc.gov/nchs/nhcs/index.htm>. If data users have queries about the public use data files, they may send their question through email to nhcs@cdc.gov, or call us at 301-458-4321. A response to data user inquiries is generally provided in 1-2 business days.

The NHCS also has a listserv, where updates and information about the most recent NHCS news and data releases are sent out. To join the listserv, there are two options:

1. Send an email message to listserv@cdc.gov. Leave the subject line blank. In the body of the message, type or paste: **subscribe NHCS-DATA your name** where “your name” is your first and last name.
2. Fill out and submit a subscription request form through the NCHS listservs webpage (https://www.cdc.gov/nchs/products/nchs_listservs.htm). You should receive an email notification of your subscription.

Contents

Section 1 About the National Hospital Care Survey	6
Section 1.1 Background	6
Section 1.2 Data Sources	6
Section 1.3 Data Confidentiality	7
Section 2 Methodology	8
Section 2.1 Brief Overview	8
Section 2.2 Hospital Frame and Sample Design	8
Section 2.3 NHCS Public Use Data Files Sample Design	8
Section 2.3.1 Sampling Inpatient Discharge Records	9
Section 2.3.2 Sampling ED Visit Records	9
Section 2.4 Data Collection Procedures	10
Section 2.5 Weighting	10
Section 3 Sample Size, Eligibility, and Response Rate	13
Section 4 Data Processing	14
Section 4.1 Diagnosis Data	14
Section 4.2 Procedure Data	15
Section 4.3 Data Quality	16
Section 4.4 Patient Age	16
Section 4.5 Newborn Age	16
Section 4.6 Sex	16
Section 4.7 Discharge Status	16
Section 4.8 Length of Stay	16
Section 4.9 Discharge Month	17
Section 5 Standard Errors and Variance Estimation	18
Section 5.1 Subpopulation Analysis – Subsetting Data	18
Section 6 Presentation Standards	20
Section 7 Data Analysis Guidance	21
Section 7.1 Patient encounter weight	21
Section 7.2 SAS Survey Procedures	22
Section 7.2.1 Specifying the VARMETHOD Option	22
Section 7.2.2 Declaring the REPWEIGHTS Statement	22

Section 7.2.3 Example SAS Code	22
Section 7.3 R Survey Procedures	23
Section 7.4 Stata Survey Procedures	23
Section 8 Survey Content	25
Section 8.1 Variable Missingness Rate by Setting.....	25
Section 8.1.1 Variable Missingness – Among All Inpatient Discharges	25
Section 8.1.2 Variable Missingness – Among All ED Visits.....	27
Section 9 Data Comparison.....	29
Section 9.1 Public Use Data Files and Restricted Use Data File	29
Section 9.2 Public Use Data Files, Restricted Use Data File, and HCUP Nationwide Samples.....	32
Section 10 Preferred Reporting Items for Complex Sample Survey Analysis (PRICSSA) Document for the 2020 National Hospital Care Survey Public Use Data File	34
Section 11 Research Data Center	35
Section 12 References	36
Appendix A Unweighted frequencies for Inpatient discharges and ED visits.....	38

Section 1 About the National Hospital Care Survey

Section 1.1 Background

The National Hospital Care Survey (NHCS) provides data on health care utilization in hospital-based settings, including inpatient departments and emergency departments (EDs). Data collection for NHCS began in 2011, integrating two long-standing National Center for Health Statistics (NCHS) surveys: the National Hospital Discharge Survey, the longest continuously fielded sample of inpatient care from 1965-2010; and the National Hospital Ambulatory Medical Care Survey (1992-2022), which was comprised of three data collection components: hospital EDs (1992-2022), outpatient departments (1992-2017), and ambulatory surgery centers/locations (2009-2017).

NHCS collects data on inpatient discharges and ED visits including information on diagnoses and procedures, length of stay, and patient demographics. NHCS aims to provide health trends and outcomes of the U.S. population's utilization of hospitals in the following ways:

- Provide nationally representative, accurate, and reliable health care data for hospital-based settings.
- Answer key questions of interest to health care professionals, researchers, and policy makers about health care quality, use of resources, and disparities of services to population subgroups.
- Monitor national trends in substance use-related emergency department visits, including opioid visits.
- Provide data that links episodes of care across hospital units, as well as to other data sources, such as the National Death Index.
- Contribute to a stronger public health infrastructure that helps address current and future public health threats.

For 2020, the NHCS sample included 608 non-institutional, non-federal hospitals with six or more staffed inpatient beds in the 50 U.S. states and the District of Columbia. In 2020, there were 205 hospitals that provided data on inpatient discharges and ED visits. Overall, 132,694 inpatient discharges and 388,753 ED visits were selected to create the 2020 NHCS public use data files.

Section 1.2 Data Sources

NHCS receives data from billing and electronic health record sources. This includes Uniform Billing (UB)-04 administrative claims data, the accepted electronic standard for hospital billing as mandated by the

Centers for Medicare & Medicaid Services (<https://www.nubc.org/>). Additionally, hospitals can submit data from Vizient, a member-driven health care services company that collects encounter data from hospitals prior to submitting data to NHCS (<https://www.vizientinc.com/>). Vizient data are similar to UB-04 administrative claims data, but do not contain patient identifiers. Alternatively, hospitals may submit electronic health record (EHR) data, which contain an unlimited number of medical diagnosis and procedure codes, laboratory and medication data, and unstructured clinical notes. NHCS only accepts EHR data in the format of HL7 CDA® R2 Implementation Guide: National Health Care Surveys Release 1, DSTU Release 1.2 – US Realm (http://www.hl7.org/implement/standards/product_brief.cfm?product_id=385).

In 2020, NHCS also received data from the hospital database of the American College of Emergency Physicians (ACEP) for sampled non-participating hospitals (<https://www.acep.org/>). The data from ACEP are only from EDs and do not include any inpatient data, nor does the data include any hospital or patient identifiers.

Section 1.3 Data Confidentiality

NCHS and its agents take the security and confidentiality of NHCS data very seriously. Strict laws have been implemented to establish minimum Federal standards for safeguarding the privacy of individually identifiable health information. Assurance of confidentiality is provided to all hospitals according to Section 308(d) of the Public Health Services Act [42 United States Code 242m (d)]. Strict procedures according to Section 3572 of the Confidential Information Protection and Statistical Efficiency Act (44 U.S.C. 3561-3583) are utilized to prevent disclosure of personal identifiable information in NHCS data. All information which could identify a participating hospital is confidential and seen only by persons associated with NHCS and is not disclosed or released to others for any other purpose. Prior to the release of public use data files, NCHS conducts extensive disclosure risk analysis to minimize the chance of inadvertent disclosure. As a result, selected characteristics and/or data elements may have been masked on the public use data files to minimize the potential risk of disclosure. Masking was performed in such a way to cause minimal impact on the data. See Section 4: Data Processing for more information on which variables in the public use data files were impacted.

The protocol for NHCS has been approved by the NCHS Research Ethics Review Board annually since the survey's establishment (2011).

Section 2 Methodology

Section 2.1 Brief Overview

The 2020 NHCS used a national probability sample of hospitals to collect data on inpatient discharges and ED visits to develop the public use data files. The 2020 NHCS public use data files sample was designed to provide estimates for encounters in both inpatient and ED settings. NHCS data are extracted from hospital billing or EHR systems and transmitted electronically to NCHS for processing. These data are sourced from non-institutional, non-federal hospitals with six or more staffed inpatient beds in the 50 U.S. states and District of Columbia.

Section 2.2 Hospital Frame and Sample Design

The initial sampling frame for NHCS was constructed in 2011 and consisted of 6,622 non-institutional and non-federal hospitals with at least six staffed inpatient beds. From that initial file, a sample of 1,000 eligible hospitals was selected.

The initial sample of 1,000 hospitals was split into two groups of 500 – a base sample and a reserve sample. The base sample hospitals were selected for data collection and the reserve sample of hospitals was held to replenish the base sample if more hospitals of certain hospital types of the fielded sample were deemed ineligible during data collection or to capture more data from a type of hospital. In 2013, 81 general acute hospitals with at least 500 staffed inpatient beds from the reserve sample were added to the base sample for data collection. In 2017, the sample and frame files were updated to include newly constructed hospitals from a new source file. The updating of the NHCS sample and frame occurs every three years. Due to the addition of newly sampled birth hospitals, the 2020 base sample included 608 hospitals and the frame included 6,906 hospitals.

Section 2.3 NHCS Public Use Data Files Sample Design

While the NHCS restricted use data file includes every hospital encounter record submitted to NHCS for the survey year, the 2020 NHCS public use data files consist of a 5% sample of NHCS encounter data from each setting (i.e., both inpatient and ED). In 2020, NHCS collected 2,781,542 inpatient and 7,960,956 ED records. Systematic random sampling was used to select the samples for the public use data files of inpatient and ED records.

This 5% sample of NHCS records was selected for the public use data files instead of the full listing of records for the following reasons: (1) inclusion of all records from inpatient discharges and ED visits from

all sampled hospitals would pose unacceptable disclosure risks for the hospitals; and (2) NHCS collects a lot of records that can be difficult for statistical software to process efficiently, so public use data files were reduced to a manageable size for data users.

Section 2.3.1 Sampling Inpatient Discharge Records

A targeted number of records was determined by taking 5% of the inpatient records (n=132,694). The sampling interval was the total number of inpatient records divided by the targeted sample. The sampling interval used to select the public use data file records in the 2020 NHCS was 20. For each hospital, a random number was selected between 0 and the sampling interval (TE). For each discharge, a random number was selected between 1 and the number of records submitted within each hospital. Inpatient discharges were then sorted by the following variables within a sampled hospital:

Admission Type → Age → Sex → Length of Stay (LOS) → Expected Source of Payer → Discharge Date → Random Discharge Number

Next from the ordered array of records for hospital (j), the records whose assigned “array sequential” numbers are equal to or first after the following numbers were selected for the public use data file sample:

$$R_j + L \times TE$$

Where:

R_j = random number between 0 and TE

L = 0, 1, 2,

TE = sampling interval

The sampled records were then sequentially numbered in the order in which they were selected from the hospital’s submitted records.

Section 2.3.2 Sampling ED Visit Records

For ED visits (n=388,753), similar to inpatient discharges, random numbers were assigned 1 through the number of records submitted by a hospital. Within each hospital, events were sorted as follows:

Hospital ID → Age → Sex → Discharge Status → Expected Source of Payment → Encounter End Date → Randomly assigned number for selection

Then within date of visit, the ED visits were sorted to the numbers randomly assigned at the start of the process. The sorted events were then sequentially numbered from each hospital in the sorted order.

Section 2.4 Data Collection Procedures

In 2020, there were multiple methods for hospitals to submit hospital billing or EHR data. Hospitals could directly submit UB-04 administrative claims or EHR data. Additionally, data could be provided by third-party entities, like Vizient or ACEP. Participating hospitals were required to submit one type of data (e.g., UB-04 administrative claims or EHR, not both). Once data were collected, several steps were required for data processing. Specifications for checking, configuring, and transmitting the data files were developed by NCHS. Once NCHS received the data files, they were processed to harmonize and integrate the data to create a database with similar variables across sources.

Section 2.5 Weighting

Weighting was done to produce hospital level, ED visit and inpatient discharge level estimates and accounts for sampling probabilities and nonresponse. Hospital level data were collected via self-completed forms from in-scope hospitals defined earlier. Inpatient discharges and ED visits were collected (without sampling) from the sampled hospitals via electronic files of their claims or EHR. Participating hospitals submitted all of their data for 2020. While NHCS restricted use data files include all (or 100%) of records sent, the public use data files include a 5% sample of those records.

All inpatient discharge and ED visit data collected for 2020, with the exception of 8 hospitals, were used to produce the inpatient and ED weighted estimates. The 8 hospitals excluded were done so due to quality issues with the submitted data.

The production of weights to produce national estimates involved combining data from the Premier Health Database (PHD) to the 2020 NHCS. The PHD was then removed, and the weights were adjusted for only the 2020 NHCS data. The PHD is a commercially available hospital-based database that collects inpatient service-level data from partner hospitals. It stores information about hospital characteristics such as admitting/attending physician specialties, point of origin, admission type, and discharge status. It also collects encounter-level information on inpatient visits, such as age, sex, race/ethnicity, International Classification of Diseases (ICD) codes, Current Procedure Terminology (CPT) codes, and Healthcare Common Procedure Coding System (HCPCS) codes.

Additionally, the Healthcare Cost and Utilization Project (HCUP) nationwide samples, National Inpatient Sample (NIS) and Nationwide Emergency Department Sample (NEDS) were used as construct controls for calibrating the 2020 encounter-level weights for each setting. NIS and NEDS collect discharge level data from participating state partners. These data sources collect information including patient demographics (e.g., sex, age, race, etc.), encounter characteristics (e.g., diagnoses, procedures, diagnosis-related groups (DRG), admission types, etc.), and hospital-level characteristics (e.g., hospital size and region).

The weighting methodology for point estimation and variance estimation consisted of the following steps:

- Computation of original NHCS weights for NHCS hospitals that reflects the original stratified sampling design and estimated response propensities from a model fitted via a likelihood-based approach.
- Computation of original Premier weights for Premier hospitals that reflects estimated participation propensities from a model fitted via a likelihood-based approach.
- For the combined data set of NHCS and Premier, creation of original combined weights as (original combined) = λ (original NHCS weight) for NHCS hospitals and (original combined) = $(1 - \lambda)$ (original Premier weight) for Premier hospitals.
- Creation of replicate combined weights for variance estimation:
 - Treat the combined NHCS and Premier data as a stratified sample with NHCS strata corresponding to the original design strata and with Premier serving as its own stratum.
 - Use the stratified delete-a-group jackknife (DAGJK), with G_A groups for NHCS and G_B groups for Premier, to construct $G_A + G_B$ sets of replicate combined weights from the original combined weights.
 - Initial assignment of groups will be individual NHCS hospitals (groups of size one hospital, so that $G_A = n_A$) for NHCS and random groups of expected size $n_B/50$ Premier hospitals (so that $G_B = 50$) for Premier.
- Estimates from the combined NHCS and Premier data are then computed using the original combined weights and variance estimates are computed using the replicate combined weights.
- Creation of NHCS calibrated weights:
 - Using the original combined weights and the combined NHCS and Premier data, compute key national estimates to serve as control totals.

- Calibrate the original combined weights for the NHCS-only subset of the combined data to the key national estimates. Unlike the original NHCS weights that are hospital-specific, these calibrated weights are encounter-specific.
- Creation of replicate NHCS calibrated weights for variance estimation:
 - Using the *g*th set of replicate combined weights and the combined NHCS and Premier data, compute the *g*th set of key national estimates to serve as control totals.
 - Calibrate the *g*th set of replicate combined weights for the NHCS-only subset of the combined data to the *g*th set of key national estimates.
- Estimates from the NHCS-only data are then computed using the NHCS calibrated weights and variance estimates are computed using the replicate NHCS calibrated weights.
- Weights and replicate weights for the public use data files were obtained by modifying the NHCS calibrated weights and replicate NHCS calibrated weights. Each replicate weight was adjusted based on the 5% public use data files sampling weight across all encounters within all hospitals.

The 2020 data included a small number of both NHCS specialty hospitals and Premier specialty hospitals. Due to the relatively small number of specialty hospitals in the 2020 NHCS, the inpatient weights for these hospitals were calibrated independently of the acute care hospitals on the weighted survey file. These hospitals were treated as two additional strata and the DAGJK method was used within each stratum.

Section 3 Sample Size, Eligibility, and Response Rate

An eligible responding NHCS hospital must have provided data for at least half the months the hospital was in-scope during the survey year. A hospital is considered a partial respondent if they provided less than half of the months for which the hospital was in scope for the NHCS in the survey year. For example, a hospital that was in-scope for all 12 months during the year must have provided at least 6 months of data.

Of the 205 participating hospitals that were included in the 2020 NHCS database sample, 197 (96.1%) were selected to create the public use data files sample. Eight hospitals that submitted data in 2020 were not included in the public use data files selection due to quality issues identified with the submitted data. From the 197 hospitals, 5% of all records were selected by setting for the public use data files. Overall, 132,694 inpatient discharges and 388,753 ED visits were selected. ED patients transferred to the inpatient department are counted in both the inpatient and ED settings. Table 3.1 presents the number of hospitals, encounters, and response rates of NHCS by setting and type of sample population. See Table 3.1 below for more information.

Table 3.1. Number of hospitals, encounters, and unweighted response rates by setting and file type, National Hospital Care Survey, 2020

	Inpatient		Emergency Department		Total		Unweighted Response Rate ¹
	Hospitals	Encounters	Hospitals	Encounters	Hospitals	Encounters	
Restricted Use Data File	150	2,781,542	200	7,960,956	205	10,010,479	33.7
Public Use Data File	144	132,694	191	388,753	197	521,447	N/A

N/A is not applicable.

¹ Response rate calculation is the number of hospitals that provided data (205) divided by the number of sampled hospitals (608), multiplied by 100.

Section 4 Data Processing

Certain NHCS data elements with missing values were imputed prior to creation of the public use data files. This included imputing values for patients with missing sex, age, and length of stay information. For the 2020 NHCS public use data files, 554 (0.42%) inpatient discharges and 1,334 (0.34%) ED visits contained at least one imputed value for age, sex, and/or length of stay. The data included in the public use data files underwent additional processing to prepare them for release. In addition to imputing missing data, suppression rules such as masking were applied for some records to protect patient confidentiality. Other variables such as age or length of stay were either top-coded or bottom-coded in accordance with NCHS confidentiality requirements; this is noted for specific variables outlined in this section.

Section 4.1 Diagnosis Data

NHCS collects data from participating hospitals from various sources. The diagnosis coding systems varies by data source. Diagnoses from UB-04 administrative claims data are exclusively coded in *International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM)*. EHR submitting hospitals provided diagnoses in SNOMED CT and ICD-10-CM. In the creation of a harmonized and integrated hospital database the SNOMED CT diagnosis codes were translated to ICD-10-CM where applicable. Translation from SNOMED CT to ICD-10-CM was the only modification to the diagnosis codes. On the public use data files, medical diagnosis codes were limited to ICD-10-CM diagnosis codes.

An ICD-10-CM code can have a maximum of 7 characters and is organized by chapters from A to Z. For the 2020 NHCS public use data files, ICD-10-CM codes have been truncated to four characters to minimize disclosure risks. Diagnosis codes are presented as original codes provided by the hospital. While the codes have been truncated, the diagnosis codes are never updated or revised to a different code that would result in a change to the original diagnosis for a visit or hospitalization. To maintain integrity of the data, any codes that appear to be invalid are kept as is.

Duplicate 4-character ICD-10-CM codes were removed for each unique encounter on both the inpatient and ED public use data files. Diagnosis codes were limited to 30 unique records per encounter (variables DX1 through DX30 in the public use data files). A flag was created for encounters with 31 or more diagnosis codes, prior to removal of diagnosis codes above the 30-diagnosis code threshold (variable DX_GT30 in the public use data files).

To account for diagnosis codes that may have been excluded after the removal of diagnosis codes above 30 per encounter, category flags from the Agency for Healthcare Research and Quality (AHRQ) Healthcare Cost and Utilization Project (HCUP) Clinical Classification Software Refined (CCSR) for ICD-10-CM diagnosis codes were created to provide information captured in the removed diagnoses. The AHRQ HCUP CCSR aggregates ICD-10-CM diagnosis codes into clinically meaningful categories. This is intended to be used analytically to examine patterns of health care among several factors.

The AHRQ HCUP CCSR categories are used to provide information from all ICD-10-CM diagnosis codes collected in NHCS, including diagnosis codes that were removed from the public use data files because an encounter had more than 30 diagnoses. All ICD-10-CM diagnosis codes for records in the inpatient and ED public use data files were mapped to a clinical category as defined by the CCSR and a CCSR category flag was created. The 20 CCSR categories with the highest counts were included on the inpatient and ED public use data files, respectively.

CCSR v2023.1 version was used for creating the CCSR categories. More information on the CCSR for ICD-10-CM diagnoses, including documentation, can be found here: https://hcup-us.ahrq.gov/toolssoftware/ccsr/ccs_refined.jsp

Section 4.2 Procedure Data

The coding system used for classifying procedures that occurred during the hospital visit depends on the hospital setting. Inpatient procedures are captured by the *International Classification of Diseases, Tenth Revision, Procedure Coding System (ICD-10-PCS)*. Procedures occurring in the ED are captured using Current Procedural Terminology (CPT®) which is created and maintained by the American Medical Association. CPT codes have been excluded from the ED public use data file at this time due to limitations of approved use in the CPT data user's agreement. Therefore, only ICD-10-PCS codes are available on the inpatient public use data file; procedure codes are unavailable on the ED public use data file.

For the NHCS inpatient public use data file, ICD-10-PCS codes have been truncated to three characters or less to minimize disclosure risks. Duplicate 3-character ICD-10-PCS codes were removed from the inpatient discharges on the public use data file. Procedure codes were limited to a maximum of 10 per encounter (listed in the public use data file as variables PD1 through PD10). A flag was created for inpatient discharges with 11 or more procedure codes, prior to removal of procedure codes above the 10-procedure code threshold (variable PD_GT10 in the public use data file).

Section 4.3 Data Quality

Data users and researchers should be aware of the possibility that the data may contain implausible combinations (e.g., a male patient with a uterine cancer diagnosis) due to coding errors during the collection process. To preserve the data's integrity, errors within the data have been retained without modification or correction. As a result, data users are urged to exercise discretion in determining how they address such records during their analyses.

Section 4.4 Patient Age

Patients with missing age underwent an imputation process that assigned an age based on donor records with a similar first listed diagnosis. Imputed age values comprised 0.26% of inpatient discharges and 0.30% of ED visits. To reduce disclosure risk, encounters of patients ages 86 and older were top coded to 86 years.

Section 4.5 Newborn Age

A newborn flag identifying encounters of infant patients under 1 month was added to both the inpatient and ED public use data files.

Section 4.6 Sex

Patients with missing sex underwent an imputation process that assigned sex based on donor records with similar first listed diagnosis. Imputed sex values comprised 0.29% of inpatient discharges and 0.31% of ED visits.

Section 4.7 Discharge Status

To minimize disclosure risk, four categories for discharge status were collapsed into the "other" category. The "missing" category is composed of three collapsed categories.

Section 4.8 Length of Stay

Length of stay (LOS) is only available for inpatient discharges and is calculated based on the start and end dates of the encounter. If a patient has a missing length of stay due to missing encounter date information, their length of stay value was imputed based on a donor record with a similar procedure as the encounter with a missing length of stay. The percentage of inpatient discharges with an imputed LOS value is 0.11%. To minimize disclosure risk, encounters for patients with inpatient LOS durations over 14 days were top-coded to 15 days. Additionally, encounters for patients with LOS of 30 or more days were flagged (variable LOS_30Days in the public use data file).

Section 4.9 Discharge Month

To minimize risk of disclosure, exact dates are not provided and instead only the month of the inpatient discharge and ED visit are provided on the inpatient and ED public use data files.

Section 5 Standard Errors and Variance Estimation

Standard error is primarily a measure of the sampling variability that occurs by chance because only a sample of hospitals are in NHCS, rather than the entire universe of hospitals. Standard errors and other measures of sampling variability are best determined by using a statistical software package that takes into account the sample designs of surveys to produce such measures.

Propensity modeling was used to develop the base weight through variable selection and harmonization. Replicate calibrated weights were created to produce variance estimates, using the stratified delete-a-group jackknife (DAGJK) method. The DAGJK method resulted in the creation of 100 replicated calibrated weights for the ED public use data file and 122 replicated calibrated weights for the inpatient public use data file to match key national estimates.

The SAS code provided below can be used to generate weighted estimates with standard errors:

```
PROC SURVEYFREQ DATA=FILE VARMETHOD=JACKKNIFE;
TABLE VAR1; * REPLACE "VAR1" WITH THE CATEGORICAL VARIABLE OF INTEREST;
REPWEIGHTS PUF_ENCWTGT_1 - PUF_ENCWTGT_XXX / JKCOEFS=1; * CHANGE "PUF_ENCWTGT_XXX" TO
PUF_ENCWTGT_100 FOR ED OR PUF_ENCWTGT_122 FOR INPATIENT;
WEIGHT PUF_ENCWTGT_BASE;
ODS OUTPUT ONEWAY= ONEWAY;
RUN;
```

See Section 7 for further guidance on how to apply weights and calculate relative standard errors to generate national estimates.

Section 5.1 Subpopulation Analysis – Subsetting Data

For data users who may have a subpopulation of interest, such as a particular age group or sex, a domain analysis, also known as a subgroup or subpopulation analysis must be performed.

For some variance estimation methods, the entire set of data containing the appropriate weights for a particular survey year must be used to obtain the correct variance estimates. Therefore, it is not recommended to drop observations from the dataset when subsetting data, as it may affect variance estimation.

Instead, the estimation procedure must indicate which records are in the subgroup of interest. For example, when examining female patients aged 35 and over, the entire dataset of examined individuals, (both male and female patients of all reported ages) must be read into the statistical software program.

The STAT and DOMAIN statements in the SAS survey procedure, or comparable statements in other programs (SUBSET in R; SUBPOP or over in Stata) must be used to indicate the subgroup of interest (i.e., females aged 35 and over).

Depending on the specifications of a data user's statistical software of choice, an indicator variable created by the data user prior to the procedure may facilitate the identification of the subgroup in the procedure statements.

Section 6 Presentation Standards

Data users should be aware of the reliability of survey estimates, particularly smaller estimates. NCHS has published guidance for the assessment of reliability and presentation of proportions (or percentages) (https://www.cdc.gov/nchs/data/series/sr_02/sr02_175.pdf) and for the presentation of rates and counts (https://www.cdc.gov/nchs/data/series/sr_02/sr02-200.pdf). For presentation or publication of NHCS estimates, we recommend estimates be rounded to the nearest thousand.

These presentation standards apply to products published by NCHS. If, according to the presentation standards, an estimate is not reliable, data users should examine the confidence interval carefully before using the estimate.

Section 7 Data Analysis Guidance

The following section provides an overview on how data users can derive visit or discharge estimates and compute variances to produce standard errors, using statistical software tools such as SAS, R, and Stata. SAS/STAT® software provides a set of procedures whose names begin with SURVEY for survey analysis. R relies on the “survey” package to conduct survey data analysis whereas Stata, uses the “svy” command. SAS, R, and Stata users can use these procedures to conduct statistical analysis on data from the 2020 NHCS public use data files.

Section 7.1 Patient encounter weight

The patient encounter weight is a vital component in the process of producing estimates from sample data and its use should be clearly understood by all data users. The statistics contained on the public use data files reflect only a sample of encounters; a 5% sample of the NHCS data collected from sampled hospitals, not a complete count of all encounters that occurred in the United States. Each inpatient discharge and ED visit record in the public use data files represents one patient encounter in the sample of 132,694 inpatient discharges and 388,753 ED visits. To obtain national estimates from the 5% sample, each record is assigned an inflation factor called the “patient encounter weight” (variable PUF_ENCWGT_BASE in the public use data files).

By aggregating the “patient encounter weights” assigned to the PUF_ENCWGT_BASE variable on the 132,694 inpatient discharges and 388,753 ED visits for 2020, the data user can obtain the estimated total of 36,241,066 inpatient discharges (standard error of 2,184,499 inpatient discharges) and 114,595,499 ED visits (standard error of 5,626,669 ED visits) made in the United States.

Note that estimates of inpatient discharges and ED visits produced from the 2020 NHCS public use data files may differ somewhat from those estimates produced from the NHCS restricted use data files. This is because of adjustments required for the public use data files, as part of the disclosure risk mitigation process. Certain variables were masked on some records for confidentiality purposes. Other variables were top and/or bottom coded in accordance with NCHS confidentiality requirements.

The tables in Section 9 compare aggregate unweighted and weighted data for selected variables between the 2020 NHCS public use data files and restricted use data file by setting.

Section 7.2 SAS Survey Procedures

Section 7.2.1 Specifying the VARMETHOD Option

To correctly use the SURVEYFREQ or SURVEYMEANS procedures, the VARMETHOD= option must be added to the PROC statement to indicate the method to develop the replicate weights. For NHCS, this method is the stratified delete-a-group jackknife (DAGJK) method. Thus, the option is always VARMETHOD=JACKKNIFE.

For example, in PROC SURVEYMEANS:

```
PROC SURVEYMEANS DATA = FILE VARMETHOD=JACKKNIFE;
```

Section 7.2.2 Declaring the REPWEIGHTS Statement

To properly use the SURVEYFREQ or SURVEYMEANS procedures, the REPWEIGHTS options must be specified. The REPWEIGHTS statement identifies the variables containing replicate weights. When using NHCS data, the statement is—

```
REPWEIGHTS PUF_ENCWTG_1 – PUF_ENCWTG_XXX / JKCOEFS=1;
```

There are 122 replicate weights on the inpatient public use data file, so “PUF_ENCWTG_XXX” would be “PUF_ENCWTG_122.” There are 100 replicate weights on the ED public use data file, so “PUF_ENCWTG_XXX” would be “PUF_ENCWTG_100.”

Section 7.2.3 Example SAS Code

The programs below demonstrate how to use replicate weights and calculate variance estimates using SAS SURVEYFREQ and SURVEYMEANS procedures:

For categorical variables:

```
PROC SURVEYFREQ DATA=FILE VARMETHOD=JACKKNIFE;
TABLE VAR1; * REPLACE “VAR1” WITH THE CATEGORICAL VARIABLE OF INTEREST;
REPWEIGHTS PUF_ENCWTG_1 – PUF_ENCWTG_XXX / JKCOEFS=1; * CHANGE “PUF_ENCWTG_XXX” TO
PUF_ENCWTG_122 FOR INPATIENT OR PUF_ENCWTG_100 FOR ED;
WEIGHT PUF_ENCWTG_BASE;
ODS OUTPUT ONEWAY=ONEWAY;
RUN;
```

For continuous variables:

```
PROC SURVEYMEANS DATA=FILE VARMETHOD=JACKKNIFE;
VAR VAR1; * REPLACE “VAR1” WITH THE CONTINUOUS VARIABLE OF INTEREST;
```

```
REPWEIGHTS PUF_ENCWTGT_1 – PUF_ENCWTGT_XXX / JKCOEFS=1; * CHANGE “PUF_ENCWTGT_XXX” TO
PUF_ENCWTGT_122 FOR INPATIENT OR PUF_ENCWTGT_100 FOR ED;
WEIGHT PUF_ENCWTGT_BASE;
ODS OUTPUT STATISTICS= STATISTICS;
RUN;
```

Section 7.3 R Survey Procedures

The R package “survey” can be utilized for complex survey analysis (<https://cran.r-project.org/web/packages/survey/index.html>). The R programs below demonstrate how to use replicate weights and calculate variance estimates in the “survey” package.

```
install.packages("survey")
library(survey)
```

For categorical variables:

```
# Replace “VAR1” with the categorical variable of interest
# Replace “FILE” with filename
# Conduct survey weighted frequency analysis
survey_freq <- svytable(~VAR1, FILE)
```

```
# View the survey frequency results
print(survey_freq)
```

For continuous variables:

```
# Replace “VAR1” with the continuous variable of interest
# Replace “FILE” with filename
# Conduct survey weighted mean analysis
survey_means <- svymean(~VAR1, FILE)
```

```
# View the survey means results
print(survey_means)
```

Section 7.4 Stata Survey Procedures

The Stata programs below demonstrate how to use replicate weights and calculate variance estimates with the svyset command (<https://www.stata.com/manuals/svsvyset.pdf>).

For categorical variables:

```
*Set survey design
svyset PUF_ID, weight(PUF_ENCWTGT_BASE)
```

*Set replicate weights, change "PUF_ENCWGT_XXX" to PUF_ENCWGT_122 for inpatient or PUF_ENCWGT_100 for ED
svyset [pweight=PUF_ENCWGT_BASE], vce(jackknife) jkrweight(PUF_ENCWGT_1-PUF_ENCWGT_XXX)

*Specify one-way tables, change "VAR1" to categorical variable of interest
svy: tab VAR1, count se

*Specify one-way tables, change "VAR1" to categorical variable of interest
svy: tab VAR1, percent

For continuous variables:

*Set survey design
svyset PUF_ID, weight(PUF_ENCWGT_BASE)

*Set replicate weights, change "PUF_ENCWGT_XXX" to PUF_ENCWGT_122 for inpatient or PUF_ENCWGT_100 for ED
svyset [pweight=PUF_ENCWGT_BASE], vce(jackknife) jkrweight(PUF_ENCWGT_1-PUF_ENCWGT_XXX)

*Specify one-way tables, change "VAR1" to continuous variable of interest
svy: mean VAR1

Section 8 Survey Content

For the 2020 NHCS public use data files, 172 variables were included in the inpatient public use data file and 159 variables were included in the ED public use data file. In the inpatient and ED public use data files, 51 (29.7% and 32.1% respectively) variables include medical diagnoses information. In the inpatient public use data file, 11 (6.4%) variables include medical procedure information and two variables measure LOS (1.2%). The ED public use data file does not include information on procedures or LOS.

Please refer to the 2020 NHCS public use data files codebooks for detailed information on the variables including variable names, variable type, variable descriptions, and variable values.

Section 8.1 Variable Missingness Rate by Setting

In the ED public use data file, 52 variables (32.7%) had an unweighted missingness rate that was greater than 5%. In the inpatient public use data files, 39 variables (22.7%) had an unweighted missingness rate that was greater than 5%. See Section 8.1.1 and 8.1.2 for additional information.

However, it is expected that most of the diagnosis variables after the first-listed diagnosis and procedure variables will have a high missingness percentage. This is due to the fact that not all encounters are expected to include more than one or multiple procedures or diagnoses listed. Excluding the non-first-listed diagnosis and procedure variables and the AHRQ CCSR variables, there is only one inpatient variables (variable PD_GT10 in the public use data file) and two ED variables (variables DISCHARGE_STATUS and DX_GT30 in the public use data file) that have a nonresponse percentage above 5%.

Section 8.1.1 Variable Missingness – Among All Inpatient Discharges

The variables in the table below had an unweighted variable missingness percentage greater than 5%. As noted above, it is expected that procedure and diagnoses missingness for some variables would be greater than 5% given that not all encounters are expected to include more than one or multiple procedures or diagnoses listed. Regardless, these are all included below to assist data users in using the NHCS public use data files. The denominator for the percent missing is all visits on the inpatient public use data file.

Table 8.1 Percent missing (unweighted) for variables in the inpatient public use data file with a missingness greater than 5%

Variable Name	Variable Description	Percent Missing
PD1	Procedure #1 (ICD-10-PCS), procedure code	34.3
PD2	Procedure #2 (ICD-10-PCS), procedure code	58.6
PD3	Procedure #3 (ICD-10-PCS), procedure code	74.8
PD4	Procedure #4 (ICD-10-PCS), procedure code	84.9
PD5	Procedure #5 (ICD-10-PCS), procedure code	91.1
PD6	Procedure #6 (ICD-10-PCS), procedure code	94.5
PD7	Procedure #7 (ICD-10-PCS), procedure code	96.6
PD8	Procedure #8 (ICD-10-PCS), procedure code	97.9
PD9	Procedure #9 (ICD-10-PCS), procedure code	98.7
PD10	Procedure # 10 (ICD-10-PCS), procedure code	99.2
PD_GT10	Procedure Greater than 10 flag; associated with encounters that have 11 or more procedures	34.3
DX3	Diagnosis #3 (ICD-10-CM), diagnosis code	6.9
DX4	Diagnosis #4 (ICD-10-CM), diagnosis code	11.2
DX5	Diagnosis #5 (ICD-10-CM), diagnosis code	15.8
DX6	Diagnosis #6 (ICD-10-CM), diagnosis code	20.4
DX7	Diagnosis #7 (ICD-10-CM), diagnosis code	25.1
DX8	Diagnosis #8 (ICD-10-CM), diagnosis code	29.9
DX9	Diagnosis #9 (ICD-10-CM), diagnosis code	34.7
DX10	Diagnosis #10 (ICD-10-CM), diagnosis code	40.6
DX11	Diagnosis #11 (ICD-10-CM), diagnosis code	54.1
DX12	Diagnosis #12 (ICD-10-CM), diagnosis code	66.5
DX13	Diagnosis #13 (ICD-10-CM), diagnosis code	71.5
DX14	Diagnosis #14 (ICD-10-CM), diagnosis code	74.6
DX15	Diagnosis #15 (ICD-10-CM), diagnosis code	76.9
DX16	Diagnosis #16 (ICD-10-CM), diagnosis code	79.0
DX17	Diagnosis #17 (ICD-10-CM), diagnosis code	80.9
DX18	Diagnosis #18 (ICD-10-CM), diagnosis code	82.8
DX19	Diagnosis #19 (ICD-10-CM), diagnosis code	84.6
DX20	Diagnosis #20 (ICD-10-CM), diagnosis code	86.3
DX21	Diagnosis #21 (ICD-10-CM), diagnosis code	87.7
DX22	Diagnosis #22 (ICD-10-CM), diagnosis code	89.0
DX23	Diagnosis #23 (ICD-10-CM), diagnosis code	90.2
DX24	Diagnosis #24 (ICD-10-CM), diagnosis code	91.4
DX25	Diagnosis #25 (ICD-10-CM), diagnosis code	92.5
DX26	Diagnosis #26 (ICD-10-CM), diagnosis code	93.6
DX27	Diagnosis #27 (ICD-10-CM), diagnosis code	94.4
DX28	Diagnosis #28 (ICD-10-CM), diagnosis code	95.1
DX29	Diagnosis #29 (ICD-10-CM), diagnosis code	95.7

Variable Name	Variable Description	Percent Missing
DX30	Diagnosis #30 (ICD-10-CM), diagnosis code	96.2

Section 8.1.2 Variable Missingness – Among All ED Visits

The variables in the table below had an unweighted variable missingness percentage greater than 5%.

The denominator for the percent missing is all visits on the ED public use data file.

Table 8.2 Percent missing (unweighted) for variables in the emergency department public use data file with a missingness greater than 5%

Variable Name	Variable Description	Percent Missing
DISCHARGE_STATUS	Discharge Status	7.2
DX1	Diagnosis #1 (ICD-10-CM), primary or first-listed diagnosis code	8.4
DX2	Diagnosis #2 (ICD-10-CM), diagnosis code	25.7
DX3	Diagnosis #3 (ICD-10-CM), diagnosis code	41.5
DX4	Diagnosis #4 (ICD-10-CM), diagnosis code	53.9
DX5	Diagnosis #5 (ICD-10-CM), diagnosis code	63.8
DX6	Diagnosis #6 (ICD-10-CM), diagnosis code	71.0
DX7	Diagnosis #7 (ICD-10-CM), diagnosis code	76.4
DX8	Diagnosis #8 (ICD-10-CM), diagnosis code	80.4
DX9	Diagnosis #9 (ICD-10-CM), diagnosis code	83.6
DX10	Diagnosis #10 (ICD-10-CM), diagnosis code	86.6
DX11	Diagnosis #11 (ICD-10-CM), diagnosis code	92.4
DX12	Diagnosis #12 (ICD-10-CM), diagnosis code	96.1
DX13	Diagnosis #13 (ICD-10-CM), diagnosis code	97.3
DX14	Diagnosis #14 (ICD-10-CM), diagnosis code	97.9
DX15	Diagnosis #15 (ICD-10-CM), diagnosis code	98.3
DX16	Diagnosis #16 (ICD-10-CM), diagnosis code	98.6
DX17	Diagnosis #17 (ICD-10-CM), diagnosis code	98.8
DX18	Diagnosis #18 (ICD-10-CM), diagnosis code	99.0
DX19	Diagnosis #19 (ICD-10-CM), diagnosis code	99.2
DX20	Diagnosis #20 (ICD-10-CM), diagnosis code	99.4
DX21	Diagnosis #21 (ICD-10-CM), diagnosis code	99.4
DX22	Diagnosis #22 (ICD-10-CM), diagnosis code	99.5
DX23	Diagnosis #23 (ICD-10-CM), diagnosis code	99.6
DX24	Diagnosis #24 (ICD-10-CM), diagnosis code	99.7
DX25	Diagnosis #25 (ICD-10-CM), diagnosis code	99.8
DX26	Diagnosis #26 (ICD-10-CM), diagnosis code	99.9
DX27	Diagnosis #27 (ICD-10-CM), diagnosis code	99.9
DX28	Diagnosis #28 (ICD-10-CM), diagnosis code	99.9
DX29	Diagnosis #29 (ICD-10-CM), diagnosis code	99.9

Variable Name	Variable Description	Percent Missing
DX30	Diagnosis #30 (ICD-10-CM), diagnosis code	99.9
DX_GT30	Diagnosis Greater than 30 flag; associated with encounters that have 31 or more diagnosis codes	8.4
CCSR_CIR007	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR007 [Essential hypertension]	8.4
CCSR_CIR008	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR008 [Hypertension with complications and secondary hypertension]	8.4
CCSR_CIR011	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR011 [Coronary atherosclerosis and other heart disease]	8.4
CCSR_CIR012	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR012 [Nonspecific chest pain]	8.4
CCSR_CIR019	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR019 [Heart failure]	8.4
CCSR_END005	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END005 [Diabetes mellitus, Type 2]	8.4
CCSR_END010	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END010 [Disorders of lipid metabolism]	8.4
CCSR_END011	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END011 [Fluid and electrolyte disorders]	8.4
CCSR_EXT020	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR_EXT020 [External cause codes: intent of injury, accidental/unintentional]	8.4
CCSR_FAC016	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR FAC016 [Exposure, encounters, screening or contact with infectious disease]	8.4
CCSR_FAC021	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR FAC021 [Personal/family history of disease]	8.4
CCSR_FAC025	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR FAC025 [Other specified status]	8.4
CCSR_GEN003	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR GEN003 [Chronic kidney disease]	8.4
CCSR_INJ017	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR INJ017 [Superficial injury; contusion, initial encounter]	8.4
CCSR_MBD024	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR MBD024 [Tobacco-related disorders]	8.4
CCSR_MUS010	Indication that at least one ICD-10-CM diagnosis on the record is -- included in CCSR MUS010 [Musculoskeletal pain, not low back pain]	8.4
CCSR_SYM004	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR SYM004 [Nausea and vomiting]	8.4
CCSR_SYM006	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR SYM006 [Abdominal pain and other digestive/abdomen signs and symptoms]	8.4
CCSR_SYM010	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR SYM010 [Nervous system signs and symptoms]	8.4
CCSR_SYM013	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR SYM013 [Respiratory signs and symptoms]	8.4

Section 9 Data Comparison

Section 9.1 Public Use Data Files and Restricted Use Data File

Of the 205 participating hospitals that were included in the 2020 NHCS restricted use data file, 197 (96.1%) were selected to create the public use data files sample. The 2020 inpatient public use data file contains 132,694 inpatient discharges, for a weighted total of 36,241,066 inpatient discharges (standard error of 2,184,449 inpatient discharges). The 2020 ED public use data file contains 388,753 ED visits, for a weighted total of 114,595,499 ED visits (standard error of 5,626,669 ED visits). The 2020 NHCS restricted use data file contains unweighted data from 150 hospitals that submitted 2,832,062 inpatient discharges, for a weighted total of 36,342,461 inpatient discharges (standard error of 2,173,127 inpatient discharges). The 2020 NHCS restricted use data file contains data from 199 hospitals that submitted 7,978,530 ED visits, for a weighted total of 114,561,914 ED visits (standard error of 5,631,753 ED visits). A comparison of weighted frequencies for inpatient discharges and ED visits in the public use data files and restricted use data file is presented in Table 9.1 and Table 9.2.

Table 9.1. Comparison of frequencies for inpatient discharges on the public use data file (weighted n=36,241,066) and restricted use data file (weighted n=36,342,461), National Hospital Care Survey, 2020

Variable	Public Use Data File				Restricted Use Data File			
	Unweighted Count	Weighted			Unweighted Count	Weighted		
		Count	Std. Error	%		Count	Std. Error	%
Age (in years)								
Under 1	14,953	3,995,014	358,556	11.0	299,588	4,019,203	357,865	11.1
1-17	5,587	1,772,950	362,002	4.9	112,332	1,822,994	401,161	5.0
18-44	33,474	9,268,002	643,320	25.6	668,435	9,266,652	618,432	25.5
45-64	32,139	8,296,364	475,751	22.9	642,647	8,313,822	480,417	22.9
65-74	21,626	5,619,587	390,756	15.5	432,504	5,641,533	380,754	15.5
75 and over	24,835	7,255,820	554,595	20.0	496,790	7,245,679	536,847	19.9
Missing	80	33,329	31,421	0.1	1,584	32,578	30,783	0.1
Sex								
Male	60,701	16,225,708	977,850	44.8	1,214,397	16,301,395	947,177	44.9
Female	71,888	19,957,691	1,242,206	55.1	1,437,897	20,000,798	1,249,506	55.0
Missing	105	57,666	40,948	0.2	1,586	40,268	34,166	0.1
Discharge status								
Routine to Home	85,777	23,890,910	1,514,092	65.9	1,717,716	23,995,053	1,487,301	66.0
Left against medical advice	1,877	554,005	111,639	1.5	36,993	535,027	63,037	1.5
Transfer to short term facility	1,269	665,653	54,815	1.8	25,296	677,526	65,479	1.9
Transfer to long term facility	4,379	1,014,693	109,292	2.8	88,405	1,088,492	125,091	3.0
Home health care	18,621	4,582,308	452,497	12.6	368,387	4,525,721	465,387	12.5
Hospice care - home or medical facility	2,298	746,310	118,157	2.1	47,334	710,366	86,710	2.0
Other	12,318	3,456,947	266,716	9.5	246,651	3,482,340	268,650	9.6
Dead	3,654	925,297	80,248	2.6	72,707	922,816	86,101	2.5
Missing	2,501	404,942	220,876	1.1	50,391	405,121	217,399	1.1
Discharge month								
January	12,532	3,380,965	220,293	9.3	250,989	3,567,564	212,643	9.8
February	11,683	3,246,016	222,134	9.0	233,131	3,303,877	201,237	9.1
March	11,116	3,233,317	286,645	8.9	222,254	3,120,442	195,190	8.6
April	8,790	2,416,304	168,918	6.7	174,397	2,446,680	145,625	6.7
May	10,066	2,777,587	196,554	7.7	200,654	2,741,202	173,324	7.5
June	10,593	2,829,008	196,434	7.8	214,088	2,929,595	185,720	8.1
July	11,531	3,211,502	238,502	8.9	229,713	3,147,032	194,042	8.7
August	11,329	3,072,858	273,846	8.5	225,440	3,022,906	195,075	8.3
September	11,150	2,969,109	265,546	8.2	224,598	3,003,929	177,470	8.3
October	11,916	3,136,200	234,282	8.7	237,831	3,186,507	203,642	8.8
November	10,817	2,876,424	222,923	7.9	215,403	2,859,902	191,226	7.9
December	11,171	3,091,776	220,937	8.5	225,382	3,012,827	210,362	8.3

Table 9.2. Comparison of frequencies for ED visits on the public use data file (weighted n=114,595,499) and restricted use data file (weighted n=114,561,914), National Hospital Care Survey, 2020

Variable	Public Use Data File				Restricted Use Data File			
	Unweighted Count	Weighted			Unweighted Count	Weighted		
		Count	Std. Error	%		Count	Std. Error	%
Age (in years)								
Under 1	6,708	1,630,620	181,009	1.4	135,018	1,625,536	182,786	1.4
1-17	48,685	14,133,681	979,858	12.3	972,895	14,136,097	978,521	12.3
18-44	154,154	47,890,772	2,191,911	41.8	3,083,235	47,881,234	2,199,155	41.8
45-64	98,212	28,608,338	1,499,687	25.0	1,963,972	28,594,994	1,499,431	25.0
65-74	38,564	10,790,224	706,467	9.4	771,265	10,789,722	710,914	9.4
75 and over	41,869	11,461,509	959,498	10.0	837,490	11,455,083	961,783	10.0
Missing	561	80,354	81,838	0.1	11,069	79,247	82,724	0.1
Sex								
Male	180,775	52,091,506	2,654,647	45.5	3,616,311	52,083,659	2,656,643	45.5
Female	207,345	62,371,631	3,014,106	54.4	4,146,674	62,351,492	3,016,048	54.4
Missing	633	132,362	86,907	0.1	11,959	126,763	89,996	0.1
Discharge status								
Routine to Home	314,022	99,216,245	4,445,255	86.6	6,280,325	99,150,961	4,445,134	86.5
Left against medical advice	7,554	2,188,161	157,802	1.9	150,898	2,189,987	158,159	1.9
Transfer to short term facility	5,279	2,181,121	200,617	1.9	105,930	2,204,930	195,949	1.9
Transfer to long term facility	2,115	407,931	71,083	0.4	42,575	409,031	72,116	0.4
Home health care	6,093	1,469,910	334,619	1.3	121,534	1,477,416	338,695	1.3
Hospice care - home or medical facility	1,185	231,975	54,247	0.2	23,621	236,296	56,557	0.2
Other	22,527	4,225,748	514,271	3.7	450,382	4,205,577	517,510	3.7
Dead	2,063	374,198	29,382	0.3	41,273	380,193	31,875	0.3
Missing	27,915	4,300,210	1,643,017	3.8	558,406	4,307,523	1,642,748	3.8
Discharge month								
January	44,757	13,139,793	767,095	11.5	895,631	13,120,600	760,780	11.5
February	40,784	12,004,237	674,673	10.5	815,526	11,994,536	682,015	10.5
March	35,874	10,470,739	573,567	9.1	716,063	10,454,374	575,262	9.1
April	20,942	6,302,189	355,409	5.5	419,492	6,243,975	322,949	5.5
May	25,203	7,562,675	443,290	6.6	503,667	7,615,398	437,545	6.6
June	27,892	8,425,972	426,967	7.4	558,200	8,421,553	416,572	7.4
July	33,246	9,749,850	494,295	8.5	664,429	9,788,941	496,944	8.5
August	34,235	10,107,168	617,427	8.8	685,486	10,062,811	620,626	8.8
September	31,859	9,423,894	573,906	8.2	636,584	9,412,833	574,582	8.2
October	33,697	10,091,948	607,176	8.8	674,036	10,098,244	605,686	8.8
November	30,689	8,828,415	503,990	7.7	614,721	8,831,836	490,408	7.7
December	29,575	8,488,620	479,126	7.4	591,109	8,516,813	487,726	7.4

Section 9.2 Public Use Data Files, Restricted Use Data File, and HCUP Nationwide

Samples

As stated in Section 2.5, the HCUP nationwide samples, the NIS and the NEDS were used as construct controls for calibrating the 2020 encounter-level weights for each setting. NIS and NEDS collect hospital discharge data from participating state partners. These data sources collect patient demographics (e.g., sex, age, race, etc.), encounter characteristics (e.g., diagnoses, procedures, DRGs, admission types, etc.), and hospital-level characteristics (e.g., hospital size and region).

Statewide data organizations voluntarily participate in NEDS by providing HCUP with data from ED visits that may or may not have resulted in a hospital admission. It is estimated that the 2020 NEDS accounts for 85% of the U.S. population and 84% of ED visits reported in the American Hospital Association Annual Survey Database. Records are selected for NEDS from two existing HCUP databases: the State Emergency Department Databases (SEDD) and the State Inpatient Databases (SID).

NIS is a sample of all-payer discharges from U.S. community hospitals, defined as “all non-Federal, short-term, general, and other specialty hospitals, excluding hospital units of institutions.” It’s estimated that the 2020 NIS covers 98% of the U.S. population and 97% of discharges from U.S. community hospitals. NIS is sampled from the SID.

A comparison of weighted total inpatient discharges on the 2020 NHCS public use data file, NHCS restricted use data file, and HCUP NIS is in Table 9.3. A comparison of weighted totals for ED visits in the public use data file, restricted use data file, and NEDS is presented in Table 9.4.

Table 9.3 Comparison of weighted total inpatient discharges on the 2020 NHCS public use data file, NHCS restricted use data file, and the HCUP NIS

Data File	Inpatient Discharges	Standard Error
National Hospital Care Survey (NHCS) public use data file	36,241,066	2,184,449
National Hospital Care Survey (NHCS) restricted use data file	36,342,461	2,173,127
Healthcare Cost and Utilization Project National Inpatient Sample (HCUP NIS)	32,355,827	284,645

Table 9.4. Comparison of weighted total emergency department visits on the 2020 NHCS public use data file, NHCS restricted use data file, and HCUP NEDS

Data File	Emergency Department Visits	Standard Error
National Hospital Care Survey (NHCS) public use data file	114,595,499	5,626,669
National Hospital Care Survey (NHCS) restricted use data file	114,561,914	5,631,753
Healthcare Cost and Utilization Project Nationwide Emergency Department Sample (HCUP NEDS)	123,278,165	2,696,916

Section 10 Preferred Reporting Items for Complex Sample Survey Analysis (PRICSSA) Document for the 2020 National Hospital Care Survey Public Use Data File

Table 10.1 below provides a Preferred Reporting Items for Complex Survey Analysis (PRICSSA) document (Seidenberg, Moser, & West 2023) for users of the 2020 NHCS public use data. This information may be helpful to users when analyzing the 2020 NHCS.

Table 10.1 Preferred Reporting Items for Complex Sample Survey Analysis

Preferred Reporting Items for Complex Sample Survey Analysis (PRICSSA)	Description
Name of survey	National Hospital Care Survey
Data collection mode	Administrative data collection of hospital UB-04 administrative claims or EHR data
Target population	Hospital inpatient discharges and ED visits to nonfederal noninstitutional hospitals with six or more staffed inpatient beds in the United States
Populations excluded	Hospital inpatient discharges and ED visits to federal hospitals, institutional hospitals, or hospitals with less than six staffed inpatient beds
Sample design	Stratified random sample
Variance and standard error estimation	Jackknife variance estimation
Base Weight	PUF_ENCWGT_BASE
Replicate Weights	Inpatient: PUF_ENCWGT_1 – PUF_ENCWGT122 ED: PUF_ENCWGT_1 – PUF_ENCWGT100
Presentation standards	Proportions or percentages: https://www.cdc.gov/nchs/data/series/sr_02/sr02_175.pdf Rates and counts: https://www.cdc.gov/nchs/data/series/sr_02/sr02-200.pdf
Unweighted total sample size	132,694 inpatient discharges 388,753 ED visits
Weighted total sample size	36,241,066 inpatient discharges 114,595,499 ED visits
Response rate (unweighted)	33.7%
Location of example code	See Section 7

Section 11 Research Data Center

NCHS operates the Research Data Center (RDC) to allow researchers access to restricted-use data. The RDC is responsible for protecting the confidentiality of survey respondents, study subjects, and institutions while providing access to restricted-use data for statistical purposes. The 2020 NHCS restricted use data file, which contains inpatient and ED data excluded from the public use data file due to disclosure risks and NCHS confidentiality requirements, can be accessed through the Federal and NCHS RDC. In addition, the 2020 NHCS restricted use data file will be linked to other vital and administrative records such as the National Death Index (NDI). This allows the linked data to expand the analytic utility of NHCS data and provides the opportunity to conduct a vast array of studies focused on the associations between a variety of health factors, health care utilization and mortality.

For information on how to access the 2020 NHCS restricted use data file through the RDC, please see: <https://www.cdc.gov/rdc/data/b1/2019-2020-NHCS-RDC-Data-Dictionary-508.pdf>.

Section 12 References

Agency for Healthcare Research and Quality (AHRQ). Clinical Classifications Software (CCS) for ICD-10-CM/PCS. Healthcare Cost and Utilization Project (HCUP). Available at: https://hcup-us.ahrq.gov/toolssoftware/ccsr/ccs_refined.jsp. Accessed November 29, 2023.

American College of Emergency Physicians (ACEP). Available at: <https://www.acep.org>. Accessed November 29, 2023.

Health Level Seven International. HL7 CDA® R2 Implementation Guide: National Health Care Surveys (NHCS), R1 STU Release 3.1 - US Realm. Available at: http://www.hl7.org/implement/standards/product_brief.cfm?product_id=385. Accessed November 29, 2023.

Lumley T (2023). "survey: analysis of complex survey samples." R package version 4.2. Available at: <https://cran.r-project.org/web/packages/survey/index.html>.

National Center for Health Statistics. Division of Health Care Statistics. National Center for Health Statistics Research Data Center National Hospital Care Survey 2019 and 2020 Restricted Use File Data Dictionary. Hyattsville, Maryland. Available at: <https://www.cdc.gov/rdc/data/b1/2019-2020-NHCS-RDC-Data-Dictionary-508.pdf>.

National Uniform Billing Committee (NUBC). Available at: <https://www.nubc.org>. Accessed November 29, 2023.

Parker JD, Talih M, Irinata KE, Zhang G, Branum AM, Davis D, et al. National Center for Health Statistics data presentation standards for rates and counts. National Center for Health Statistics. Vital Health Stat 2(200). 2023. DOI: <https://dx.doi.org/10.15620/cdc:124368>.

Parker JD, Talih M, Malec DJ, et al. National Center for Health Statistics data presentation standards for proportions. National Center for Health Statistics. Vital Health Stat 2(175). 2017. Available at: https://www.cdc.gov/nchs/data/series/sr_02/sr02_175.pdf.

Seidenberg AB, Moser RP, West BT. Preferred Reporting Items for Complex Sample Survey Analysis (PRICSSA). *Journal of Survey Statistics and Methodology* 2023;11(4):743-757.

StataCorp. 2023. Stata 18 Survey Data Reference Manual. College Station, TX: Stata Press. Available at: <https://www.stata.com/manuals/svsvyset.pdf>.

Vizient Inc. Available at: <https://www.vizientinc.com>. Accessed November 29, 2023.

Appendix A Unweighted frequencies for Inpatient discharges and ED visits

Appendix Table A.1. Unweighted frequencies for inpatient discharges on the public use data file, National Hospital Care Survey, 2020 (n=132,694)

Variable	Description	Count	%
Year	Survey year		
2020		132,694	100.0
Sex	Patient sex		
1	Male	60,701	44.8
2	Female	71,888	55.1
-9	Missing	105	0.2
Newborn	Newborn flag		
1	Yes	13,963	10.5
2	No	118,731	89.5
LOS_30DAYS	Length of stay in hospital (days) greater than 30 days		
1	Yes	2,342	1.8
2	No	130,337	98.2
-9	Missing	15	0.0
Discharge_month	Discharge month		
1	January	12,532	9.4
2	February	11,683	8.8
3	March	11,116	8.4
4	April	8,790	6.6
5	May	10,066	7.6
6	June	10,593	8.0
7	July	11,531	8.7
8	August	11,329	8.5
9	September	11,150	8.4
10	October	11,916	9.0
11	November	10,817	8.2
12	December	11,171	8.4
Discharge_status	Discharge status		
1	Routine to Home	85,777	64.6
2	Left against medical advice	1,877	1.4
3	Transfer to short term facility	1,269	1.0
4	Transfer to long term facility	4,379	3.3
5	Home health care	18,621	14.0
6	Hospice care - home or medical facility	2,298	1.7
7	Other	12,318	9.3
8	Dead	3,654	2.8
-9	Missing	2,501	1.9
PD_GT10	Greater than 10 procedures associated with the encounters		
1	Yes	779	0.6
2	No	86,422	65.1
-9	Missing	45,493	34.3
DX_GT30	Encounters that had DX more than 30 diagnoses		
1	Yes	4,483	3.4
2	No	126,870	95.6
-9	Missing	1,341	1.0

Variable	Description	Count	%
CCSR_CIR007	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR007 [Essential hypertension]		
1	Yes	31,698	23.9
2	No	99,656	75.1
-9	Missing	1,340	1.0
CCSR_CIR008	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR008		
1	Yes	28,695	21.6
2	No	102,659	77.4
-9	Missing	1,340	1.0
CCSR_CIR011	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR011		
1	Yes	20,589	15.5
2	No	110,765	83.5
-9	Missing	1,340	1.0
CCSR_CIR017	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR017		
1	Yes	19,211	14.5
2	No	112,143	84.5
-9	Missing	1,340	1.0
CCSR_CIR019	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR019		
1	Yes	20,556	15.5
2	No	110,798	83.5
-9	Missing	1,340	1.0
CCSR_DIG004	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR DIG004		
1	Yes	19,513	14.7
2	No	111,841	84.3
-9	Missing	1,340	1.0
CCSR_END003	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END003		
1	Yes	18,383	13.9
2	No	112,971	85.1
-9	Missing	1,340	1.0
CCSR_END005	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END005		
1	Yes	25,299	19.1
2	No	106,055	79.9
-9	Missing	1,340	1.0
CCSR_END009	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END009		
1	Yes	19,050	14.4
2	No	112,304	84.6
-9	Missing	1,340	1.0
CCSR_END010	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END010		
1	Yes	33,757	25.4
2	No	97,597	73.6
-9	Missing	1,340	1.0
CCSR_END011	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END011		
1	Yes	33,730	25.4

Variable	Description	Count	%
2	No	97,624	73.6
-9	Missing	1,340	1.0
CCSR_FAC009	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR FAC009		
1	Yes	17,976	13.5
2	No	113,378	85.4
-9	Missing	1,340	1.0
CCSR_FAC016	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR FAC016		
1	Yes	41,802	31.5
2	No	89,552	67.5
-9	Missing	1,340	1.0
CCSR_FAC021	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR FAC021		
1	Yes	43,519	32.8
2	No	87,835	66.2
-9	Missing	1,340	1.0
CCSR_FAC025	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR FAC025		
1	Yes	55,186	41.6
2	No	76,168	57.4
-9	Missing	1,340	1.0
CCSR_GEN002	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR GEN002		
1	Yes	20,750	15.6
2	No	110,604	83.4
-9	Missing	1,340	1.0
CCSR_GEN003	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR GEN003		
1	Yes	20,735	15.6
2	No	110,619	83.4
-9	Missing	1,340	1.0
CCSR_INF003	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR INF003		
1	Yes	17,630	13.3
2	No	113,724	85.7
-9	Missing	1,340	1.0
CCSR_PRG023	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR PRG023		
1	Yes	10,589	8.0
2	No	120,765	91.0
-9	Missing	1,340	1.0

Appendix Table A.2. Unweighted frequencies for ED visits on the public use data file, National Hospital Care Survey, 2020 (n=388,753)

Variable	Description	Count	%
Year	Survey year		
2020		388,753	100.0
Sex	Patient sex		
1	Male	180,776	46.5
2	Female	207,345	53.3
-9	Missing	632	0.2
Newborn	Newborn flag		
1	Yes	1,099	0.3
2	No	387,654	99.7
Discharge_month	Discharge month		
1	January	44,757	11.5
2	February	40,784	10.5
3	March	35,874	9.2
4	April	20,942	5.4
5	May	25,203	6.5
6	June	27,892	7.2
7	July	33,246	8.6
8	August	34,235	8.8
9	September	31,859	8.2
10	October	33,697	8.7
11	November	30,689	7.9
12	December	29,575	7.6
Discharge_status	Discharge status		
1	Routine to Home	314,022	80.8
2	Left against medical advice	7,554	1.9
3	Transfer to short term facility	5,279	1.4
4	Transfer to long term facility	2,115	0.5
5	Home health care	6,093	1.6
6	Hospice care - home or medical facility	1,185	0.3
7	Other	22,527	5.8
8	Dead	2,063	0.5
-9	Missing	27,915	7.2
DX_GT30	Encounters that had more than 30 diagnoses		
1	Yes	249	0.1
2	No	355,727	91.5
-9	Missing	32,777	8.4
CCSR_CIR007	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR007		
1	Yes	56,280	14.5
2	No	299,728	77.1
-9	Missing	32,745	8.4
CCSR_CIR008	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR008		
1	Yes	21,077	5.4
2	No	334,931	86.2
-9	Missing	32,745	8.4
CCSR_CIR011	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR011		
1	Yes	16,111	4.1
2	No	339,897	87.4
-9	Missing	32,745	8.4

Variable	Description	Count	%
CCSR_CIR012	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR012		
1	Yes	26,811	6.9
2	No	329,197	84.7
-9	Missing	32,745	8.4
CCSR_CIR019	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR CIR019		
1	Yes	15,479	4.0
2	No	340,529	87.6
-9	Missing	32,745	8.4
CCSR_END005	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END005		
1	Yes	32,214	8.3
2	No	323,794	83.3
-9	Missing	32,745	8.4
CCSR_END010	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END010		
1	Yes	30,135	7.8
2	No	325,873	83.8
-9	Missing	32,745	8.4
CCSR_END011	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR END011		
1	Yes	25,008	6.4
2	No	331,000	85.1
-9	Missing	32,745	8.4
CCSR_EXT020	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR_EXT020		
1	Yes	47,643	12.3
2	No	308,365	79.3
-9	Missing	32,745	8.4
CCSR_FAC016	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR FAC016		
1	Yes	43,414	11.2
2	No	312,594	80.4
-9	Missing	32,745	8.4
CCSR_FAC021	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR FAC021		
1	Yes	51,828	13.3
2	No	304,180	78.2
-9	Missing	32,745	8.4
CCSR_FAC025	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR FAC025		
1	Yes	73,047	18.8
2	No	282,961	72.8
-9	Missing	32,745	8.4
CCSR_GEN003	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR GEN003		
1	Yes	15,824	4.1
2	No	340,184	87.5
-9	Missing	32,745	8.4
CCSR_INJ017	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR INJ017		
1	Yes	16,771	4.3

Variable	Description	Count	%
2	No	339,237	87.3
-9	Missing	32,745	8.4
CCSR_MBD024	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR MBD024		
1	Yes	39,412	10.1
2	No	316,596	81.4
-9	Missing	32,745	8.4
CCSR_MUS010	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR MUS010		
1	Yes	28,392	7.3
2	No	327,616	84.3
-9	Missing	32,745	8.4
CCSR_SYM004	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR SYM004		
1	Yes	22,848	5.9
2	No	333,160	85.7
-9	Missing	32,745	8.4
CCSR_SYM006	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR SYM006		
1	Yes	40,246	10.4
2	No	315,762	81.2
-9	Missing	32,745	8.4
CCSR_SYM010	Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR SYM010		
1	Yes	23,815	6.1
2	No	332,193	85.5
-9	Missing	32,745	8.4