

Deaths: Preliminary Data for 2008

by Arialdi M. Miniño, M.P.H.; Jiaquan Xu, M.D.; and Kenneth D. Kochanek, M.A., Division of Vital Statistics

Abstract

Objectives—This report presents preliminary U.S. data on deaths, death rates, life expectancy, leading causes of death, and infant mortality for 2008 by selected characteristics such as age, sex, race, and Hispanic origin.

Methods—Data in this report are based on death records comprising more than 99 percent of the demographic and medical files for all deaths in the United States in 2008. The records are weighted to independent control counts for 2008. For certain causes of death such as unintentional injuries, homicides, suicides, drug-induced deaths, and sudden infant death syndrome, preliminary and final data may differ because of the truncated nature of the preliminary file. Comparisons are made with 2007 final data.

Results—The age-adjusted death rate decreased from 760.2 deaths per 100,000 population in 2007 to 758.7 deaths per 100,000 population in 2008. From 2007 to 2008, age-adjusted death rates decreased significantly for 6 of the 15 leading causes of death: Diseases of heart, Malignant neoplasms, Cerebrovascular diseases, Accidents (unintentional injuries), Diabetes mellitus, and Assault (homicide). From 2007 to 2008, age-adjusted death rates increased significantly for 6 of the 15 leading causes of death: Chronic lower respiratory diseases; Alzheimer's disease; Influenza and pneumonia; Nephritis, nephrotic syndrome and nephrosis; Intentional self-harm (suicide); and Essential hypertension and hypertensive renal disease. Life expectancy decreased by 0.1 year from 77.9 years in 2007 to 77.8 in 2008.

Keywords: death rates • life expectancy • vital statistics • preliminary

Introduction

This report presents preliminary mortality data for the United States based on vital records for a substantial proportion of deaths occurring in 2008. Statistics in preliminary reports are generally considered reliable; past analyses reveal that most statistics shown in preliminary reports for 1995–2007 were confirmed by the final statistics for each of those years (1–13).

Data Sources and Methods

Preliminary data in this report are based on records of deaths that occurred in calendar year 2008, which were received from state vital statistics offices and processed by the Centers for Disease Control and Prevention's National Center for Health Statistics (NCHS) as of September 22, 2010. Estimates of the level of completeness of preliminary data for each state are shown in [Table I](#). Detailed information on the nature, sources, and qualifications of the preliminary data are given in "[Technical Notes](#)."

Each state vital statistics office reported to NCHS the number of deaths registered and processed for calendar year 2008. Those state counts were used as independent control counts for NCHS's 2008 preliminary national mortality file. A comparison of a) the number of 2008 death records received from the states for processing by NCHS with b) the states' independent control counts of the number of deaths in 2008 indicates that demographic information for the United States was available for an estimated 99 percent of infant deaths (under age 1 year) and 100 percent of deaths of persons aged 1 year and over occurring in calendar year 2008 (see [Table I](#) in "[Technical Notes](#)"). Medical (or cause-of-death) information was processed separately and available for an estimated 99 percent of infant deaths and 99 percent of deaths of persons aged 1 year and over in 2008.

To produce the preliminary estimates shown in this report, 2008 records were weighted using 2008 state-specific, independent control counts of infant deaths and deaths of those aged 1 year and over received in state vital statistics offices. Two separate sets of weights were applied to the death records—one set for demographic information and another for medical information. This results in inconsistencies between demographic data from the mortality demographic tables and the medical tables showing causes of death (see "Nature and sources of data" in "[Technical Notes](#)"). Preliminary estimates are subject to sampling variation as well as random variation.

Cause-of-death information is not always available when preliminary data are sent to NCHS but is available later for final data processing. As a result, estimates of cause of death based on preliminary mortality data may differ from statistics developed from the final mortality data (see [Tables II](#) and [III](#) in "[Technical Notes](#)"). Such differences may affect certain causes of death where the cause is pending

investigation, such as for Assault (homicide), Intentional self-harm (suicide), Accidents (unintentional injuries), Drug-induced deaths, and Sudden infant death syndrome (SIDS); see “Nonsampling error” in [“Technical Notes.”](#)

This preliminary report includes national and state estimates of total deaths and death rates, as well as statistics on life expectancy, infant mortality, and causes of death. Data are shown for the following race and ethnic groups: white, non-Hispanic white, black, non-Hispanic black, American Indian or Alaska Native (AIAN), Asian or Pacific Islander (API), and Hispanic populations. Tabulations by race and ethnic group are based on the race and ethnic group reported for the decedent. Race and Hispanic origin are reported as separate items on the death certificate. Death rates for AIAN, API, and, to a lesser extent, Hispanic populations are known to be too low because of reporting problems (see “Race and Hispanic origin” in [“Technical Notes.”](#)).

Changes in death rates from 2007 to 2008 were tested for statistical significance. Differences in death rates across demographic groups (but occurring in 2008 only) were also tested for statistical significance. Unless otherwise specified, reported differences in death rates are statistically significant.

Age-adjusted death rates are better indicators than crude death rates for showing changes in the risk of death over time when the age distribution of the population is changing, and for comparing the mortality of population subgroups that have different age compositions. All age-adjusted death rates are standardized to the year 2000 population (see “Computing rates and percentages” in [“Technical Notes.”](#)).

Two measures of infant mortality are shown: the infant death rate and the infant mortality rate (see “Infant mortality” in [“Technical Notes.”](#)). These measures typically are similar, although they can differ because they have different denominators. The denominator of the 2008 infant death rate is the estimated population under age 1 year on the reference date of July 1, 2008 (14). This estimated population includes a combination of infants born in 2007 who had not reached their first birthday by July 1, 2008, and infants born in 2008 before July 1. In contrast, the denominator of the 2008 infant mortality rate is all live births occurring in 2008. The infant mortality rate is a better indicator of the risk of dying during the first year of life than the infant death rate.

This report includes data for the District of Columbia and 30 states—Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Michigan, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Dakota, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Vermont, Washington, and Wyoming—that had implemented the 2003 revision of the U.S. Standard Certificate of Death by 2008. The remaining 20 states collected and reported death data in 2008 based on the 1989 revision of the U.S. Standard Certificate of Death. The 2003 revision is described in detail elsewhere (15,16). In this report, revised data are combined with unrevised but comparable data. More details on procedures used to combine revised with unrevised data on race are given in [“Technical Notes.”](#)

Because the 2000 U.S. census allowed for selection of multiple races, death certificate data by race (i.e., the numerators for death rates) are not wholly compatible with the population data collected in this census. Data from the census are necessary to produce denominators for computing death rates.

Multiple-race data were collected in 2008 by the 30 states and the District of Columbia that used the 2003 revision of the U.S. Standard

Certificate of Death in a manner that is consistent with the population data collected in the 2000 census. Four additional states—Hawaii, Maine, Minnesota, and Wisconsin—that used the 1989 revision of the U.S. Standard Certificate of Death also reported multiple-race data. The remaining 16 states did not collect multiple-race data in 2008. In order to produce national death rates for 2007 and 2008 for this report, multiple-race data from death certificates were “bridged” to be consistent with 1977 Office of Management and Budget (OMB) single-race categories (17); that is, the multiple-race categories were bridged back to single-race categories. Data for Asian and Native Hawaiian or Other Pacific Islander (NHOP) persons were reported as the combined category API (18). The populations used to calculate death rates are also bridged to single-race categories. These populations are produced under a collaborative arrangement with the U.S. Census Bureau and are based on year 2000 census counts. The procedures used to produce the bridged populations are described in separate publications (19,20). As the remaining 16 reporting areas gradually begin to collect data on race according to the 1997 OMB standards (21), use of the bridged populations is expected to be discontinued.

Note that the population data used to compile the death rates by race shown in this report are based on special estimation procedures. They are not true counts. The estimation procedures used to develop these populations are subject to error. Smaller population groups are affected much more than larger populations by this measurement error, especially the AIAN population (19).

Results

Trends in numbers and rates

The preliminary number of deaths in the United States for 2008 was 2,473,018 ([Tables A and 1](#)). The crude death rate of 813.3 per 100,000 population was 1.2 percent larger than the rate of 803.6 per 100,000 in 2007. The estimated age-adjusted death rate, which accounts for changes in the age distribution of the population, reached a record low of 758.7 per 100,000 U.S. standard population, 0.2 percent lower than the 2007 rate of 760.2 ([Tables A and 1](#)) (13). [Figure 1](#) illustrates the general pattern of decline in both crude and age-adjusted death rates from 1980 through 2008. The age-adjusted death rate decreased from 2007 to 2008 by 0.5 percent for males. The observed age-adjusted death rate increase of less than half of 1 percent for females was not statistically significant. The relative magnitudes of significant changes in age-adjusted death rates by sex, race, and Hispanic origin ([Table 1](#)) are:

- White females—0.4 percent increase
- Non-Hispanic white males—0.2 percent increase
- Non-Hispanic white females—0.5 percent increase
- Black males—2.8 percent decrease
- Black females—1.9 percent decrease
- Non-Hispanic black males—2.7 percent decrease
- Non-Hispanic black females—1.8 percent decrease
- Hispanic males—3.0 percent decrease

Among the major race and Hispanic origin groups, the lowest mortality was reported for the API, Hispanic, and AIAN populations. Compared with the non-Hispanic white population, preliminary

Table A. Deaths, age-adjusted death rates, and life expectancy at birth, by race and sex; and infant deaths and mortality rates, by race: United States, final 2007 and preliminary 2008

[Data are based on a continuous file of records received from the states. Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals]

Measure and sex	All races ¹		White ²		Black ²	
	2008	2007	2008	2007	2008	2007
All deaths	2,473,018	2,423,712	2,120,961	2,074,151	289,306	289,585
Male	1,226,721	1,203,968	1,046,532	1,023,951	147,275	148,309
Female	1,246,297	1,219,744	1,074,429	1,050,200	142,031	141,276
Age-adjusted death rate ³	758.7	760.2	750.6	749.4	935.6	958.0
Male	901.0	905.6	889.5	890.5	1,151.3	1,184.4
Female	643.7	643.4	637.1	634.8	779.0	793.8
Life expectancy at birth (in years) ⁴	77.8	77.9	78.2	78.4	73.6	73.6
Male	75.3	75.4	75.7	75.9	70.2	70.0
Female	80.3	80.4	80.6	80.8	76.8	76.8
All infant deaths	28,033	29,138	18,162	18,807	8,513	8,944
Infant mortality rate ⁵	6.59	6.75	5.54	5.64	12.68	13.24

¹Includes races other than white and black.

²Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported in 2008 for Arkansas, California, Connecticut, Delaware, District of Columbia, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Dakota, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Vermont, Washington, Wisconsin, and Wyoming; and in 2007 for California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. Multiple-race data for these reporting areas were bridged to single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

³Per 100,000 U.S. standard population, based on the year 2000 standard.

⁴Life expectancies were calculated using a revised methodology and may differ from those previously published; see "Technical Notes."

⁵Deaths under age 1 year per 1,000 live births in specified group.

age-adjusted death rates were 46 percent lower for the API population, 30 percent lower for the Hispanic population, and 20 percent lower for the AIAN population. In contrast, the age-adjusted death rate for the non-Hispanic black population was 25 percent higher than that for the non-Hispanic white population (Table 1). It is important to keep in mind, however, that mortality for races other than white and black may be seriously understated in some cases due to underreporting for some race groups and Hispanic origin on death certificates (22–24).




Figure 1. Crude and age-adjusted death rates: United States, 1980–2007 final and 2008 preliminary

Statistically significant decreases in mortality from 2007 to 2008 were registered for those under age 1 year and across age groups 5–44 and 65–74. The death rate increased for those aged 85 and over, while other age groups did not experience significant change. The magnitude of the significant changes in mortality by age group is (Table 1):

- Under 1 year—5.1 percent decrease
- 5–14 years—7.8 percent decrease
- 15–24 years—5.3 percent decrease
- 25–34 years—1.4 percent decrease
- 35–44 years—2.5 percent decrease
- 65–74 years—0.7 percent decrease
- 85 years and over—0.6 percent increase

The death rate for "under 1 year" is based on a population estimate and is different from the infant mortality rate, which is based on live births (see "Infant mortality").

Life expectancy data shown in this report for data years 2007–2008 are based on a newly revised methodology, which is similar to that developed for the 1999–2001 decennial life tables; see "Technical Notes." The preliminary estimate of life expectancy at birth for the total population in 2008 is 77.8 years. This represents a decrease in life expectancy of 0.1 year relative to 2007 (Tables A and 6). Life expectancy for males and females decreased about one-tenth of a year for each group from 2007 to 2008: from 75.4 to 75.3 years for males in 2008, and from 80.4 to 80.3 years for females. The difference between male and female life expectancy at birth has been generally decreasing since its peak of 7.8 years in 1979 (25). The gap between male and female life expectancy was 5.0 years in 2008, the same as in 2007. A record-high life expectancy was reached for black males (70.2 years), rising by 0.2 years from 70.0 years in 2007. The difference

in life expectancy between the white and black populations in 2008 was 4.6 years, a 0.2-year decrease from the 2007 gap between the two races (13).

White females have the highest life expectancy (Figure 2), followed by, in order, black females, white males, and black males. Figure 2 also shows that this pattern has not changed from 1976 through 2008, even though life expectancy for all groups has generally increased over this time period.

By state of residence, Hawaii had the lowest mortality in 2008 with an age-adjusted death rate of 589.0 deaths per 100,000 standard population (Table 3). Mortality was highest in West Virginia, with an age-adjusted death rate of 958.1 per 100,000 standard population.

Causes of death

The 15 leading causes of death in 2008 (Table B) were:

1. Diseases of heart
2. Malignant neoplasms
3. Chronic lower respiratory diseases
4. Cerebrovascular diseases
5. Accidents (unintentional injuries)
6. Alzheimer's disease

7. Diabetes mellitus
8. Influenza and pneumonia
9. Nephritis, nephrotic syndrome and nephrosis
10. Septicemia
11. Intentional self-harm (suicide)
12. Chronic liver disease and cirrhosis
13. Essential hypertension and hypertensive renal disease
14. Parkinson's disease
15. Assault (homicide)

Cerebrovascular diseases (stroke) had been the third leading cause of death in the United States for over five decades, continually trailing Diseases of heart and Malignant neoplasms. In 2008, however, Chronic lower respiratory diseases (CLRD) replaced stroke as the third leading cause of death. Stroke is now the fourth leading cause of death. The change in the ranks of these two causes of death is mostly driven by a decreasing trend in the numbers of deaths from stroke (from 167,661 deaths in 2000 to 133,750 in 2008) (Figure 3). A corresponding increase in the number of deaths from CLRD during this period, particularly from 2007 to 2008, also helped to bring about this transposition in ranks.

The increase in deaths from CLRD from 2007 to 2008 needs to be interpreted with caution, however. Changes were made in 2008 to




Figure 2. Life expectancy at birth, by race and sex: United States, 1975–2007 final and 2008 preliminary

Table B. Deaths and death rates for 2008 and age-adjusted death rates and percent changes in age-adjusted rates from 2007 to 2008 for the 15 leading causes of death: United States, final 2007 and preliminary 2008

[Data are based on a continuous file of records received from the states. Rates are per 100,000 population; age-adjusted rates per 100,000 U.S. standard population based on the year 2000 standard; see "Technical Notes." For explanation of asterisks (*) preceding cause-of-death codes, see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals]

Rank ¹	Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Number	Death rate	Age-adjusted death rate		
				2008	2007	Percent change
...	All causes	2,472,699	813.2	758.6	760.2	-0.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	617,527	203.1	186.7	190.9	-2.2
2	Malignant neoplasms (C00-C97)	566,137	186.2	175.5	178.4	-1.6
3	Chronic lower respiratory diseases (J40-J47)	141,075	46.4	44.0	40.8	7.8
4	Cerebrovascular diseases (I60-I69)	133,750	44.0	40.6	42.2	-3.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86) ²	121,207	39.9	38.6	40.0	-3.5
6	Alzheimer's disease (G30)	82,476	27.1	24.4	22.7	7.5
7	Diabetes mellitus (E10-E14)	70,601	23.2	21.8	22.5	-3.1
8	Influenza and pneumonia (J09-J18) ³	56,335	18.5	17.0	16.2	4.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48,283	15.9	14.8	14.5	2.1
10	Septicemia (A40-A41)	35,961	11.8	11.1	11.0	0.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) ²	35,933	11.8	11.6	11.3	2.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29,963	9.9	9.2	9.1	1.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25,823	8.5	7.7	7.4	4.1
14	Parkinson's disease (G20-G21)	20,507	6.7	6.4	6.4	0.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) ²	17,837	5.9	5.9	6.1	-3.3
...	All other causes (residual)	469,284	154.3

... Category not applicable.

¹Based on number of deaths.

²Preliminary and final data may differ significantly because of the truncated nature of the preliminary file.

³New code J09 (Influenza due to identified avian influenza virus) was added to the category in 2007.

NOTES: Data are subject to sampling and random variation. For information regarding calculation of standard errors and further discussion of variability of the data, see "Technical Notes."

the way that CLRDs are coded and classified, but it is unclear at this point to what extent these changes contributed to the increase shown for this cause of death (7.8 percent). NCHS periodically revises its cause-of-death coding practices to reflect and follow changes to coding rules made by the World Health Organization (WHO). Starting with 2008, death certificates that independently mention pneumonia in conjunction with a chronic lower respiratory problem are now classified as "Chronic obstructive pulmonary disease with acute lower respiratory

infection," which is one of the conditions that make up the larger category known as CLRD. This condition has been underused in the past, and the recent change aims to address this underuse (see "Technical Notes"). Preliminary analysis of death certificates classified to CLRD suggest that, had this change not been implemented, a substantial number of these deaths would have been counted as deaths from pneumonia. The final report on mortality for 2008 will include a more detailed analysis of the effects that this change in coding and classification had on counts of deaths from CLRD.

From 2007 to 2008, the age-adjusted death rate declined significantly for 6 of the 15 leading causes of death. The preliminary age-adjusted death rate for the leading cause of death, Diseases of heart, decreased by 2.2 percent. The age-adjusted death rate for Malignant neoplasms decreased by 1.6 percent (Tables B and 2). Deaths from these two diseases combined accounted for 48.0 percent of deaths in the United States in 2008. Although heart disease mortality has exhibited a downward trend since 1950, cancer mortality began to decline only in the early 1990s (10,13). The preliminary age-adjusted death rate also decreased significantly for stroke (3.8 percent).

Other leading causes of death that showed significant decreases in 2008 relative to 2007 were: Accidents (unintentional injuries), 3.5 percent; Diabetes mellitus, 3.1 percent; and Assault (homicide), 3.3 percent.

The age-adjusted death rates for CLRD increased significantly (by 7.8 percent) between 2007 and 2008. As explained above, this single-year increase must be interpreted with caution until the coding changes that affected this condition can be evaluated. The preliminary age-adjusted death rate also increased for Alzheimer's disease (7.5 percent); Influenza and pneumonia (4.9 percent); Nephritis, nephrotic




Figure 3. Deaths from Cerebrovascular diseases and Chronic lower respiratory diseases: United States, 1999–2007 final and 2008 preliminary

syndrome and nephrosis (2.1 percent); Intentional self-harm (suicide) (2.7 percent); and Essential hypertension and hypertensive renal disease (4.1 percent).

Observed increases for the age-adjusted death rates for Septicemia and for Chronic liver disease and cirrhosis were not significant. The age-adjusted death rate for Parkinson's disease (14th leading cause of death) remained unchanged from 2007 to 2008.

Human immunodeficiency virus (HIV) disease was not among the 15 leading causes of death in 2008. The preliminary age-adjusted death rate for HIV disease declined by 10.8 percent from 2007 to 2008 (Table 2). Following a period of increase from 1987 through 1994, HIV disease mortality reached a plateau in 1995. Subsequently, the rate for this disease decreased an average of 33.0 percent per year from 1995 through 1998, and 4.4 percent per year from 1999 through 2007 (13). For all races combined in the age group 15–24, HIV disease was the 10th leading cause of death in 2008, climbing up one position relative to its rank as 11th leading cause for those aged 15–24 in 2007. HIV disease remained the sixth leading cause of death for the age group 25–44, unchanged in rank from 2007. Among decedents aged 45–64, HIV disease dropped from 11th leading cause in 2007 to 12th leading cause.

Enterocolitis due to *Clostridium difficile* (*C. difficile*)—a predominantly antibiotic-associated inflammation of the intestines caused by *C. difficile*, a gram-positive, anaerobic, spore-forming bacillus—is of growing concern. The disease is often acquired in hospitals or other health-care facilities with long-term patients or residents and accounts for an increasing number of deaths (26,27). In 1999, 793 deaths were due to *C. difficile*, compared with 6,372 *C. difficile* deaths in 2007 (13) and 7,483 deaths in 2008. The age-adjusted death rate for this cause increased from 2.0 deaths per 100,000 standard population in 2007 to 2.3 deaths per 100,000 standard population in 2008 (15.0 percent). In 2008, *C. difficile* ranked as the 18th leading cause of death for the population aged 65 and over. Approximately 93 percent of deaths from *C. difficile* occurred to people aged 65 and older (data not shown).

The preliminary age-adjusted death rate for drug-induced deaths declined by 2.4 percent from 12.6 in 2007 to 12.3 in 2008. However, the final number of drug-induced deaths in 2008 may be substantially higher because information on cause of death in these cases is often delayed pending investigation. Additional information based on toxicology or autopsy reports is often not available in the preliminary file. The age-adjusted death rate for injury at work declined 9.5 percent from 2.1 to 1.9. The age-adjusted death rate increased for alcohol-induced deaths by 2.8 percent. The observed increase in mortality from firearm injuries (Table 2) is not statistically significant.

Infant mortality

The preliminary infant mortality rate for 2008 was 6.59 infant deaths per 1,000 live births (Tables A and 4). This represents a decrease of 2.4 percent from the 2007 rate of 6.75. With the exception of 2002 and 2005, the infant mortality rate has statistically remained the same or decreased significantly each successive year from 1958 through 2008 (13,28). The neonatal (i.e., infants aged less than 28 days) mortality rate decreased 3.4 percent, from 4.42 per 1,000 live births in 2007 to 4.27 per 1,000 live births in 2008. The postneonatal (i.e., infants aged 28 days or more) mortality rate did not change significantly between 2007 and 2008.

The 2008 preliminary infant mortality rate for black infants was 12.68 infant deaths per 1,000 live births, compared with 13.24 per 1,000 live births in 2007. This represents a decrease of 4.2 percent.

The infant mortality rate for white infants in 2008 was 5.54 per 1,000 live births, an insignificant decrease compared with the rate of 5.64 per 1,000 live births in 2007. The mortality rate for black infants was 2.3 times the rate for white infants (Tables A and 4). Because of inconsistencies in the reporting of race groups on birth and death certificates (especially for races other than white and black and for Hispanic origin), infant mortality rates for these groups are likely to be underestimated (24). The linked birth/infant death data set provides a better source of data for infant deaths and mortality rates by maternal race and ethnicity (29).

Although the infant mortality rate is the preferred indicator of the risk of dying during the first year of life, the infant death rate is also shown in this report. While similar, these two rates vary based on differences in their denominators. The denominator of the 2008 infant death rate is the estimated population under age 1 year as of the reference date, July 1, 2008 (14). This population estimate includes a combination of infants born in 2007 who had not reached their first birthday before July 1, 2008, and infants born in 2008 before July 1, 2008. In contrast, the denominator of the 2008 infant mortality rate is all live births occurring during 2008 (30). For example, the preliminary number of live births for 2008 ($n = 4,251,095$) is 1.4 percent lower than the midyear infant population in 2008 ($n = 4,313,132$). Therefore, the infant mortality rate for 2008 (659.4 deaths per 100,000 live births) is higher than the infant death rate for 2008 (649.9 deaths per 100,000 population). For 2008, both the infant mortality rate and the infant death rate decreased significantly from 2007.

The 10 leading causes of infant mortality for 2008 were:

1. Congenital malformations, deformations and chromosomal abnormalities
2. Disorders related to short gestation and low birth weight, not elsewhere classified
3. Sudden infant death syndrome
4. Newborn affected by maternal complications of pregnancy
5. Accidents (unintentional injuries)
6. Newborn affected by complications of placenta, cord and membranes
7. Bacterial sepsis of newborn
8. Respiratory distress of newborn
9. Diseases of the circulatory system
10. Neonatal hemorrhage

The only difference between the leading causes for 2007 (13) and those for 2008 (Table 8) is that Diseases of the circulatory system and Neonatal hemorrhage exchanged ranks, the former moving from 10th to 9th and the latter moving from 9th to 10th.

The infant mortality rate decreased for 2 of 10 leading causes of death from 2007 to 2008 (Table 8). The infant mortality rate for Bacterial sepsis of newborn—the seventh leading cause of death—decreased by 13.7 percent (Tables 5 and 8). The infant mortality rate for Respiratory distress of newborn—the eighth leading cause of death—decreased by 19.7 percent.

Deaths due to SIDS, currently the third leading cause of infant death, have been declining since 1988 (4). The observed decrease in SIDS from 56.8 to 53.9 is not statistically significant. Because SIDS

deaths often involve lengthy investigations, the mortality rate due to SIDS is typically lower based on preliminary data than that based on the final data. Recent declines in mortality due to SIDS also may reflect primarily a change in the way SIDS is diagnosed and reported by medical examiners and coroners (31).

Observed declines in the infant mortality rate for: Congenital malformations, deformations and chromosomal abnormalities (0.9 percent); Disorders related to short gestation and low birth weight, not elsewhere classified (1.1 percent); Newborn affected by complications of placenta, cord and membranes (4.2 percent); Diseases of the circulatory system (4.1 percent); and Neonatal hemorrhage (5.8 percent) were not statistically significant (Tables 5 and 8).

Observed increases in the infant mortality rate for Newborn affected by maternal complications of pregnancy (1.2 percent) and Accidents (unintentional injuries) (2.7 percent) were not statistically significant (Tables 5 and 8).

References

- Anderson RN, Kochanek KD, Murphy SL. Report of final mortality statistics, 1995. Monthly vital statistics report; vol 45 no 11 suppl 2. Hyattsville, MD: National Center for Health Statistics. 1997. Available from: http://www.cdc.gov/nchs/data/mvstr/supp/mv45_11s2.pdf.
- Peters KD, Kochanek KD, Murphy SL. Deaths: Final data for 1996. National vital statistics reports; vol 47 no 9. Hyattsville, MD: National Center for Health Statistics. 1998. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr47/nvsr47_09.pdf.
- Hoyert DL, Kochanek KD, Murphy SL. Deaths: Final data for 1997. National vital statistics reports; vol 47 no 19. Hyattsville, MD: National Center for Health Statistics. 1999. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr47/nvsr47_19.pdf.
- Murphy SL. Deaths: Final data for 1998. National vital statistics reports; vol 48 no 11. Hyattsville, MD: National Center for Health Statistics. 2000. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr48/nvsr48_11.pdf.
- Hoyert DL, Arias E, Smith BL, et al. Deaths: Final data for 1999. National vital statistics reports; vol 49 no 8. Hyattsville, MD: National Center for Health Statistics. 2001. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr49/nvsr49_08.pdf.
- Miniño AM, Arias E, Kochanek KD, et al. Deaths: Final data for 2000. National vital statistics reports; vol 50 no 15. Hyattsville, MD: National Center for Health Statistics. 2002. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr50/nvsr50_15.pdf.
- Arias E, Anderson RN, Kung HC, et al. Deaths: Final data for 2001. National vital statistics reports; vol 52 no 3. Hyattsville, MD: National Center for Health Statistics. 2003. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr52/nvsr52_03.pdf.
- Kochanek KD, Murphy SL, Anderson RN, Scott C. Deaths: Final data for 2002. National vital statistics reports; vol 53 no 5. Hyattsville, MD: National Center for Health Statistics. 2004. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr53/nvsr53_05.pdf.
- Hoyert DL, Heron MP, Murphy SL, Kung HC. Deaths: Final data for 2003. National vital statistics reports; vol 54 no 13. Hyattsville, MD: National Center for Health Statistics. 2006. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr54/nvsr54_13.pdf.
- Miniño AM, Heron MP, Murphy SL, Kochanek KD. Deaths: Final data for 2004. National vital statistics reports; vol 55 no 19. Hyattsville, MD: National Center for Health Statistics. 2007. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr55/nvsr55_19.pdf.
- Kung HC, Hoyert DL, Xu J, Murphy SL. Deaths: Final data for 2005. National vital statistics reports; vol 56 no 10. Hyattsville, MD: National Center for Health Statistics. 2008. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr56/nvsr56_10.pdf.
- Heron M, Hoyert DL, Murphy SL, et al. Deaths: Final data for 2006. National vital statistics reports; vol 57 no 14. Hyattsville, MD: National Center for Health Statistics. 2009. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr57/nvsr57_14.pdf.
- Xu J, Kochanek KD, Murphy SL, Tejada-Vera B. Deaths: Final data for 2007. National vital statistics reports; vol 58 no 19. Hyattsville, MD: National Center for Health Statistics. 2010. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_19.pdf.
- National Center for Health Statistics. Vintage 2008 bridged-race postcensal population estimates of the resident population of the United States for July 1, 2005–July 1, 2008, by year, county, single-year of age, bridged race, Hispanic origin, and sex (pcen_v2008_y0508.txt). Prepared under a collaborative agreement with the U.S. Census Bureau. 2009. Available from: http://www.cdc.gov/nchs/nvss/bridged_race/data_documentation.htm#vintage2008.
- National Center for Health Statistics. 2003 revision of the U.S. Standard Certificate of Death. 2003. Available from: <http://www.cdc.gov/nchs/data/dvs/DEATH11-03final-acc.pdf>.
- National Center for Health Statistics. Report of the panel to evaluate the U.S. standard certificates. 2000. Available from: http://www.cdc.gov/nchs/data/dvs/panelreport_acc.pdf.
- Office of Management and Budget. Race and ethnic standards for federal statistics and administrative reporting. Statistical Policy Directive 15. May 12, 1977.
- U.S. Census Bureau. Age, sex, race and Hispanic origin information from the 1990 census: A comparison of census results with results where age and race have been modified, 1990. CPH-L-74. Washington, DC: U.S. Department of Commerce. 1991.
- Ingram DD, Parker JD, Schenker N, et al. United States census 2000 population with bridged race categories. National Center for Health Statistics. Vital Health Stat 2(135). 2003.
- Schenker N, Parker JD. From single-race reporting to multiple-race reporting: Using imputation methods to bridge the transition. *Stat Med* 22(9):1571–87. 2003.
- Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. *Fed Regist* (62FR58782–58790). October 30, 1997.
- Sorlie PD, Rogot E, Johnson NJ. Validity of demographic characteristics on the death certificate. *Epidemiology* 3(2):181–4. 1992.
- Rosenberg HM, Maurer JD, Sorlie PD, et al. Quality of death rates by race and Hispanic origin: A summary of current research, 1999. National Center for Health Statistics. *Vital Health Stat* 2(128). 1999.
- Arias E, Schauman WS, Eschbach K, et al. The validity of race and Hispanic origin reporting on death certificates in the United States. National Center for Health Statistics. *Vital Health Stat* 2(148):1–23. 2008. Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_148.pdf.
- Arias E. United States life tables, 2006. National vital statistics reports; vol 58 no 21. Hyattsville, MD: National Center for Health Statistics. 2010. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_21.pdf.
- Sunenshine RH, McDonald LC. Clostridium difficile-associated disease: New challenges from an established pathogen. *Cleve Clin J Med* 73(2):187–97. 2006. Available from: http://www.cdc.gov/ncidod/dhqp/pdf/infDis/Cdiff_CCJM02_06.pdf.
- Redelings MD, Sorvillo F, Mascola L. Increase in Clostridium difficile-related mortality rates, United States, 1999–2004. *Emerg Infect Dis* [online series] 13(9):1417–9. 2007. Available from: <http://www.cdc.gov/EID/content/13/9/1417.htm>.

28. Kochanek KD, Martin JA. Supplemental analyses of recent trends in infant mortality. Health E-stat. Hyattsville, MD: National Center for Health Statistics. 2003. Available from: <http://www.cdc.gov/nchs/products/pubs/pubd/hestats/infantmort/infantmort.htm>.
29. Mathews TJ, MacDorman MF. Infant mortality statistics from the 2006 period linked birth/infant death data set. National vital statistics reports; vol 58 no 17. Hyattsville, MD: National Center for Health Statistics. 2010. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_17.pdf.
30. Hamilton BE, Martin JA, Ventura SJ. Births: Preliminary data for 2008. National vital statistics reports; vol 58 no 16. Hyattsville, MD: National Center for Health Statistics. 2010. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_16.pdf.
31. Shapiro-Mendoza CK, Tomashek KM, Anderson RN, Wingo J. Recent national trends in sudden, unexpected infant deaths: More evidence supporting a change in classification or reporting. Am J Epidemiol 163(8):762–9. 2006.
32. National Center for Health Statistics. Technical appendix. Vital statistics of the United States: Mortality, 1999. Available from: <http://www.cdc.gov/nchs/data/statab/techap99.pdf>.
33. National Center for Health Statistics. Vital statistics, instructions for classifying the underlying cause of death. NCHS instruction manual, part 2a. Hyattsville, MD. Published annually.
34. World Health Organization. International Statistical Classification of Diseases and Related Health Problems, Tenth Revision. Geneva. 1992.
35. World Health Organization. International Statistical Classification of Diseases and Related Health Problems, Tenth Revision, Second Edition. Geneva. 2004.
36. World Health Organization. Manual of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Ninth Revision. Geneva. 1977.
37. Anderson RN, Miniño AM, Hoyert DL, Rosenberg HM. Comparability of cause of death between ICD–9 and ICD–10: Preliminary estimates. National vital statistics reports; vol 49 no 2. Hyattsville, MD: National Center for Health Statistics. 2001. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr49/nvsr49_02.pdf.
38. National Center for Health Statistics. ICD–10 cause-of-death lists for tabulating mortality statistics (updated October 2007 to include WHO updates to ICD–10 for data year 2007). NCHS instruction manual, part 9. Hyattsville, MD. 2007. Available from: http://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
39. National Center for Health Statistics. ICD–10 ACME Decision Tables for Classifying Underlying Causes of Death, 2008. NCHS instruction manual, part 2c. Hyattsville, MD: Public Health Service. 2008. Available from: http://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
40. Heron, M. Deaths: Leading causes for 2006. National vital statistics reports; vol 58 no 14. Hyattsville, MD: National Center for Health Statistics. 2010. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_14.pdf.
41. Sirken M. Comparison of two methods of constructing abridged life tables by reference to a “standard” table. National Center for Health Statistics. Vital Health Stat 2(4). 1966 [revised]. Available from: www.cdc.gov/nchs/data/series/sr_02/sr02_004.pdf.
42. Anderson RN. Method for constructing complete annual U.S. life tables. National Center for Health Statistics. Vital Health Stat 2(129). 1999. Available from: www.cdc.gov/nchs/data/series/sr_02/sr02_129.pdf.
43. National Center for Health Statistics. U.S. decennial life tables for 1989–91, vol 1 no 2, methodology of the national and state life tables. Hyattsville, MD. 1998. Available from: www.cdc.gov/nchs/data/lifetables/life89_1_2.pdf.
44. Kestenbaum B. A description of the extreme aged population based on improved Medicare enrollment data. Demography 29(4):565–80. 1992.
45. Wei R, Curtin LR, Arias E, Anderson RN. U.S. decennial life tables for 1999–2001: Methodology of the United States life tables. National vital statistics reports; vol 57 no 4. Hyattsville, MD: National Center for Health Statistics. 2008. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr57/nvsr57_04.pdf.
46. Arias E, Rostron BL, Tejada-Vera B. United States Life Tables, 2005. National vital statistics reports; vol 58 no 10. Hyattsville, MD: National Center for Health Statistics. 2010. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_10.pdf.
47. National Center for Health Statistics. Vintage 2007 bridged-race postcensal population estimates of the resident population of the United States for July 1, 2000–July 1, 2007, by year, county, single-year of age, bridged race, Hispanic origin, and sex (pcen_v2007.txt). Prepared under a collaborative agreement with the U.S. Census Bureau. 2008. Available from: http://www.cdc.gov/nchs/nvss/bridged_race/data_documentation.htm.
48. Anderson RN, Rosenberg HM. Age standardization of death rates: Implementation of the year 2000 standard. National vital statistics reports; vol 47 no 3. Hyattsville, MD: National Center for Health Statistics. 1998. Available from: www.cdc.gov/nchs/data/nvsr/nvsr47/nvsr47_03.pdf.

List of Detailed Tables

1. Deaths and death rates, by age, sex, and race and Hispanic origin, and age-adjusted death rates, by sex and race and Hispanic origin: United States, final 2007 and preliminary 2008	9
2. Deaths, death rates, and age-adjusted death rates for 113 selected causes, Injury by firearms, Drug-induced deaths, Alcohol-induced deaths, Injury at work, and Enterocolitis due to <i>Clostridium difficile</i> : United States, final 2007 and preliminary 2008	17
3. Deaths, death rates, and age-adjusted death rates: United States, and each state and territory, final 2007 and preliminary 2008	21
4. Infant deaths and infant mortality rates, by age and race and Hispanic origin: United States, final 2007 and preliminary 2008	22
5. Infant deaths and infant mortality rates for 130 selected causes: United States, final 2007 and preliminary 2008	23
6. Expectation of life, by age, race, and sex: United States, final 2007 and preliminary 2008	27
7. Deaths and death rates for the 10 leading causes of death in specified age groups: United States, preliminary 2008	29
8. Infant deaths and infant mortality rates for the 10 leading causes of infant death, by race and Hispanic origin: United States, preliminary 2008	32

Table 1. Deaths and death rates, by age, sex, and race and Hispanic origin, and age-adjusted death rates, by sex and race and Hispanic origin: United States, final 2007 and preliminary 2008

[Data are based on a continuous file of records received from the states. Age-specific rates are per 100,000 population in specified group. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals. Race and Hispanic origin are reported separately on the death certificate. Data for Hispanic origin and specified races other than white and black should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 34 states and the District of Columbia in 2008 and by 27 states and the District of Columbia in 2007; see "Technical Notes." The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes." Data for persons of Hispanic origin are included in the data for each race group, according to the decedent's reported race; see "Technical Notes"]

Age, sex, and race and Hispanic origin	2008		2007	
	Number	Rate	Number	Rate
All races, both sexes				
All ages	2,473,018	813.3	2,423,712	803.6
Under 1 year ¹	28,033	649.9	29,138	684.5
1-4 years	4,747	28.4	4,703	28.6
5-14 years	5,663	14.1	6,147	15.3
15-24 years	32,208	75.7	33,982	79.9
25-34 years	42,309	103.4	42,572	104.9
35-44 years	76,409	179.8	79,606	184.4
45-54 years	186,643	420.6	184,686	420.9
55-64 years	296,269	879.5	287,110	877.7
65-74 years	401,750	1,996.5	389,238	2,011.3
75-84 years	653,845	5,019.9	652,682	5,011.6
85 years and over	744,976	13,020.0	713,647	12,946.5
Not stated	166	...	201	...
Age-adjusted rate ²	758.7	...	760.2
All races, male				
All ages	1,226,721	818.2	1,203,968	809.9
Under 1 year ¹	15,651	708.9	16,293	747.8
1-4 years	2,703	31.7	2,634	31.3
5-14 years	3,284	16.0	3,585	17.4
15-24 years	24,030	109.9	25,316	115.8
25-34 years	29,662	141.9	29,792	144.0
35-44 years	47,717	223.9	50,105	231.8
45-54 years	115,098	526.7	114,456	530.0
55-64 years	179,599	1,105.2	173,618	1,100.6
65-74 years	225,512	2,434.0	218,344	2,456.9
75-84 years	322,031	6,034.7	320,803	6,038.4
85 years and over	261,319	14,022.5	248,866	14,006.4
Not stated	115	...	156	...
Age-adjusted rate ²	901.0	...	905.6
All races, female				
All ages	1,246,297	808.6	1,219,744	797.4
Under 1 year ¹	12,382	588.1	12,845	618.1
1-4 years	2,044	25.1	2,069	25.7
5-14 years	2,379	12.1	2,562	13.1
15-24 years	8,178	39.5	8,666	42.0
25-34 years	12,647	63.1	12,780	64.2
35-44 years	28,692	135.4	29,501	136.9
45-54 years	71,546	317.7	70,230	315.2
55-64 years	116,670	669.2	113,492	670.1
65-74 years	176,238	1,623.1	170,894	1,633.0
75-84 years	331,814	4,315.6	331,879	4,304.1
85 years and over	483,656	12,535.8	464,781	12,442.3
Not stated	51	...	45	...
Age-adjusted rate ²	643.7	...	643.4

See footnotes at end of table.

Table 1. Deaths and death rates, by age, sex, and race and Hispanic origin, and age-adjusted death rates, by sex and race and Hispanic origin: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Age-specific rates are per 100,000 population in specified group. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals. Race and Hispanic origin are reported separately on the death certificate. Data for Hispanic origin and specified races other than white and black should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 34 states and the District of Columbia in 2008 and by 27 states and the District of Columbia in 2007; see "Technical Notes." The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes." Data for persons of Hispanic origin are included in the data for each race group, according to the decedent's reported race; see "Technical Notes"]

Age, sex, and race and Hispanic origin	2008		2007	
	Number	Rate	Number	Rate
Total white, both sexes				
All ages	2,120,961	864.9	2,074,151	851.5
Under 1 year ¹	18,162	549.8	18,807	573.7
1–4 years	3,350	26.0	3,287	25.8
5–14 years	4,072	13.0	4,480	14.4
15–24 years	23,627	71.2	25,137	75.7
25–34 years	31,123	96.9	31,181	98.0
35–44 years	57,627	170.4	59,571	172.8
45–54 years	144,834	397.6	142,233	394.0
55–64 years	238,951	840.6	231,287	835.4
65–74 years	338,817	1,959.2	327,377	1,970.3
75–84 years	579,658	5,051.9	579,122	5,030.8
85 years and over	680,608	13,288.2	651,521	13,176.5
Not stated	130	...	148	...
Age-adjusted rate ²	750.6	...	749.4
White male				
All ages	1,046,532	860.6	1,023,951	848.1
Under 1 year ¹	10,151	600.2	10,540	627.8
1–4 years	1,920	29.1	1,847	28.3
5–14 years	2,340	14.6	2,590	16.2
15–24 years	17,546	102.6	18,513	108.1
25–34 years	22,075	133.2	21,998	134.2
35–44 years	36,677	213.6	38,121	218.2
45–54 years	90,857	500.8	89,556	498.4
55–64 years	145,915	1,051.9	140,785	1,042.7
65–74 years	191,321	2,377.5	184,828	2,396.7
75–84 years	287,530	6,054.3	286,824	6,049.2
85 years and over	240,108	14,358.6	228,227	14,286.4
Not stated	93	...	122	...
Age-adjusted rate ²	889.5	...	890.5
White female				
All ages	1,074,429	869.0	1,050,200	854.9
Under 1 year ¹	8,011	496.8	8,267	516.8
1–4 years	1,430	22.8	1,440	23.1
5–14 years	1,733	11.4	1,890	12.4
15–24 years	6,081	37.8	6,624	41.2
25–34 years	9,049	58.2	9,183	59.6
35–44 years	20,949	125.8	21,450	126.2
45–54 years	53,978	295.2	52,677	290.5
55–64 years	93,037	639.2	90,502	638.0
65–74 years	147,496	1,595.1	142,549	1,600.9
75–84 years	292,128	4,344.1	292,298	4,317.6
85 years and over	440,501	12,769.4	423,294	12,646.7
Not stated	37	...	26	...
Age-adjusted rate ²	637.1	...	634.8

See footnotes at end of table.

Table 1. Deaths and death rates, by age, sex, and race and Hispanic origin, and age-adjusted death rates, by sex and race and Hispanic origin: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Age-specific rates are per 100,000 population in specified group. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals. Race and Hispanic origin are reported separately on the death certificate. Data for Hispanic origin and specified races other than white and black should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 34 states and the District of Columbia in 2008 and by 27 states and the District of Columbia in 2007; see "Technical Notes." The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes." Data for persons of Hispanic origin are included in the data for each race group, according to the decedent's reported race; see "Technical Notes"]

Age, sex, and race and Hispanic origin	2008		2007	
	Number	Rate	Number	Rate
Non-Hispanic white, both sexes				
All ages	1,981,198	982.0	1,939,606	964.1
Under 1 year ¹	12,545	544.0	12,998	559.7
1–4 years	2,380	26.2	2,325	25.5
5–14 years	3,082	13.2	3,362	14.2
15–24 years	18,570	70.7	19,607	74.5
25–34 years	24,868	101.6	24,757	102.4
35–44 years	49,109	180.0	50,966	181.5
45–54 years	130,455	410.4	127,898	403.8
55–64 years	220,247	854.9	213,876	849.6
65–74 years	316,695	1,994.5	306,342	2,006.2
75–84 years	549,133	5,143.4	549,752	5,113.1
85 years and over	654,028	13,556.3	627,642	13,413.3
Not stated	88	...	81	...
Age-adjusted rate ²	766.2	...	763.3
Non-Hispanic white male				
All ages	969,419	978.4	948,662	960.4
Under 1 year ¹	7,044	596.3	7,340	616.8
1–4 years	1,367	29.3	1,316	28.1
5–14 years	1,811	15.1	1,945	16.1
15–24 years	13,578	100.8	14,129	104.6
25–34 years	17,348	140.6	17,138	140.8
35–44 years	30,831	225.5	32,139	228.4
45–54 years	81,342	515.9	79,931	508.7
55–64 years	134,286	1,067.2	129,977	1,057.5
65–74 years	178,750	2,413.2	172,867	2,432.7
75–84 years	272,518	6,167.3	272,398	6,152.7
85 years and over	230,484	14,721.1	219,419	14,588.3
Not stated	58	...	63	...
Age-adjusted rate ²	908.6	...	906.8
Non-Hispanic white female				
All ages	1,011,780	985.6	990,944	967.6
Under 1 year ¹	5,500	488.9	5,658	499.6
1–4 years	1,012	22.8	1,009	22.7
5–14 years	1,271	11.1	1,417	12.3
15–24 years	4,993	39.1	5,478	42.7
25–34 years	7,520	62.0	7,619	63.4
35–44 years	18,278	134.3	18,827	134.4
45–54 years	49,112	306.6	47,967	300.5
55–64 years	85,961	652.3	83,899	651.3
65–74 years	137,944	1,628.4	133,475	1,634.9
75–84 years	276,614	4,420.4	277,354	4,385.4
85 years and over	423,543	12,996.7	408,223	12,856.7
Not stated	30	...	18	...
Age-adjusted rate ²	650.9	...	647.7

See footnotes at end of table.

Table 1. Deaths and death rates, by age, sex, and race and Hispanic origin, and age-adjusted death rates, by sex and race and Hispanic origin: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Age-specific rates are per 100,000 population in specified group. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals. Race and Hispanic origin are reported separately on the death certificate. Data for Hispanic origin and specified races other than white and black should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 34 states and the District of Columbia in 2008 and by 27 states and the District of Columbia in 2007; see "Technical Notes." The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes." Data for persons of Hispanic origin are included in the data for each race group, according to the decedent's reported race; see "Technical Notes"]

Age, sex, and race and Hispanic origin	2008		2007	
	Number	Rate	Number	Rate
Total black, both sexes				
All ages	289,306	716.7	289,585	723.4
Under 1 year ¹	8,513	1,190.7	8,944	1,250.0
1–4 years	1,131	42.1	1,137	42.2
5–14 years	1,287	20.1	1,363	20.9
15–24 years	7,222	105.6	7,380	109.3
25–34 years	9,287	159.8	9,662	168.9
35–44 years	15,797	281.1	16,990	299.5
45–54 years	35,994	665.4	36,818	694.2
55–64 years	48,748	1,385.6	47,682	1,414.7
65–74 years	51,810	2,734.4	51,384	2,791.4
75–84 years	59,172	5,652.7	58,986	5,725.7
85 years and over	50,315	12,071.2	49,194	12,281.5
Not stated	30	...	45	...
Age-adjusted rate ²	935.6	...	958.0
Black male				
All ages	147,275	763.4	148,309	775.6
Under 1 year ¹	4,728	1,293.2	4,975	1,363.2
1–4 years	654	47.8	621	45.3
5–14 years	769	23.6	816	24.6
15–24 years	5,522	159.4	5,751	168.1
25–34 years	6,378	225.4	6,648	240.3
35–44 years	9,201	348.0	10,106	378.9
45–54 years	20,703	828.2	21,495	876.7
55–64 years	28,686	1,827.7	28,127	1,870.8
65–74 years	28,001	3,541.4	27,672	3,604.9
75–84 years	27,094	7,107.5	26,815	7,169.0
85 years and over	15,518	12,538.0	15,255	12,964.7
Not stated	20	...	28	...
Age-adjusted rate ²	1,151.3	...	1,184.4
Black female				
All ages	142,031	674.0	141,276	675.7
Under 1 year ¹	3,785	1,083.5	3,969	1,132.2
1–4 years	477	36.1	516	39.0
5–14 years	518	16.4	547	17.0
15–24 years	1,701	50.4	1,629	48.9
25–34 years	2,909	97.6	3,014	102.1
35–44 years	6,596	221.7	6,884	229.1
45–54 years	15,290	525.5	15,323	537.2
55–64 years	20,062	1,029.5	19,555	1,047.4
65–74 years	23,809	2,156.4	23,712	2,209.5
75–84 years	32,077	4,819.3	32,171	4,902.9
85 years and over	34,797	11,874.0	33,939	11,997.4
Not stated	10	...	17	...
Age-adjusted rate ²	779.0	...	793.8

See footnotes at end of table.

Table 1. Deaths and death rates, by age, sex, and race and Hispanic origin, and age-adjusted death rates, by sex and race and Hispanic origin: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Age-specific rates are per 100,000 population in specified group. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals. Race and Hispanic origin are reported separately on the death certificate. Data for Hispanic origin and specified races other than white and black should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 34 states and the District of Columbia in 2008 and by 27 states and the District of Columbia in 2007; see "Technical Notes." The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes." Data for persons of Hispanic origin are included in the data for each race group, according to the decedent's reported race; see "Technical Notes"]

Age, sex, and race and Hispanic origin	2008		2007	
	Number	Rate	Number	Rate
Non-Hispanic black, both sexes				
All ages	285,959	746.4	286,366	750.7
Under 1 year ¹	8,193	1,271.8	8,629	1,309.1
1–4 years	1,085	44.6	1,102	43.7
5–14 years	1,260	20.9	1,335	21.6
15–24 years	7,096	109.1	7,237	112.6
25–34 years	9,130	166.2	9,521	176.2
35–44 years	15,600	291.9	16,797	310.9
45–54 years	35,605	683.5	36,465	713.2
55–64 years	48,289	1,418.6	47,199	1,446.2
65–74 years	51,303	2,791.6	50,890	2,847.8
75–84 years	58,551	5,753.3	58,398	5,822.9
85 years and over	49,823	12,259.9	48,762	12,468.1
Not stated	24	...	31	...
Age-adjusted rate ²	956.6	...	978.6
Non-Hispanic black male				
All ages	145,377	795.5	146,474	805.1
Under 1 year ¹	4,542	1,379.5	4,794	1,426.3
1–4 years	624	50.4	609	47.5
5–14 years	752	24.6	801	25.5
15–24 years	5,428	164.8	5,635	173.1
25–34 years	6,252	234.2	6,538	250.6
35–44 years	9,080	361.5	9,981	393.4
45–54 years	20,450	850.4	21,260	900.4
55–64 years	28,378	1,871.3	27,823	1,913.7
65–74 years	27,702	3,617.3	27,377	3,678.3
75–84 years	26,797	7,243.7	26,523	7,295.1
85 years and over	15,358	12,760.9	15,115	13,189.0
Not stated	14	...	18	...
Age-adjusted rate ²	1,178.3	...	1,210.9
Non-Hispanic black female				
All ages	140,582	701.6	139,892	701.0
Under 1 year ¹	3,651	1,159.1	3,835	1,187.1
1–4 years	461	38.6	493	39.8
5–14 years	508	17.1	534	17.6
15–24 years	1,669	52.0	1,602	50.5
25–34 years	2,878	101.9	2,983	106.7
35–44 years	6,520	230.2	6,816	237.9
45–54 years	15,155	540.4	15,205	552.5
55–64 years	19,911	1,054.8	19,376	1,070.7
65–74 years	23,600	2,201.6	23,513	2,255.0
75–84 years	31,753	4,902.0	31,875	4,985.7
85 years and over	34,465	12,049.1	33,647	12,169.3
Not stated	10	...	13	...
Age-adjusted rate ²	796.1	...	810.4

See footnotes at end of table.

Table 1. Deaths and death rates, by age, sex, and race and Hispanic origin, and age-adjusted death rates, by sex and race and Hispanic origin: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Age-specific rates are per 100,000 population in specified group. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals. Race and Hispanic origin are reported separately on the death certificate. Data for Hispanic origin and specified races other than white and black should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 34 states and the District of Columbia in 2008 and by 27 states and the District of Columbia in 2007; see "Technical Notes." The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes." Data for persons of Hispanic origin are included in the data for each race group, according to the decedent's reported race; see "Technical Notes"]

Age, sex, and race and Hispanic origin	2008		2007	
	Number	Rate	Number	Rate
Total AIAN, ³ both sexes ⁴				
All ages	14,785	432.1	14,367	444.0
Under 1 year ¹	405	582.0	432	921.7
1–4 years	100	38.9	97	54.9
5–14 years	97	17.5	94	18.2
15–24 years	636	105.5	622	103.3
25–34 years	774	151.0	731	146.6
35–44 years	1,214	266.9	1,213	264.8
45–54 years	2,090	477.0	1,958	456.4
55–64 years	2,446	847.0	2,367	858.3
65–74 years	2,668	1,844.7	2,536	1,853.8
75–84 years	2,589	3,646.9	2,546	3,760.3
85 years and over	1,762	6,155.5	1,771	6,708.1
Not stated	3	...	—	...
Age-adjusted rate ²	610.3	...	627.2
AIAN male ⁴				
All ages	8,170	478.0	7,885	488.2
Under 1 year ¹	234	659.7	241	1,009.9
1–4 years	50	38.4	57	63.6
5–14 years	52	18.5	61	23.2
15–24 years	462	150.7	441	143.7
25–34 years	530	199.0	514	198.3
35–44 years	725	315.0	768	332.5
45–54 years	1,318	618.3	1,193	573.0
55–64 years	1,440	1,039.3	1,372	1,037.0
65–74 years	1,466	2,172.8	1,357	2,131.7
75–84 years	1,271	4,156.7	1,214	4,193.4
85 years and over	621	6,503.3	667	7,638.6
Not stated	—	...	—	...
Age-adjusted rate ²	717.8	...	736.7
AIAN female ⁴				
All ages	6,615	386.3	6,482	400.0
Under 1 year ¹	170	498.3	191	830.3
1–4 years	50	39.5	40	46.0
5–14 years	45	16.5	33	13.0
15–24 years	174	58.7	181	61.3
25–34 years	244	99.1	217	90.6
35–44 years	488	217.2	445	196.0
45–54 years	772	343.2	765	346.4
55–64 years	1,006	669.6	995	693.5
65–74 years	1,202	1,557.9	1,179	1,611.9
75–84 years	1,319	3,263.7	1,332	3,436.8
85 years and over	1,141	5,981.3	1,104	6,248.2
Not stated	3	...	—	...
Age-adjusted rate ²	515.0	...	533.2

See footnotes at end of table.

Table 1. Deaths and death rates, by age, sex, and race and Hispanic origin, and age-adjusted death rates, by sex and race and Hispanic origin: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Age-specific rates are per 100,000 population in specified group. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals. Race and Hispanic origin are reported separately on the death certificate. Data for Hispanic origin and specified races other than white and black should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 34 states and the District of Columbia in 2008 and by 27 states and the District of Columbia in 2007; see "Technical Notes." The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes." Data for persons of Hispanic origin are included in the data for each race group, according to the decedent's reported race; see "Technical Notes"]

Age, sex, and race and Hispanic origin	2008		2007	
	Number	Rate	Number	Rate
Total API, ⁵ both sexes				
All ages	47,966	319.1	45,609	308.7
Under 1 year ¹	953	423.6	955	441.8
1–4 years	165	18.8	182	21.7
5–14 years	206	10.6	210	11.0
15–24 years	722	37.1	843	43.2
25–34 years	1,125	45.0	998	39.0
35–44 years	1,772	68.0	1,832	71.5
45–54 years	3,725	177.7	3,677	180.2
55–64 years	6,124	421.6	5,774	418.8
65–74 years	8,455	1,070.8	7,941	1,046.6
75–84 years	12,426	2,868.2	12,028	2,904.2
85 years and over	12,291	7,958.5	11,161	7,929.2
Not stated	3	...	8	...
Age-adjusted rate ²	414.2	...	415.0
API male				
All ages	24,744	338.1	23,823	331.4
Under 1 year ¹	536	464.2	537	483.5
1–4 years	79	17.5	109	25.3
5–14 years	123	12.4	118	12.2
15–24 years	500	50.1	611	61.0
25–34 years	679	55.0	632	50.1
35–44 years	1,114	87.7	1,110	88.9
45–54 years	2,220	223.0	2,212	229.1
55–64 years	3,558	530.4	3,334	523.1
65–74 years	4,724	1,313.4	4,487	1,304.7
75–84 years	6,136	3,500.0	5,950	3,538.4
85 years and over	5,073	8,742.2	4,717	8,918.0
Not stated	2	...	6	...
Age-adjusted rate ²	493.4	...	499.2
API female				
All ages	23,222	301.0	21,786	287.2
Under 1 year ¹	417	380.9	418	397.6
1–4 years	86	20.1	73	17.9
5–14 years	83	8.6	92	9.8
15–24 years	222	23.4	232	24.4
25–34 years	445	35.1	366	28.1
35–44 years	658	49.3	722	54.9
45–54 years	1,505	136.7	1,465	136.2
55–64 years	2,566	328.2	2,440	329.2
65–74 years	3,731	867.9	3,454	832.7
75–84 years	6,290	2,438.8	6,078	2,470.6
85 years and over	7,218	7,486.9	6,444	7,334.0
Not stated	1	...	2	...
Age-adjusted rate ²	353.5	...	350.6

See footnotes at end of table.

Table 1. Deaths and death rates, by age, sex, and race and Hispanic origin, and age-adjusted death rates, by sex and race and Hispanic origin: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Age-specific rates are per 100,000 population in specified group. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals. Race and Hispanic origin are reported separately on the death certificate. Data for Hispanic origin and specified races other than white and black should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported by 34 states and the District of Columbia in 2008 and by 27 states and the District of Columbia in 2007; see "Technical Notes." The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes." Data for persons of Hispanic origin are included in the data for each race group, according to the decedent's reported race; see "Technical Notes"]

Age, sex, and race and Hispanic origin	2008		2007	
	Number	Rate	Number	Rate
Hispanic, both sexes ⁶				
All ages	140,103	298.4	135,519	297.8
Under 1 year ¹	5,891	531.5	6,068	587.4
1–4 years	1,030	24.6	1,009	26.0
5–14 years	1,019	12.1	1,157	14.1
15–24 years	5,202	69.2	5,686	76.5
25–34 years	6,408	78.3	6,546	80.1
35–44 years	8,644	123.3	8,695	127.1
45–54 years	14,428	289.6	14,400	303.2
55–64 years	18,602	651.8	17,424	647.4
65–74 years	22,092	1,467.3	21,127	1,477.1
75–84 years	30,492	3,619.5	29,490	3,678.4
85 years and over	26,283	8,390.9	23,890	8,542.2
Not stated	11	...	27	...
Age-adjusted rate ²	536.0	...	546.1
Hispanic male ⁶				
All ages	77,261	318.5	75,708	321.8
Under 1 year ¹	3,279	578.5	3,343	632.7
1–4 years	588	27.5	556	28.0
5–14 years	548	12.7	663	15.8
15–24 years	4,076	103.8	4,506	115.3
25–34 years	4,848	107.4	4,956	110.1
35–44 years	5,934	159.1	6,037	166.3
45–54 years	9,513	374.3	9,634	399.2
55–64 years	11,518	835.5	10,764	831.4
65–74 years	12,506	1,839.4	11,978	1,862.7
75–84 years	14,938	4,288.9	14,438	4,364.8
85 years and over	9,502	8,475.4	8,809	8,953.7
Not stated	11	...	24	...
Age-adjusted rate ²	634.8	...	654.5
Hispanic female ⁶				
All ages	62,842	277.0	59,811	272.1
Under 1 year ¹	2,612	482.3	2,725	539.9
1–4 years	442	21.6	453	23.8
5–14 years	471	11.4	494	12.3
15–24 years	1,126	31.4	1,180	33.5
25–34 years	1,561	42.6	1,590	43.4
35–44 years	2,710	82.6	2,658	82.7
45–54 years	4,915	201.4	4,766	204.0
55–64 years	7,085	480.2	6,660	476.9
65–74 years	9,586	1,160.9	9,149	1,162.1
75–84 years	15,555	3,147.9	15,052	3,196.2
85 years and over	16,780	8,343.4	15,081	8,318.9
Not stated	—	...	3	...
Age-adjusted rate ²	449.3	...	452.7

... Category not applicable.

— Quantity zero.

¹Death rates for under 1 year (based on population estimates) differ from infant mortality rates (based on live births). See text for additional information on the infant mortality rate.

²For method of computation, see "Technical Notes."

³American Indian or Alaska Native.

⁴Includes deaths among Aleut and Eskimo persons.

⁵Asian or Pacific Islander.

⁶Includes all persons of Hispanic origin of any race; see "Technical Notes."

NOTE: Data are subject to sampling or random variation. For information regarding the calculation of standard errors and further discussion of variability of the data, see "Technical Notes."

Table 2. Deaths, death rates, and age-adjusted death rates for 113 selected causes, Injury by firearms, Drug-induced deaths, Alcohol-induced deaths, Injury at work, and Enterocolitis due to *Clostridium difficile*: United States, final 2007 and preliminary 2008

[Data are based on a continuous file of records received from the states. Rates per 100,000 population. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	2008			2007		
	Number	Rate	Age-adjusted rate	Number	Rate	Age-adjusted rate
All causes	2,472,699	813.2	758.6	2,423,712	803.6	760.2
Salmonella infections (A01–A02)	43	0.0	0.0	30	0.0	0.0
Shigellosis and amebiasis (A03,A06)	6	*	*	4	*	*
Certain other intestinal infections (A04,A07–A09)	7,883	2.6	2.4	6,758	2.2	2.1
Tuberculosis (A16–A19)	590	0.2	0.2	554	0.2	0.2
Respiratory tuberculosis (A16)	452	0.1	0.1	424	0.1	0.1
Other tuberculosis (A17–A19)	138	0.0	0.0	130	0.0	0.0
Whooping cough (A37)	20	0.0	0.0	9	*	*
Scarlet fever and erysipelas (A38,A46)	3	*	*	3	*	*
Meningococcal infection (A39)	102	0.0	0.0	87	0.0	0.0
Septicemia (A40–A41)	35,961	11.8	11.1	34,828	11.5	11.0
Syphilis (A50–A53)	34	0.0	0.0	42	0.0	0.0
Acute poliomyelitis (A80)	–	*	*	–	*	*
Arthropod-borne viral encephalitis (A83–A84,A85.2)	2	*	*	3	*	*
Measles (B05)	–	*	*	–	*	*
Viral hepatitis (B15–B19)	7,631	2.5	2.3	7,407	2.5	2.3
Human immunodeficiency virus (HIV) disease (B20–B24)	10,295	3.4	3.3	11,295	3.7	3.7
Malaria (B50–B54)	5	*	*	5	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20–A36, A42–A44,A48–A49,A54–A79,A81–A82,A85.0–A85.1,A85.8,A86–B04,B06–B09,B25–B49,B55–B99)	5,933	2.0	1.8	5,825	1.9	1.8
Malignant neoplasms (C00–C97)	566,137	186.2	175.5	562,875	186.6	178.4
Malignant neoplasms of lip, oral cavity and pharynx (C00–C14)	8,031	2.6	2.4	8,067	2.7	2.5
Malignant neoplasm of esophagus (C15)	13,739	4.5	4.2	13,592	4.5	4.3
Malignant neoplasm of stomach (C16)	11,381	3.7	3.5	11,388	3.8	3.6
Malignant neoplasms of colon, rectum and anus (C18–C21)	53,337	17.5	16.5	53,586	17.8	16.9
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	18,243	6.0	5.6	17,146	5.7	5.4
Malignant neoplasm of pancreas (C25)	35,267	11.6	10.9	34,117	11.3	10.8
Malignant neoplasm of larynx (C32)	3,759	1.2	1.2	3,634	1.2	1.1
Malignant neoplasms of trachea, bronchus and lung (C33–C34)	158,873	52.3	49.6	158,760	52.6	50.6
Malignant melanoma of skin (C43)	8,643	2.8	2.7	8,461	2.8	2.7
Malignant neoplasm of breast (C50)	41,049	13.5	12.6	40,970	13.6	12.9
Malignant neoplasm of cervix uteri (C53)	4,018	1.3	1.3	4,021	1.3	1.3
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54–C55)	7,682	2.5	2.4	7,456	2.5	2.3
Malignant neoplasm of ovary (C56)	14,373	4.7	4.4	14,621	4.8	4.6
Malignant neoplasm of prostate (C61)	28,517	9.4	8.8	29,093	9.6	9.2
Malignant neoplasms of kidney and renal pelvis (C64–C65)	12,915	4.2	4.0	12,703	4.2	4.0
Malignant neoplasm of bladder (C67)	14,053	4.6	4.3	13,843	4.6	4.4
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70–C72)	13,739	4.5	4.3	13,234	4.4	4.2
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81–C96)	54,998	18.1	17.2	54,991	18.2	17.6
Hodgkin's disease (C81)	1,170	0.4	0.4	1,271	0.4	0.4
Non-Hodgkin's lymphoma (C82–C85)	20,374	6.7	6.3	20,528	6.8	6.5
Leukemia (C91–C95)	22,357	7.4	7.0	21,825	7.2	7.0
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	11,038	3.6	3.4	11,307	3.7	3.6

See footnotes at end of table.

Table 2. Deaths, death rates, and age-adjusted death rates for 113 selected causes, Injury by firearms, Drug-induced deaths, Alcohol-induced deaths, Injury at work, and Enterocolitis due to *Clostridium difficile*: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Rates per 100,000 population. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	2008			2007		
	Number	Rate	Age-adjusted rate	Number	Rate	Age-adjusted rate
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue. . . . (C96)	59	0.0	0.0	60	0.0	0.0
All other and unspecified malignant neoplasms. . . . (C17,C23–C24,C26–C31,C37–C41,C44–C49, C51–C52,C57–C60,C62–C63,C66,C68–C69,C73–C80,C97)	63,519	20.9	19.7	63,192	21.0	20.0
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior . . . (D00–D48)	14,481	4.8	4.5	14,204	4.7	4.5
Anemias (D50–D64)	5,033	1.7	1.5	4,829	1.6	1.5
Diabetes mellitus (E10–E14)	70,601	23.2	21.8	71,382	23.7	22.5
Nutritional deficiencies (E40–E64)	2,981	1.0	0.9	2,852	0.9	0.9
Malnutrition (E40–E46)	2,767	0.9	0.8	2,644	0.9	0.8
Other nutritional deficiencies (E50–E64)	214	0.1	0.1	208	0.1	0.1
Meningitis. . . . (G00,G03)	633	0.2	0.2	655	0.2	0.2
Parkinson's disease. . . . (G20–G21)	20,507	6.7	6.4	20,058	6.7	6.4
Alzheimer's disease. . . . (G30)	82,476	27.1	24.4	74,632	24.7	22.7
Major cardiovascular diseases. . . . (I00–I78)	804,899	264.7	243.6	806,156	267.3	249.9
Diseases of heart. . . . (I00–I09,I11,I13,I20–I51)	617,527	203.1	186.7	616,067	204.3	190.9
Acute rheumatic fever and chronic rheumatic heart diseases. . . . (I00–I09)	3,149	1.0	1.0	3,201	1.1	1.0
Hypertensive heart disease (I11)	32,374	10.6	9.7	30,780	10.2	9.5
Hypertensive heart and renal disease (I13)	2,867	0.9	0.9	2,987	1.0	0.9
Ischemic heart diseases (I20–I25)	405,019	133.2	122.6	406,351	134.7	126.0
Acute myocardial infarction (I21–I22)	133,723	44.0	40.7	132,968	44.1	41.4
Other acute ischemic heart diseases (I24)	4,233	1.4	1.3	4,092	1.4	1.3
Other forms of chronic ischemic heart disease. . . . (I20,I25)	267,063	87.8	80.6	269,291	89.3	83.3
Atherosclerotic cardiovascular disease, so described. . . . (I25.0)	58,517	19.2	17.6	59,051	19.6	18.2
All other forms of chronic ischemic heart disease. . . . (I20,I25.1–I25.9)	208,545	68.6	63.0	210,240	69.7	65.1
Other heart diseases (I26–I51)	174,118	57.3	52.6	172,748	57.3	53.4
Acute and subacute endocarditis. . . . (I33)	1,179	0.4	0.4	1,225	0.4	0.4
Diseases of pericardium and acute myocarditis (I30–I31,I40)	829	0.3	0.3	867	0.3	0.3
Heart failure (I50)	57,215	18.8	17.0	56,565	18.8	17.3
All other forms of heart disease (I26–I28,I34–I38,I42–I49,I51)	114,895	37.8	34.9	114,091	37.8	35.5
Essential hypertension and hypertensive renal disease. . . . (I10,I12,I15)	25,823	8.5	7.7	23,965	7.9	7.4
Cerebrovascular diseases. . . . (I60–I69)	133,750	44.0	40.6	135,952	45.1	42.2
Atherosclerosis (I70)	7,846	2.6	2.3	8,232	2.7	2.5
Other diseases of circulatory system (I71–I78)	19,952	6.6	6.1	21,940	7.3	6.9
Aortic aneurysm and dissection (I71)	11,088	3.6	3.4	12,986	4.3	4.1
Other diseases of arteries, arterioles and capillaries (I72–I78)	8,864	2.9	2.7	8,954	3.0	2.8
Other disorders of circulatory system (I80–I99)	4,034	1.3	1.2	4,101	1.4	1.3
Influenza and pneumonia (J09–J18) ¹	56,335	18.5	17.0	52,717	17.5	16.2
Influenza (J09–J11) ¹	1,721	0.6	0.5	411	0.1	0.1
Pneumonia (J12–J18)	54,614	18.0	16.4	52,306	17.3	16.1
Other acute lower respiratory infections (J20–J22,U04) ²	285	0.1	0.1	255	0.1	0.1
Acute bronchitis and bronchiolitis (J20–J21)	235	0.1	0.1	213	0.1	0.1
Other and unspecified acute lower respiratory infections. . . . (J22,U04) ^{2,3}	50	0.0	0.0	42	0.0	0.0
Chronic lower respiratory diseases (J40–J47)	141,075	46.4	44.0	127,924	42.4	40.8

See footnotes at end of table.

Table 2. Deaths, death rates, and age-adjusted death rates for 113 selected causes, Injury by firearms, Drug-induced deaths, Alcohol-induced deaths, Injury at work, and Enterocolitis due to *Clostridium difficile*: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Rates per 100,000 population. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	2008			2007		
	Number	Rate	Age-adjusted rate	Number	Rate	Age-adjusted rate
Bronchitis, chronic and unspecified (J40–J42)	733	0.2	0.2	667	0.2	0.2
Emphysema (J43)	12,459	4.1	3.9	12,790	4.2	4.1
Asthma (J45–J46)	3,395	1.1	1.0	3,447	1.1	1.1
Other chronic lower respiratory diseases (J44, J47)	124,489	40.9	38.8	111,020	36.8	35.3
Pneumoconioses and chemical effects (J60–J66, J68)	905	0.3	0.3	915	0.3	0.3
Pneumonitis due to solids and liquids (J69)	16,622	5.5	5.0	16,988	5.6	5.2
Other diseases of respiratory system (J00–J06, J30–J39, J67, J70–J98)	30,000	9.9	9.3	28,508	9.5	9.0
Peptic ulcer (K25–K28)	3,070	1.0	0.9	3,045	1.0	1.0
Diseases of appendix (K35–K38)	420	0.1	0.1	426	0.1	0.1
Hernia (K40–K46)	1,682	0.6	0.5	1,698	0.6	0.5
Chronic liver disease and cirrhosis (K70, K73–K74)	29,963	9.9	9.2	29,165	9.7	9.1
Alcoholic liver disease (K70)	14,859	4.9	4.5	14,406	4.8	4.5
Other chronic liver disease and cirrhosis (K73–K74)	15,104	5.0	4.6	14,759	4.9	4.6
Cholelithiasis and other disorders of gallbladder (K80–K82)	3,425	1.1	1.0	3,237	1.1	1.0
Nephritis, nephrotic syndrome and nephrosis (N00–N07, N17–N19, N25–N27)	48,283	15.9	14.8	46,448	15.4	14.5
Acute and rapidly progressive nephritic and nephrotic syndrome (N00–N01, N04)	165	0.1	0.0	206	0.1	0.1
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02–N03, N05–N07, N26)	4,120	1.4	1.2	2,958	1.0	0.9
Renal failure (N17–N19)	43,965	14.5	13.5	43,263	14.3	13.6
Other disorders of kidney (N25, N27)	33	0.0	0.0	21	0.0	0.0
Infections of kidney (N10–N12, N13.6, N15.1)	629	0.2	0.2	628	0.2	0.2
Hyperplasia of prostate (N40)	504	0.2	0.1	491	0.2	0.1
Inflammatory diseases of female pelvic organs (N70–N76)	133	0.0	0.0	116	0.0	0.0
Pregnancy, childbirth and the puerperium (O00–O99)	774	0.3	0.3	769	0.3	0.3
Pregnancy with abortive outcome (O00–O07)	34	0.0	0.0	31	0.0	0.0
Other complications of pregnancy, childbirth and the puerperium (O10–O99)	739	0.2	0.2	738	0.2	0.3
Certain conditions originating in the perinatal period (P00–P96)	13,889	4.6	4.4	14,599	4.8	4.7
Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	10,284	3.4	3.3	10,421	3.5	3.4
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99)	38,455	12.6	11.7	33,500	11.1	10.4
All other diseases (residual)	252,706	83.1	76.4	238,192	79.0	73.8
Accidents (unintentional injuries) (V01–X59, Y85–Y86)	121,207	39.9	38.6	123,706	41.0	40.0
Transport accidents (V01–V99, Y85)	42,742	14.1	13.9	46,844	15.5	15.3
Motor vehicle accidents (V02–V04, V09.0, V09.2, V12–V14, V19.0–V19.2, V19.4–V19.6, V20–V79, V80.3–V80.5, V81.0–V81.1, V82.0–V82.1, V83–V86, V87.0–V87.8, V88.0–V88.8, V89.0, V89.2)	39,831	13.1	12.9	43,945	14.6	14.4
Other land transport accidents (V01, V05–V06, V09.1, V09.3–V09.9, V10–V11, V15–V18, V19.3, V19.8–V19.9, V80.0–V80.2, V80.6–V80.9, V81.2–V81.9, V82.2–V82.9, V87.9, V88.9, V89.1, V89.3, V89.9)	1,146	0.4	0.4	1,083	0.4	0.3
Water, air and space, and other and unspecified transport accidents and their sequelae (V90–V99, Y85)	1,765	0.6	0.6	1,816	0.6	0.6
Nontransport accidents (W00–X59, Y86)	78,465	25.8	24.7	76,862	25.5	24.6
Falls (W00–W19)	24,062	7.9	7.3	22,631	7.5	7.0
Accidental discharge of firearms (W32–W34)	587	0.2	0.2	613	0.2	0.2

See footnotes at end of table.

Table 2. Deaths, death rates, and age-adjusted death rates for 113 selected causes, Injury by firearms, Drug-induced deaths, Alcohol-induced deaths, Injury at work, and Enterocolitis due to *Clostridium difficile*: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Rates per 100,000 population. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	2008			2007		
	Number	Rate	Age-adjusted rate	Number	Rate	Age-adjusted rate
Accidental drowning and submersion (W65–W74)	3,549	1.2	1.2	3,443	1.1	1.1
Accidental exposure to smoke, fire and flames (X00–X09)	2,907	1.0	0.9	3,286	1.1	1.1
Accidental poisoning and exposure to noxious substances (X40–X49)	30,306	10.0	9.9	29,846	9.9	9.8
Other and unspecified nontransport accidents and their sequelae (W20–W31, W35–W64, W75–W99, X10–X39, X50–X59, Y86)	17,054	5.6	5.3	17,043	5.7	5.4
Intentional self-harm (suicide) (*U03, X60–X84, Y87.0)	35,933	11.8	11.6	34,598	11.5	11.3
Intentional self-harm (suicide) by discharge of firearms (X72–X74)	18,251	6.0	5.8	17,352	5.8	5.6
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03, X60–X71, X75–X84, Y87.0)	17,681	5.8	5.7	17,246	5.7	5.6
Assault (homicide) (*U01–*U02, X85–Y09, Y87.1)	17,837	5.9	5.9	18,361	6.1	6.1
Assault (homicide) by discharge of firearms (*U01.4, X93–X95)	12,209	4.0	4.0	12,632	4.2	4.2
Assault (homicide) by other and unspecified means and their sequelae (*U01.0–*U01.3, *U01.5–*U01.9, *U02, X85–X92, X96–Y09, Y87.1)	5,628	1.9	1.8	5,729	1.9	1.9
Legal intervention (Y35, Y89.0)	380	0.1	0.1	412	0.1	0.1
Events of undetermined intent (Y10–Y34, Y87.2, Y89.9)	4,979	1.6	1.6	5,381	1.8	1.8
Discharge of firearms, undetermined intent (Y22–Y24)	276	0.1	0.1	276	0.1	0.1
Other and unspecified events of undetermined intent and their sequelae (Y10–Y21, Y25–Y34, Y87.2, Y89.9)	4,703	1.5	1.5	5,105	1.7	1.7
Operations of war and their sequelae (Y36, Y89.1)	31	0.0	0.0	21	0.0	0.0
Complications of medical and surgical care (Y40–Y84, Y88)	2,602	0.9	0.8	2,597	0.9	0.8
Injury by firearms (*U01.4, W32–W34, X72–X74, X93–X95, Y22–Y24, Y35.0)4	31,651	10.4	10.3	31,224	10.4	10.2
Drug-induced deaths (D52.1, D59.0, D59.2, D61.1, D64.2, E06.4, E16.0, E23.1, E24.2, E27.3, E66.1, F11.0–F11.5, F11.7–F11.9, F12.0–F12.5, F12.7–F12.9, F13.0–F13.5, F13.7–F13.9, F14.0–F14.5, F14.7–F14.9, F15.0–F15.5, F15.7–F15.9, F16.0–F16.5, F16.7–F16.9, F17.0, F17.3–F17.5, F17.7–F17.9, F18.0–F18.5, F18.7–18.9, F19.0–F19.5, F19.7–F19.9, G21.1, G24.0, G25.1, G25.4, G25.6, G44.4, G62.0, G72.0, I95.2, J70.2–J70.4, K85.3, L10.5, L27.0–L27.1, M10.2, M32.0, M80.4, M81.4, M83.5, M87.1, R50.2, R78.1–R78.5, X40–X44, X60–X64, X85, Y10–14)4	37,777	12.4	12.3	38,371	12.7	12.6
Alcohol-induced deaths (E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K85.2, K86.0, R78.0, X45, X65, Y15)4	24,081	7.9	7.4	23,199	7.7	7.3
Injury at work5	4,689	1.9	1.9	5,025	2.1	2.1
Enterocolitis due to <i>Clostridium difficile</i> (A047)6	7,483	2.5	2.3	6,372	2.1	2.0

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

– Quantity zero.

1 New ICD–10 code J09 (Influenza due to identified avian influenza virus) was added to the category in 2007.

2 New ICD–10 code U04 (Severe acute respiratory syndrome [SARS], unspecified) was added to the category in 2007.

3 Cause-of-death title has been changed in 2007 to reflect the addition of SARS (U04).

4 Included in selected categories above.

5 Described in "Technical Notes."

6 Included in "Certain other intestinal infections (A04, A07–A09)" shown above; see "Technical Notes."

NOTES: For certain causes of death such as unintentional injuries, homicides, suicides, and respiratory diseases, preliminary and final data differ because of the truncated nature of the preliminary file. Data are subject to sampling or random variation. For information regarding the calculation of standard errors and further discussion of variability of the data, see "Technical Notes."

Table 3. Deaths, death rates, and age-adjusted death rates: United States, and each state and territory, final 2007 and preliminary 2008

[By place of residence. Data are based on a continuous file of records received from the states. Rates are per 100,000 population. Age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals]

Area	2008			2007		
	Number	Rate	Age-adjusted rate	Number	Rate	Age-adjusted rate
United States ¹	2,473,018	813.3	758.7	2,423,712	803.6	760.2
Alabama	47,712	1,023.4	930.3	46,696	1,009.0	930.7
Alaska	3,483	507.5	739.6	3,463	506.7	755.1
Arizona	45,610	701.7	650.6	45,554	718.7	682.1
Arkansas	29,310	1,026.5	899.2	28,191	994.5	882.8
California	234,229	637.2	658.8	233,720	639.4	674.2
Colorado	31,256	632.8	708.6	29,993	616.9	700.8
Connecticut	28,797	822.5	691.4	28,651	818.1	694.1
Delaware	7,623	873.1	780.8	7,327	847.3	773.6
District of Columbia	5,139	868.3	849.9	5,188	881.9	866.9
Florida	170,668	931.2	679.0	168,096	921.0	685.9
Georgia	69,942	722.1	835.4	68,331	715.9	839.8
Hawaii	9,475	735.5	589.0	9,495	739.8	607.4
Idaho	10,942	718.1	721.7	10,822	721.8	734.6
Illinois	103,615	803.1	772.0	100,503	782.0	759.8
Indiana	56,743	889.8	835.1	54,000	851.0	809.9
Iowa	28,533	950.3	744.0	27,221	911.0	718.6
Kansas	24,969	891.1	784.7	24,491	882.2	783.0
Kentucky	41,280	966.9	901.2	40,090	945.2	896.9
Louisiana	41,217	934.5	922.0	39,966	930.9	926.4
Maine	12,531	951.9	764.1	12,493	948.4	773.6
Maryland	43,885	779.0	771.6	43,757	778.8	782.7
Massachusetts	53,521	823.7	705.9	52,917	820.4	707.5
Michigan	88,418	883.9	811.7	86,721	861.0	806.1
Minnesota	38,487	737.2	675.2	37,138	714.5	661.5
Mississippi	28,980	986.2	950.0	28,255	968.0	943.0
Missouri	56,566	956.9	847.0	54,166	921.4	826.7
Montana	8,903	920.3	785.9	8,624	900.3	772.7
Nebraska	15,455	866.6	741.1	15,263	860.1	743.7
Nevada ²	20,790	799.6	868.2	18,687	728.4	803.5
New Hampshire	10,268	780.4	712.5	10,303	783.0	727.0
New Jersey	69,993	806.1	716.8	69,662	802.0	724.2
New Mexico	15,996	806.1	758.2	15,482	785.9	755.9
New York	148,660	762.7	675.8	147,680	765.3	686.4
North Carolina	77,277	837.9	825.6	76,046	839.3	834.4
North Dakota	5,870	915.1	713.0	5,561	869.3	679.5
Ohio	109,749	955.5	844.0	106,534	929.1	830.8
Oklahoma	37,061	1,017.5	932.2	36,032	996.1	920.4
Oregon	31,939	842.7	747.9	31,403	838.0	753.9
Pennsylvania	127,450	1,023.8	796.5	125,104	1,006.2	790.1
Rhode Island	9,740	926.9	749.6	9,723	919.1	750.0
South Carolina	40,305	899.7	839.7	39,439	894.8	849.7
South Dakota	7,080	880.4	708.4	6,826	857.3	693.5
Tennessee	58,882	947.4	889.7	57,087	927.2	885.2
Texas	165,197	679.1	777.3	160,548	671.6	777.7
Utah	13,991	511.3	656.9	14,143	534.6	694.2
Vermont	5,213	839.1	722.2	5,179	833.6	729.3
Virginia	59,093	760.6	762.6	58,225	755.0	770.6
Washington	48,603	742.1	723.3	47,323	731.6	722.2
West Virginia	21,549	1,187.6	958.1	21,086	1,163.7	951.7
Wisconsin	46,799	831.5	729.7	46,241	825.5	732.3
Wyoming	4,222	792.6	772.5	4,266	815.9	802.0
Puerto Rico	28,781	727.9	700.4	29,169	739.9	726.7
Virgin Islands	699	636.4	682.9	703	640.1	700.2
Guam	762	433.0	683.1	778	448.5	694.9
American Samoa	241	371.8	958.9	250	390.5	1,054.4
Northern Marianas	---	---	---	137	162.0	913.0

--- Data not available.

¹Excludes data for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas.

²Data are based on the state of occurrence due to the lack of geographic code for the state; see "Technical Notes."

NOTE: Data are subject to sampling or random variation. For information regarding calculation of standard errors and further discussion of variability of the data, see "Technical Notes."

Table 4. Infant deaths and infant mortality rates, by age and race and Hispanic origin: United States, final 2007 and preliminary 2008

[Data are based on the continuous file of records received from the states. Rates per 1,000 live births. Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals. Race and Hispanic origin are reported separately on both the birth and death certificate. Rates for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported for deaths by 34 states and the District of Columbia in 2008 and 27 states and the District of Columbia in 2007, and were reported for births by 30 states in 2008 and 25 states in 2007; see "Technical Notes." Multiple-race data for these states were bridged to single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes"]

Age and race and Hispanic origin	2008		2007	
	Number	Rate	Number	Rate
All races ¹				
Under 1 year	28,033	6.59	29,138	6.75
Under 28 days	18,154	4.27	19,058	4.42
28 days–11 months	9,879	2.32	10,080	2.34
Total white				
Under 1 year	18,162	5.54	18,807	5.64
Under 28 days	11,820	3.61	12,333	3.70
28 days–11 months	6,343	1.94	6,474	1.94
Non-Hispanic white				
Under 1 year	12,545	5.52	12,998	5.63
Under 28 days	8,022	3.53	8,417	3.64
28 days–11 months	4,522	1.99	4,581	1.98
Total black				
Under 1 year	8,513	12.68	8,944	13.24
Under 28 days	5,483	8.17	5,842	8.65
28 days–11 months	3,030	4.51	3,102	4.59
Hispanic ²				
Under 1 year	5,891	5.67	6,068	5.71
Under 28 days	3,959	3.81	4,060	3.82
28 days–11 months	1,932	1.86	2,008	1.89

¹Includes races other than white or black.

²Includes all persons of Hispanic origin of any race; see "Technical Notes."

NOTES: Data are subject to sampling or random variation. For information regarding the calculation of standard errors and further discussion of variability of the data, see "Technical Notes." Although the infant mortality rate is the preferred indicator of the risk of dying during the first year of life, another measure of infant mortality, the infant death rate, is shown elsewhere in this report. The two measures typically are similar but use different denominators. For more information on these measures of risk, see "Infant mortality" section in "Technical Notes."

Table 5. Infant deaths and infant mortality rates for 130 selected causes: United States, final 2007 and preliminary 2008

[Data are based on a continuous file of records from the states. Rates per 100,000 live births. For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	2008		2007	
	Number	Rate	Number	Rate
All causes	28,029	659.3	29,138	675.1
Certain infectious and parasitic diseases (A00–B99)	485	11.4	482	11.2
Certain intestinal infectious diseases (A00–A08)	12	*	13	*
Diarrhea and gastroenteritis of infectious origin (A09)	-	*	-	*
Tuberculosis (A16–A19)	-	*	2	*
Tetanus (A33,A35)	-	*	-	*
Diphtheria (A36)	-	*	-	*
Whooping cough (A37)	18	*	8	*
Meningococcal infection (A39)	9	*	11	*
Septicemia (A40–A41)	293	6.9	283	6.6
Congenital syphilis (A50)	-	*	5	*
Gonococcal infection (A54)	-	*	-	*
Viral diseases (A80–B34)	102	2.4	115	2.7
Acute poliomyelitis (A80)	-	*	-	*
Varicella (chickenpox) (B01)	-	*	-	*
Measles (B05)	-	*	-	*
Human immunodeficiency virus (HIV) disease (B20–B24)	-	*	5	*
Mumps (B26)	-	*	-	*
Other and unspecified viral diseases (A81–B00,B02–B04,B06–B19,B25,B27–B34)	102	2.4	110	2.5
Candidiasis (B37)	7	*	13	*
Malaria (B50–B54)	-	*	-	*
Pneumocystosis (B59)	3	*	1	*
All other and unspecified infectious and parasitic diseases (A20–A32,A38,A42–A49, A51–A53,A55–A79,B35–B36,B38–B49,B55–B58,B60–B99)	40	0.9	31	0.7
Neoplasms (C00–D48)	128	3.0	131	3.0
Malignant neoplasms (C00–C97)	68	1.6	72	1.7
Hodgkin's disease and non-Hodgkin's lymphomas (C81–C85)	1	*	2	*
Leukemia (C91–C95)	27	0.6	21	0.5
Other and unspecified malignant neoplasms (C00–C80,C88,C90,C96–C97)	39	0.9	49	1.1
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	60	1.4	59	1.4
Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50–D89)	81	1.9	116	2.7
Anemias (D50–D64)	15	*	17	*
Hemorrhagic conditions and other diseases of blood and blood-forming organs (D65–D76)	56	1.3	77	1.8
Certain disorders involving the immune mechanism (D80–D89)	9	*	22	0.5
Endocrine, nutritional and metabolic diseases (E00–E88)	247	5.8	252	5.8
Short stature, not elsewhere classified (E34.3)	9	*	5	*
Nutritional deficiencies (E40–E64)	9	*	7	*
Cystic fibrosis (E84)	4	*	11	*
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86–E87)	80	1.9	60	1.4
All other endocrine, nutritional and metabolic diseases (E00–E32,E34.0–E34.2, E34.4–E34.9,E65–E83,E85,E88)	145	3.4	169	3.9
Diseases of the nervous system (G00–G98)	414	9.7	413	9.6
Meningitis (G00,G03)	67	1.6	82	1.9
Infantile spinal muscular atrophy, type I (Werdnig-Hoffman) (G12.0)	5	*	13	*

See footnotes at end of table.

Table 5. Infant deaths and infant mortality rates for 130 selected causes: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records from the states. Rates per 100,000 live births. For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	2008		2007	
	Number	Rate	Number	Rate
Infantile cerebral palsy (G80)	8	*	11	*
Anoxic brain damage, not elsewhere classified (G93.1)	51	1.2	64	1.5
Other diseases of nervous system (G04,G06–G11,G12.1G12.9,G20–G72,G81–G92,G93.0, G93.2–G93.9,G95–G98)	283	6.7	243	5.6
Diseases of the ear and mastoid process (H60–H93)	6	*	3	*
Diseases of the circulatory system (I00–I99)	590	13.9	624	14.5
Pulmonary heart disease and diseases of pulmonary circulation (I26–I28)	88	2.1	100	2.3
Pericarditis, endocarditis and myocarditis (I30,I33,I40)	19	*	21	0.5
Cardiomyopathy (I42)	114	2.7	120	2.8
Cardiac arrest (I46)	25	0.6	29	0.7
Cerebrovascular diseases (I60–I69)	144	3.4	132	3.1
All other diseases of circulatory system (I00–I25,I31,I34–I38,I44–I45,I47–I51,I70–I99)	201	4.7	222	5.1
Diseases of the respiratory system (J00–J98,U04) ¹	578	13.6	640	14.8
Acute upper respiratory infections (J00–J06)	12	*	14	*
Influenza and pneumonia (J09–J18) ²	225	5.3	222	5.1
Influenza (J09–J11) ²	16	*	13	*
Pneumonia (J12–J18)	208	4.9	209	4.8
Acute bronchitis and acute bronchiolitis (J20–J21)	43	1.0	45	1.0
Bronchitis, chronic and unspecified (J40–J42)	22	0.5	24	0.6
Asthma (J45–J46)	6	*	4	*
Pneumonitis due to solids and liquids (J69)	10	*	10	*
Other and unspecified diseases of respiratory system (J22,J30–J39,J43–J44,J47–J68,J70–J98,U04) ¹	260	6.1	321	7.4
Diseases of the digestive system (K00–K92)	578	13.6	677	15.7
Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50–K55)	355	8.4	413	9.6
Hernia of abdominal cavity and intestinal obstruction without hernia (K40–K46,K56)	47	1.1	68	1.6
All other and unspecified diseases of digestive system (K00–K28,K30–K38,K57–K92)	175	4.1	196	4.5
Diseases of the genitourinary system (N00–N95)	172	4.0	169	3.9
Renal failure and other disorders of kidney (N17–N19,N25,N27)	140	3.3	138	3.2
Other and unspecified diseases of genitourinary system (N00–N15,N20–N23,N26,N28–N95)	32	0.8	31	0.7
Certain conditions originating in the perinatal period (P00–P96)	13,738	323.2	14,466	335.2
Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00–P04)	3,153	74.2	3,274	75.9
Newborn affected by maternal hypertensive disorders (P00.0)	84	2.0	89	2.1
Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1–P00.9)	89	2.1	91	2.1
Newborn affected by maternal complications of pregnancy (P01)	1,764	41.5	1,769	41.0
Newborn affected by incompetent cervix (P01.0)	447	10.5	488	11.3
Newborn affected by premature rupture of membranes (P01.1)	840	19.8	851	19.7
Newborn affected by multiple pregnancy (P01.5)	257	6.0	238	5.5
Newborn affected by other maternal complications of pregnancy (P01.2–P01.4,P01.6–P01.9)	220	5.2	192	4.4
Newborn affected by complications of placenta, cord and membranes (P02)	1,073	25.2	1,135	26.3
Newborn affected by complications involving placenta (P02.0–P02.3)	531	12.5	579	13.4
Newborn affected by complications involving cord (P02.4–P02.6)	55	1.3	43	1.0
Newborn affected by chorioamnionitis (P02.7)	486	11.4	511	11.8
Newborn affected by other and unspecified abnormalities of membranes (P02.8–P02.9)	1	*	2	*
Newborn affected by other complications of labor and delivery (P03)	95	2.2	127	2.9

See footnotes at end of table.

Table 5. Infant deaths and infant mortality rates for 130 selected causes: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records from the states. Rates per 100,000 live births. For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	2008		2007	
	Number	Rate	Number	Rate
Newborn affected by noxious influences transmitted via placenta or breast milk(P04)	48	1.1	63	1.5
Disorders related to length of gestation and fetal malnutrition (P05–P08)	4,816	113.3	4,961	114.9
Slow fetal growth and fetal malnutrition(P05)	83	2.0	104	2.4
Disorders related to short gestation and low birth weight, not elsewhere classified(P07)	4,733	111.3	4,857	112.5
Extremely low birth weight or extreme immaturity(P07.0,P07.2)	3,636	85.5	3,706	85.9
Other low birth weight or preterm(P07.1,P07.3)	1,097	25.8	1,151	26.7
Disorders related to long gestation and high birth weight(P08)	-	*	-	*
Birth trauma (P10–P15)	18	*	12	*
Intrauterine hypoxia and birth asphyxia (P20–P21)	382	9.0	356	8.2
Intrauterine hypoxia(P20)	144	3.4	106	2.5
Birth asphyxia(P21)	238	5.6	250	5.8
Respiratory distress of newborn(P22)	625	14.7	789	18.3
Other respiratory conditions originating in the perinatal period (P23–P28)	1,102	25.9	1,117	25.9
Congenital pneumonia(P23)	74	1.7	103	2.4
Neonatal aspiration syndromes(P24)	58	1.4	51	1.2
Interstitial emphysema and related conditions originating in the perinatal period(P25)	121	2.8	124	2.9
Pulmonary hemorrhage originating in the perinatal period(P26)	199	4.7	161	3.7
Chronic respiratory disease originating in the perinatal period(P27)	239	5.6	243	5.6
Atelectasis(P28.0–P28.1)	334	7.9	366	8.5
All other respiratory conditions originating in the perinatal period (P28.2–P28.9)	77	1.8	69	1.6
Infections specific to the perinatal period (P35–P39)	896	21.1	1,057	24.5
Bacterial sepsis of newborn(P36)	696	16.4	820	19.0
Omphalitis of newborn with or without mild hemorrhage(P38)	2	*	4	*
All other infections specific to the perinatal period (P35,P37,P39)	198	4.7	233	5.4
Hemorrhagic and hematological disorders of newborn (P50–P61)	642	15.1	711	16.5
Neonatal hemorrhage(P50–P52,P54)	551	13.0	597	13.8
Hemorrhagic disease of newborn(P53)	2	*	-	*
Hemolytic disease of newborn due to isoimmunization and other perinatal jaundice (P55–P59)	10	*	15	*
Hematological disorders (P60–P61)	78	1.8	99	2.3
Syndrome of infant of a diabetic mother and neonatal diabetes mellitus (P70.0–P70.2)	11	*	14	*
Necrotizing enterocolitis of newborn(P77)	547	12.9	554	12.8
Hydrops fetalis not due to hemolytic disease (P83.2)	170	4.0	177	4.1
Other perinatal conditions (P29,P70.3–P76,P78–P81,P83.0–P83.1,P83.3–P96)	1,375	32.3	1,444	33.5
Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	5,647	132.8	5,785	134.0
Anencephaly and similar malformations (Q00)	340	8.0	321	7.4
Congenital hydrocephalus (Q03)	104	2.4	93	2.2
Spina bifida (Q05)	22	0.5	19	*
Other congenital malformations of nervous system (Q01–Q02,Q04,Q06–Q07)	356	8.4	393	9.1
Congenital malformations of heart (Q20–Q24)	1,307	30.7	1,363	31.6
Other congenital malformations of circulatory system (Q25–Q28)	219	5.2	254	5.9
Congenital malformations of respiratory system (Q30–Q34)	378	8.9	410	9.5
Congenital malformations of digestive system (Q35–Q45)	85	2.0	132	3.1
Congenital malformations of genitourinary system (Q50–Q64)	516	12.1	514	11.9
Congenital malformations and deformations of musculoskeletal system, limbs and integument (Q65–Q85)	665	15.6	623	14.4
Down's syndrome (Q90)	87	2.0	82	1.9

See footnotes at end of table.

Table 5. Infant deaths and infant mortality rates for 130 selected causes: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records from the states. Rates per 100,000 live births. For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures for 2008 are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	2008		2007	
	Number	Rate	Number	Rate
Edward's syndrome (Q91.0–Q91.3)	556	13.1	525	12.2
Patau's syndrome (Q91.4–Q91.7)	278	6.5	295	6.8
Other congenital malformations and deformations. (Q10–Q18,Q86–Q89)	535	12.6	552	12.8
Other chromosomal abnormalities, not elsewhere classified (Q92–Q99)	200	4.7	209	4.8
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99)	3,582	84.3	3,617	83.8
Sudden infant death syndrome (R95)	2,292	53.9	2,453	56.8
Other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R53,R55–594,R96–R99)	1,290	30.3	1,164	27.0
All other diseases (residual)	34	0.8	16	*
External causes of mortality (*U01,V01–Y84)	1,750	41.2	1,747	40.5
Accidents (unintentional injuries) (V01–X59)	1,299	30.6	1,285	29.8
Transport accidents (V01–V99)	105	2.5	127	2.9
Motor vehicle accidents (V02–V04,V09.0,V09.2,V12–V14,V19.0–V19.2,V19.4–V19.6,V20–V79,V80.3–V80.5,V81.0–V81.1,V82.0–V82.1,V83–V86,V87.0–V87.8,V88.0–V88.8,V89.0,V89.2)	104	2.4	124	2.9
Other and unspecified transport accidents (V01,V05–V06,V09.1,V09.3–V09.9,V10–V11,V15–V18,V19.3,V19.8–V19.9,V80.0–V80.2,V80.6–V80.9,V81.2–V81.9,V82.2–V82.9,V87.9,V88.9,V89.1,V89.3,V89.9,V90–V99)	1	*	3	*
Falls (W00–W19)	19	*	24	0.6
Accidental discharge of firearms. (W32–W34)	-	*	1	*
Accidental drowning and submersion. (W65–W74)	37	0.9	57	1.3
Accidental suffocation and strangulation in bed. (W75)	730	17.2	669	15.5
Other accidental suffocation and strangulation (W76–W77,W81–W84)	249	5.9	220	5.1
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract. (W78–W80)	60	1.4	70	1.6
Accidents caused by exposure to smoke, fire and flames. (X00–X09)	19	*	38	0.9
Accidental poisoning and exposure to noxious substances (X40–X49)	11	*	19	*
Other and unspecified accidents (W20–W31,W35–W64,W85–W99,X10–X39,X50–X59)	67	1.6	60	1.4
Assault (homicide) (*U01,X85–Y09)	337	7.9	352	8.2
Assault (homicide) by hanging, strangulation and suffocation. (X91)	31	0.7	30	0.7
Assault (homicide) by discharge of firearms. (*U01.4,X93–X95)	8	*	15	*
Neglect, abandonment and other maltreatment syndromes (Y06–Y07)	99	2.3	86	2.0
Assault (homicide) by other and unspecified means (*U01.0–*U01.3,*U01.5–*U01.9,X85–X90,X92,X96–X99,Y00–Y05,Y08–Y09)	199	4.7	221	5.1
Complications of medical and surgical care (Y40–Y84)	23	0.5	22	0.5
Other external causes. (Y10–Y36)	91	2.1	88	2.0

* Figure does not meet standards of reliability or precision; see "Technical Notes."

- Quantity zero.

¹New ICD-10 code U04 (Severe acute respiratory syndrome [SARS], unspecified) was added to the category in 2007.

²New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category in 2007.

NOTES: For certain causes of death such as unintentional injuries, homicides, suicides, and respiratory diseases, preliminary and final data differ because of the truncated nature of the preliminary file. Data are subject to sampling or random variation. For information regarding the calculation of standard errors and further discussion of the variability of the data, see "Technical Notes." Although the infant mortality rate is the preferred indicator of the risk of dying during the first year of life, another measure of infant mortality, the infant death rate, is shown elsewhere in this report. The two measures typically are similar, yet they can differ because the denominators used for these measures are different. For more information on these measures of risk, see "Infant mortality" section in "Technical Notes."

Table 6. Expectation of life, by age, race, and sex: United States, final 2007 and preliminary 2008

[Data are based on a continuous file of records from the states. Calculations of life expectancy employ populations estimated as of July 1 for 2008 and 2007; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported in 2008 by 34 states and the District of Columbia, and in 2007 by 27 states and the District of Columbia; see "Technical Notes." Multiple-race data for these states were bridged to single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes."]

Age (years) and race	Both sexes		Male		Female	
	2008	2007	2008	2007	2008	2007
All races¹						
0	77.8	77.9	75.3	75.4	80.3	80.4
1	77.3	77.5	74.8	74.9	79.8	79.9
5	73.4	73.6	70.9	71.0	75.8	76.0
10	68.5	68.6	66.0	66.1	70.9	71.0
15	63.5	63.7	61.0	61.1	65.9	66.1
20	58.7	58.8	56.3	56.4	61.0	61.2
25	54.0	54.1	51.6	51.8	56.2	56.3
30	49.2	49.4	47.0	47.1	51.3	51.5
35	44.5	44.6	42.3	42.5	46.5	46.7
40	39.8	39.9	37.7	37.8	41.7	41.9
45	35.2	35.4	33.2	33.3	37.1	37.2
50	30.8	30.9	28.8	29.0	32.5	32.7
55	26.5	26.7	24.7	24.9	28.1	28.2
60	22.4	22.5	20.8	20.9	23.8	23.9
65	18.6	18.6	17.1	17.2	19.8	19.9
70	14.9	15.0	13.6	13.7	15.9	16.0
75	11.6	11.7	10.5	10.6	12.4	12.5
80	8.8	8.8	7.9	7.9	9.3	9.4
85	6.4	6.5	5.8	5.8	6.8	6.8
90	4.6	4.6	4.1	4.1	4.8	4.8
95	3.2	3.2	2.9	2.9	3.3	3.3
100	2.3	2.3	2.1	2.1	2.3	2.3
White						
0	78.2	78.4	75.7	75.9	80.6	80.8
1	77.6	77.8	75.2	75.4	80.0	80.2
5	73.7	73.9	71.3	71.4	76.1	76.3
10	68.7	68.9	66.3	66.5	71.1	71.3
15	63.8	64.0	61.4	61.6	66.2	66.3
20	59.0	59.2	56.6	56.8	61.3	61.5
25	54.2	54.4	51.9	52.2	56.4	56.6
30	49.5	49.7	47.3	47.5	51.5	51.7
35	44.7	44.9	42.6	42.8	46.7	46.9
40	40.0	40.2	37.9	38.1	41.9	42.1
45	35.4	35.6	33.4	33.6	37.2	37.4
50	30.9	31.1	29.1	29.2	32.7	32.8
55	26.6	26.8	24.9	25.1	28.2	28.4
60	22.5	22.6	20.9	21.0	23.9	24.0
65	18.6	18.7	17.2	17.3	19.8	19.9
70	14.9	15.0	13.7	13.8	15.9	16.0
75	11.6	11.7	10.5	10.6	12.4	12.4
80	8.7	8.8	7.9	7.9	9.3	9.3
85	6.4	6.4	5.7	5.7	6.8	6.8
90	4.5	4.6	4.1	4.1	4.8	4.8
95	3.2	3.2	2.9	2.9	3.3	3.3
100	2.2	2.2	2.0	2.0	2.2	2.2

See footnotes at end of table.

Table 6. Expectation of life, by age, race, and sex: United States, final 2007 and preliminary 2008—Con.

[Data are based on a continuous file of records from the states. Calculations of life expectancy employ populations estimated as of July 1 for 2008 and 2007; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported in 2008 by 34 states and the District of Columbia, and in 2007 by 27 states and the District of Columbia; see "Technical Notes." Multiple-race data for these states were bridged to single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes"]

Age (years) and race	Both sexes		Male		Female	
	2008	2007	2008	2007	2008	2007
Black						
0	73.6	73.6	70.2	70.0	76.8	76.8
1	73.6	73.6	70.2	70.1	76.7	76.8
5	69.7	69.7	66.3	66.2	72.8	72.9
10	64.8	64.7	61.4	61.2	67.9	67.9
15	59.9	59.8	56.5	56.3	63.0	63.0
20	55.1	55.0	51.8	51.7	58.1	58.1
25	50.5	50.4	47.3	47.2	53.3	53.3
30	45.8	45.8	42.8	42.7	48.5	48.5
35	41.2	41.2	38.3	38.2	43.7	43.8
40	36.7	36.7	33.9	33.8	39.1	39.1
45	32.3	32.3	29.5	29.5	34.6	34.6
50	28.1	28.1	25.4	25.4	30.3	30.4
55	24.2	24.2	21.7	21.7	26.3	26.3
60	20.6	20.6	18.3	18.3	22.3	22.4
65	17.2	17.2	15.2	15.2	18.7	18.7
70	14.0	14.0	12.4	12.4	15.2	15.2
75	11.2	11.2	9.9	9.9	12.1	12.1
80	8.7	8.7	7.7	7.7	9.4	9.4
85	6.7	6.7	6.0	6.0	7.1	7.1
90	5.1	5.1	4.6	4.6	5.3	5.3
95	3.8	3.8	3.5	3.5	3.9	3.9
100	2.8	2.8	2.6	2.6	2.8	2.8

¹Includes races other than white and black.

NOTES: Life expectancies were calculated using a revised methodology and may differ from those previously published; see "Technical Notes." Data are subject to sampling or random variation.

Table 7. Deaths and death rates for the 10 leading causes of death in specified age groups: United States, preliminary 2008

[Data are based on a continuous file of records received from the states. Rates are per 100,000 population in specified group. For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Rank ¹	Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>) and age	Number	Rate
All ages ²			
...	All causes	2,472,699	813.2
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	617,527	203.1
2	Malignant neoplasms (C00–C97)	566,137	186.2
3	Chronic lower respiratory diseases (J40–J47)	141,075	46.4
4	Cerebrovascular diseases (I60–I69)	133,750	44.0
5	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	121,207	39.9
...	Motor vehicle accidents . . (V02–V04,V09.0,V09.2,V12–V14,V19.0–V19.2,V19.4–V19.6,V20–V79, V80.3–V80.5, V81.0–V81.1,V82.0–V82.1,V83–V86,V87.0–V87.8,V88.0–V88.8,V89.0,V89.2)	39,831	13.1
...	All other accidents . . (V01,V05–V06,V09.1,V09.3–V09.9,V10–V12,V15–V18,V19.3,V19.8–V19.9, V80.0–V80.2, V80.6–V80.9,V81.2–V81.9, V82.2–V82.9,V87.9,V88.9,V89.1,V89.3,V89.9,V90–V99,W00–X59,Y85–Y86)	81,376	26.8
6	Alzheimer's disease (G30)	82,476	27.1
7	Diabetes mellitus (E10–E14)	70,601	23.2
8	Influenza and pneumonia (J09–J18)	56,335	18.5
9	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	48,283	15.9
10	Septicemia (A40–A41)	35,961	11.8
...	All other causes (residual)	599,347	197.1
1–4 years			
...	All causes	4,752	28.5
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	1,469	8.8
...	Motor vehicle accidents . . (V02–V04,V09.0,V09.2,V12–V14,V19.0–V19.2,V19.4–V19.6,V20–V79, V80.3–V80.5,V81.0–V81.1,V82.0–V82.1,V83–V86,V87.0–V87.8,V88.0–V88.8,V89.0,V89.2)	467	2.8
...	All other accidents . . (V01,V05–V06,V09.1,V09.3–V09.9,V10–V12,V15–V18,V19.3,V19.8–V19.9,V80.0–V80.2, V80.6–V80.9,V81.2–V81.9,V82.2–V82.9,V87.9,V88.9,V89.1,V89.3,V89.9,V90–V99,W00–X59,Y85–Y86)	1,002	6.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	520	3.1
3	Assault (homicide) (*U01–*U02,X85–Y09,Y87.1)	421	2.5
4	Malignant neoplasms (C00–C97)	394	2.4
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	182	1.1
6	Influenza and pneumonia (J09–J18)	144	0.9
7	Septicemia (A40–A41)	95	0.6
8	Cerebrovascular diseases (I60–I69)	63	0.4
9	Certain conditions originating in the perinatal period (P00–P96)	56	0.3
10	Chronic lower respiratory diseases (J40–J47)	53	0.3
...	All other causes (residual)	1,355	8.1
5–14 years			
...	All causes	5,659	14.1
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	1,862	4.6
...	Motor vehicle accidents . . (V02–V04,V09.0,V09.2,V12–V14,V19.0–V19.2,V19.4–V19.6,V20–V79,V80.3–V80.5, V81.0–V81.1,V82.0–V82.1,V83–V86,V87.0–V87.8,V88.0–V88.8,V89.0,V89.2)	1,028	2.6
...	All other accidents . . (V01,V05–V06,V09.1,V09.3–V09.9,V10–V12,V15–V18,V19.3,V19.8–V19.9,V80.0–V80.2, V80.6–V80.9,V81.2–V81.9,V82.2–V82.9,V87.9,V88.9,V89.1,V89.3,V89.9,V90–V99,W00–X59,Y85–Y86)	834	2.1
2	Malignant neoplasms (C00–C97)	890	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	331	0.8
4	Assault (homicide) (*U01–*U02,X85–Y09,Y87.1)	320	0.8
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	228	0.6
6	Intentional self-harm (suicide) (*U03,X60–X84,Y87.0)	221	0.6
7	Chronic lower respiratory diseases (J40–J47)	117	0.3
8	Cerebrovascular diseases (I60–I69)	96	0.2
9	Influenza and pneumonia (J09–J18)	90	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	88	0.2
...	All other causes (residual)	1,416	3.51

See footnotes at end of table.

Table 7. Deaths and death rates for the 10 leading causes of death in specified age groups: United States, preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Rates are per 100,000 population in specified group. For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Rank ¹	Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>) and age	Number	Rate
15–24 years			
...	All causes	32,233	75.7
1	Accidents (unintentional injuries) (V01–X59, Y85–Y86)	14,020	32.9
...	Motor vehicle accidents . . (V02–V04, V09.0, V09.2, V12–V14, V19.0–V19.2, V19.4–V19.6, V20–V79, V80.3–V80.5, V81.0–V81.1, V82.0–V82.1, V83–V86, V87.0–V87.8, V88.0–V88.8, V89.0, V89.2)	8,959	21.0
...	All other accidents . . (V01, V05–V06, V09.1, V09.3–V09.9, V10–V12, V15–V18, V19.3, V19.8–V19.9, V80.0–V80.2, V80.6–V80.9, V81.2–V81.9, V82.2–V82.9, V87.9, V88.9, V89.1, V89.3, V89.9, V90–V99, W00–X59, Y85–Y86)	5,062	11.9
2	Assault (homicide) (*U01–*U02, X85–Y09, Y87.1)	5,285	12.4
3	Intentional self-harm (suicide) (*U03, X60–X84, Y87.0)	4,297	10.1
4	Malignant neoplasms (C00–C97)	1,659	3.9
5	Diseases of heart (I00–I09, I11, I13, I20–I51)	1,059	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	466	1.1
7	Influenza and pneumonia (J09–J18)	210	0.5
8	Diabetes mellitus (E10–E14)	204	0.5
9	Cerebrovascular diseases (I60–I69)	189	0.4
10	Human immunodeficiency virus (HIV) disease (B20–B24)	171	0.4
...	All other causes (residual)	4,673	11.0
25–44 years			
...	All causes	118,755	142.3
1	Accidents (unintentional injuries) (V01–X59, Y85–Y86)	30,318	36.3
...	Motor vehicle accidents . . (V02–V04, V09.0, V09.2, V12–V14, V19.0–V19.2, V19.4–V19.6, V20–V79, V80.3–V80.5, V81.0–V81.1, V82.0–V82.1, V83–V86, V87.0–V87.8, V88.0–V88.8, V89.0, V89.2)	12,262	14.7
...	All other accidents . . (V01, V05–V06, V09.1, V09.3–V09.9, V10–V12, V15–V18, V19.3, V19.8–V19.9, V80.0–V80.2, V80.6–V80.9, V81.2–V81.9, V82.2–V82.9, V87.9, V88.9, V89.1, V89.3, V89.9, V90–V99, W00–X59, Y85–Y86)	18,056	21.6
2	Malignant neoplasms (C00–C97)	16,235	19.5
3	Diseases of heart (I00–I09, I11, I13, I20–I51)	14,495	17.4
4	Intentional self-harm (suicide) (*U03, X60–X84, Y87.0)	11,959	14.3
5	Assault (homicide) (*U01–*U02, X85–Y09, Y87.1)	7,534	9.0
6	Human immunodeficiency virus (HIV) disease (B20–B24)	3,825	4.6
7	Chronic liver disease and cirrhosis (K70, K73–K74)	2,985	3.6
8	Cerebrovascular diseases (I60–I69)	2,571	3.1
9	Diabetes mellitus (E10–E14)	2,422	2.9
10	Influenza and pneumonia (J09–J18)	1,254	1.5
...	All other causes (residual)	25,157	30.2
45–64 years			
...	All causes	482,845	618.6
1	Malignant neoplasms (C00–C97)	154,697	198.2
2	Diseases of heart (I00–I09, I11, I13, I20–I51)	104,479	133.8
3	Accidents (unintentional injuries) (V01–X59, Y85–Y86)	32,817	42.0
...	Motor vehicle accidents . . (V02–V04, V09.0, V09.2, V12–V14, V19.0–V19.2, V19.4–V19.6, V20–V79, V80.3–V80.5, V81.0–V81.1, V82.0–V82.1, V83–V86, V87.0–V87.8, V88.0–V88.8, V89.0, V89.2)	10,463	13.4
...	All other accidents . . (V01, V05–V06, V09.1, V09.3–V09.9, V10–V12, V15–V18, V19.3, V19.8–V19.9, V80.0–V80.2, V80.6–V80.9, V81.2–V81.9, V82.2–V82.9, V87.9, V88.9, V89.1, V89.3, V89.9, V90–V99, W00–X59, Y85–Y86)	22,353	28.6
4	Chronic lower respiratory diseases (J40–J47)	18,416	23.6
5	Diabetes mellitus (E10–E14)	16,990	21.8
6	Chronic liver disease and cirrhosis (K70, K73–K74)	16,749	21.5
7	Cerebrovascular diseases (I60–I69)	16,595	21.3
8	Intentional self-harm (suicide) (*U03, X60–X84, Y87.0)	13,703	17.6
9	Septicemia (A40–A41)	7,068	9.1
10	Nephritis, nephrotic syndrome and nephrosis (N00–N07, N17–N19, N25–N27)	7,021	9.0
...	All other causes (residual)	94,310	120.8

See footnotes at end of table.

Table 7. Deaths and death rates for the 10 leading causes of death in specified age groups: United States, preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Rates are per 100,000 population in specified group. For explanation of asterisk (*) preceding cause-of-death codes, see "Technical Notes." Figures are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals]

Rank ¹	Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>) and age	Number	Rate
	65 years and over		
...	All causes	1,800,261	4,631.5
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	496,662	1,277.8
2	Malignant neoplasms (C00–C97)	392,184	1,009.0
3	Chronic lower respiratory diseases (J40–J47)	121,236	311.9
4	Cerebrovascular diseases (I60–I69)	114,088	293.5
5	Alzheimer's disease (G30)	81,624	210.0
6	Diabetes mellitus (E10–E14)	50,940	131.1
7	Influenza and pneumonia (J09–J18)	48,436	124.6
8	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	39,964	102.8
9	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	39,395	101.4
...	Motor vehicle accidents . . (V02–V04,V09.0,V09.2,V12–V14,V19.0–V19.2,V19.4–V19.6,V20–V79,V80.3–V80.5, V81.0–V81.1,V82.0–V82.1,V83–V86,V87.0–V87.8,V88.0–V88.8,V89.0,V89.2)	6,544	16.8
...	All other accidents . . (V01,V05–V06,V09.1,V09.3–V09.9,V10–V12,V15–V18,V19.3,V19.8–V19.9,V80.0–V80.2, V80.6–V80.9,V81.2–V81.9,V82.2–V82.9,V87.9,V88.9,V89.1,V89.3,V89.9,V90–V99,W00–X59,Y85–Y86)	32,851	84.5
10	Septicemia (A40–A41)	27,053	69.6
...	All other causes (residual)	388,679	1,000.0

... Category not applicable.

¹Based on number of deaths; see "Technical Notes."

²Includes deaths under age 1 year.

NOTES: For certain causes of death such as unintentional injuries, homicides, suicides, and respiratory diseases, preliminary and final data differ because of the truncated nature of the preliminary file. Data are subject to sampling or random variation. For information regarding the calculation of standard errors and further discussion of variability of the data, see "Technical Notes."

Table 8. Infant deaths and infant mortality rates for the 10 leading causes of infant death, by race and Hispanic origin: United States, preliminary 2008

[Data are based on a continuous file of records received from the states. Rates are per 100,000 live births. Figures are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals. Race and Hispanic origin are reported separately on both the birth and death certificate. Rates for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported for deaths by 34 states and the District of Columbia and for births by 30 states; see "Technical Notes." The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes." Data for persons of Hispanic origin are included in the data for each race group, according to the decedent's reported race; see "Technical Notes"]

Rank ¹	Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>) and race and Hispanic origin	Number	Rate
All races ²			
...	All causes	28,029	659.3
1	Congenital malformations, deformations and chromosomal abnormalities. (Q00–Q99)	5,647	132.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified. (P07)	4,733	111.3
3	Sudden infant death syndrome (R95)	2,292	53.9
4	Newborn affected by maternal complications of pregnancy (P01)	1,764	41.5
5	Accidents (unintentional injuries) (V01–X59)	1,299	30.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	1,073	25.2
7	Bacterial sepsis of newborn (P36)	696	16.4
8	Respiratory distress of newborn. (P22)	625	14.7
9	Diseases of the circulatory system (I00–I99)	590	13.9
10	Neonatal hemorrhage. (P50–P52,P54)	551	13.0
...	All other causes (residual)	8,759	206.0
Total white			
...	All causes	18,165	554.3
1	Congenital malformations, deformations and chromosomal abnormalities. (Q00–Q99)	4,289	130.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified. (P07)	2,656	81.1
3	Sudden infant death syndrome (R95)	1,453	44.3
4	Newborn affected by maternal complications of pregnancy (P01)	1,119	34.1
5	Accidents (unintentional injuries) (V01–X59)	837	25.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	693	21.1
7	Bacterial sepsis of newborn (P36)	431	13.2
8	Respiratory distress of newborn. (P22)	393	12.0
9	Diseases of the circulatory system (I00–I99)	383	11.7
10	Neonatal hemorrhage. (P50–P52,P54)	363	11.1
...	All other causes (residual)	5,548	169.3
Non-Hispanic white			
...	All causes	12,553	552.2
1	Congenital malformations, deformations and chromosomal abnormalities. (Q00–Q99)	2,833	124.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified. (P07)	1,751	77.0
3	Sudden infant death syndrome (R95)	1,168	51.4
4	Newborn affected by maternal complications of pregnancy (P01)	767	33.7
5	Accidents (unintentional injuries) (V01–X59)	667	29.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	469	20.6
7	Bacterial sepsis of newborn (P36)	300	13.2
8	Respiratory distress of newborn. (P22)	262	11.5
9	Diseases of the circulatory system (I00–I99)	257	11.3
10	Neonatal hemorrhage (P50–P52,P54)	245	10.8
...	All other causes (residual)	3,834	168.7
Total black			
...	All causes	8,509	1,267.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified. (P07)	1,858	276.8
2	Congenital malformations, deformations and chromosomal abnormalities. (Q00–Q99)	1,078	160.6
3	Sudden infant death syndrome (R95)	742	110.5
4	Newborn affected by maternal complications of pregnancy (P01)	578	86.1
5	Accidents (unintentional injuries) (V01–X59)	406	60.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	329	49.0
7	Bacterial sepsis of newborn (P36)	224	33.4
8	Necrotizing enterocolitis of newborn (P77)	218	32.5
9	Respiratory distress of newborn. (P22)	214	31.9
10	Diseases of the circulatory system (I00–I99)	168	25.0
...	All other causes (residual)	2,694	401.3

See footnotes at end of table.

Table 8. Infant deaths and infant mortality rates for the 10 leading causes of infant death, by race and Hispanic origin: United States, preliminary 2008—Con.

[Data are based on a continuous file of records received from the states. Rates are per 100,000 live births. Figures are based on weighted data rounded to the nearest individual, so categories may not add to totals or subtotals. Race and Hispanic origin are reported separately on both the birth and death certificate. Rates for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates; see "Technical Notes." Race categories are consistent with 1977 Office of Management and Budget (OMB) standards. Multiple-race data were reported for deaths by 34 states and the District of Columbia and for births by 30 states; see "Technical Notes." The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards for comparability with other states; see "Technical Notes." Data for persons of Hispanic origin are included in the data for each race group, according to the decedent's reported race; see "Technical Notes"]

Rank ¹	Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>) and race and Hispanic origin	Number	Rate
	Hispanic ³		
...	All causes	5,884	566.4
1	Congenital malformations, deformations and chromosomal abnormalities. (Q00–Q99)	1,503	144.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified. (P07)	937	90.2
3	Newborn affected by maternal complications of pregnancy (P01)	362	34.8
4	Sudden infant death syndrome (R95)	314	30.2
5	Newborn affected by complications of placenta, cord and membranes (P02)	236	22.7
6	Accidents (unintentional injuries) (V01–X59)	181	17.4
7	Bacterial sepsis of newborn (P36)	138	13.3
8	Respiratory distress of newborn. (P22)	137	13.2
9	Diseases of the circulatory system (I00–I99)	131	12.6
10	Neonatal hemorrhage. (P50–P52,P54)	129	12.4
...	All other causes (residual)	1,816	174.8

... Category not applicable.

¹Based on number of deaths; see "Technical Notes."

²Includes races other than white and black.

³Includes all persons of Hispanic origin of any race; see "Technical Notes."

NOTES: For certain causes of death such as unintentional injuries, homicides, suicides, and sudden infant death syndrome, preliminary and final data differ because of the truncated nature of the preliminary file. Data are subject to sampling or random variation. For information regarding the calculation of standard errors and further discussion of variability of the data, see "Technical Notes." Although the infant mortality rate is the preferred indicator of the risk of dying during the first year of life, another measure of infant mortality, the infant death rate, is shown elsewhere in this report. The two measures typically are similar but use different denominators.

Technical Notes

Nature and sources of data

Preliminary mortality data for 2008 are based on a continuous receipt and processing of statistical records by the Centers for Disease Control and Prevention's National Center for Health Statistics (NCHS) through September 22, 2010. NCHS received the data from the states' vital registration systems through the Vital Statistics Cooperative Program. Demographic information for the United States was available in calendar year 2008 for an estimated 99.9 percent of infant decedents and 100 percent of decedents aged 1 year and over. Medical information for the United States was available in calendar year 2008 for an estimated 99.2 percent of infant decedents and 99.5 percent of decedents aged 1 year and over. In this report, U.S. totals include only events occurring within the 50 states and the District of Columbia. Data for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas included in tables showing data by state are not included in U.S. totals.

For 2008, individual records of infant deaths (deaths under age 1 year) and deaths of persons aged 1 year and over are weighted (when necessary) to independent counts of deaths occurring in each state. These state-specific counts serve as control totals and are the basis for the record weights in the preliminary file. If the number of records in the preliminary file is greater than the count received from the state, the state-specific number of records in the preliminary file is used instead and the weight is set at 1.0.

For this report, two separate files are processed. The medical file, or cause-of-death file, contains records that include both demographic and medical information used to generate tables showing cause of death. The demographic file, which includes records from the medical file as well as additional records containing demographic information only, is used to generate tables showing mortality by demographic characteristics only. A state-specific weight is computed for each file by dividing the state control total by the number of records in the preliminary sample.

Each record is assigned two weights, a state-specific weight and a U.S. weight. State weights are used for state-specific tabulations, and U.S. weights are used for national tabulations. For the medical file, the state weight makes the death counts comparable with those in the demographic file. The U.S. weight combines two factors: one to make the medical file counts for the individual record's state comparable with those for the demographic file, and one to compensate for any states not represented in the file. This second factor is equivalent to zero if all states are represented in the file. Thus, when all states are represented in the preliminary files, the state and U.S. weights are the same.

Because there are two separate files, each with two separate sets of weights, slight inconsistencies may occur between the demographic and medical tables in this report. [Table 1](#) shows the percent completeness of the preliminary files by place of occurrence for the United States and each state. The percent completeness is obtained by dividing the number of records in the preliminary files by the control total and multiplying by 100. Although data by place of occurrence are used to compute the weights, all data in this report are tabulated by place of residence. Because preliminary records for Nevada were missing geographic information indicating whether the decedent resided in that state, all records filed by Nevada are considered as residents of

Nevada. Therefore, data shown in [Table 3](#) (which shows deaths and death rates by state) for Nevada are approximately 8 percent higher than if nonresidents for that state were excluded. (The estimate for the overcount is based on the percentage of deaths occurring to nonresidents reported by Nevada in 2006 and 2007.)

For selected variables in the mortality files, unknown or not-stated values are imputed. The percentage not stated was less than 1.7 percent for all variables discussed in this report. Detailed information on reporting completeness and imputation procedures may be found in "Technical Appendix, Vital Statistics of the United States: Mortality, 1999" (32).

2003 revision of U.S. Standard Certificate of Death

This report includes data for the 30 states and the District of Columbia that implemented the 2003 revision of the U.S. Standard Certificate of Death by 2008: Arkansas, California, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Michigan, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Dakota, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Vermont, Washington, and Wyoming. (Because Vermont implemented the 2003 certificate revision in July 2008, a portion of the state's data for that year is based on the 1989 revision of the certificate.) The remaining 20 states collected and reported death data in 2008 based on the 1989 revision of the U.S. Standard Certificate of Death. The 2003 revision is described in detail elsewhere (15,16). Because the items presented in this report appear largely comparable despite changes to item wording and format in the 2003 revision, data from both groups of states are combined.

Race and Hispanic origin

The 2003 revision of the U.S. Standard Certificate of Death allows the reporting of more than one race (multiple races) (15). This change was implemented to reflect the increasing diversity of the population of the United States, to be consistent with the decennial census, and to reflect standards issued in 1997 by the Office of Management and Budget (OMB). OMB standards mandate the collection of more than one race for federal data (see "Population denominators") (21). In addition, the new certificate is compliant with the OMB-mandated minimum set of five races to be reported for federal data. These are white, black or African American, American Indian or Alaska Native (AIAN), Asian, and Native Hawaiian or Other Pacific Islander (NHOPI).

For 2008 mortality data, multiple races were reported on the revised death certificates of Arkansas, California, Connecticut, Delaware, District of Columbia, Florida, Georgia, Idaho, Illinois, Indiana, Kansas, Michigan, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Dakota, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Vermont, Washington, and Wyoming. Multiple races were also reported on the unrevised certificates of Hawaii, Maine, Minnesota, and Wisconsin. Data from the vital records of the remaining 16 states are based on the 1989 revision of the U.S. Standard Certificate of Death, which follows the 1977 OMB standards, allowing only a single race to be reported (16,17). In addition, these 16 states report a minimum set of

Table I. Total count of records and percent completeness of preliminary files of infant deaths and deaths to those aged 1 year and over: United States, each state and territory, preliminary 2008

[By place of occurrence]

Area	Infant deaths (aged under 1 year)			Deaths to those aged 1 year and over		
	Count of records	Percent completeness		Count of records	Percent completeness	
		Demographic file	Medical file		Demographic file	Medical file
United States ¹	28,095	99.9	99.2	2,449,367	100.0	99.5
Alabama	616	100.0	100.0	46,533	100.0	100.0
Alaska	61	100.0	100.0	3,389	100.0	100.0
Arizona	635	99.7	99.7	45,960	99.9	99.9
Arkansas	281	100.0	100.0	28,559	100.0	100.0
California	2,830	100.0	100.0	232,867	100.0	100.0
Colorado	468	100.0	100.0	31,358	100.0	100.0
Connecticut	227	100.0	100.0	28,710	100.0	100.0
Delaware	115	100.0	100.0	7,466	100.0	100.0
District of Columbia	226	99.6	97.3	6,104	100.0	99.9
Florida	1,712	100.0	100.0	170,110	100.0	100.0
Georgia	1,169	97.5	87.3	68,994	100.0	85.1
Hawaii	107	100.0	100.0	9,465	100.0	100.0
Idaho	129	100.0	100.0	10,690	100.0	100.0
Illinois	1,120	100.0	100.0	99,635	99.9	99.9
Indiana	614	100.0	100.0	56,578	100.0	100.0
Iowa	210	100.0	100.0	28,275	100.0	100.0
Kansas	278	100.0	100.0	24,000	100.0	100.0
Kentucky	352	100.0	100.0	40,739	100.0	100.0
Louisiana	603	100.0	99.2	40,553	100.0	100.0
Maine	74	100.0	100.0	12,371	100.0	99.8
Maryland	538	100.0	100.0	43,195	100.0	100.0
Massachusetts	400	100.0	100.0	53,940	100.0	100.0
Michigan	900	100.0	100.0	86,362	100.0	100.0
Minnesota	449	100.0	100.0	38,151	100.0	100.0
Mississippi	395	100.0	100.0	28,003	100.0	100.0
Missouri	671	100.0	100.0	57,706	100.0	100.0
Montana	78	98.7	97.4	8,835	100.0	100.0
Nebraska	157	100.0	100.0	15,502	100.0	100.0
Nevada	212	93.4	93.4	19,856	97.7	97.7
New Hampshire	52	100.0	100.0	10,102	100.0	100.0
New Jersey	558	100.0	98.7	68,121	100.0	99.5
New Mexico	159	100.0	95.6	15,433	100.0	99.5
New York	1,391	100.0	100.0	146,756	100.0	100.0
New York excluding New York City . .	693	100.0	100.0	93,253	100.0	100.0
New York City	698	100.0	100.0	53,503	100.0	100.0
North Carolina	1,090	100.0	100.0	77,017	100.0	100.0
North Dakota	54	100.0	100.0	6,286	100.0	100.0
Ohio	1,178	100.0	98.6	108,544	100.0	99.9
Oklahoma	404	100.0	97.0	35,724	100.0	99.9
Oregon	262	99.6	99.6	31,594	100.0	100.0
Pennsylvania	1,152	100.0	100.0	127,359	100.0	100.0
Rhode Island	84	100.0	98.8	9,892	100.0	99.7
South Carolina	473	100.0	100.0	38,890	99.9	99.9
South Dakota	105	100.0	100.0	7,141	100.0	100.0
Tennessee	784	100.0	100.0	61,063	99.9	99.9
Texas	2,540	100.0	100.0	165,118	100.0	99.9
Utah	292	100.0	100.0	14,148	100.0	100.0
Vermont	24	100.0	100.0	5,154	100.0	100.0
Virginia	698	100.0	100.0	57,769	100.0	100.0
Washington	502	100.0	100.0	48,254	100.0	100.0
West Virginia	158	100.0	100.0	21,056	100.0	99.8
Wisconsin	487	100.0	100.0	46,124	100.0	100.0
Wyoming	29	96.6	96.6	3,946	100.0	100.0
Puerto Rico	384	100.0	99.7	28,545	99.7	99.5
Virgin Islands	6	100.0	100.0	692	100.0	100.0
Guam	29	100.0	37.9	758	100.0	60.0
American Samoa	13	100.0	100.0	232	100.0	100.0
Northern Marianas	---	---	---	---	---	---

--- Data not available.

¹Excludes data for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas.

NOTE: Percent completeness equals 100 times the number of records in the preliminary file divided by the count of records.

four races as stipulated in the 1977 standards. These are white, black or African American, AIAN, and Asian or Pacific Islander (API).

In order to provide uniformity and comparability of data during the transition period, before all or most of the data becomes available in the new multiple-race format, it is necessary to adjust the data for those states reporting multiple race by “bridging” the multiple-race information reported for decedents to a single race. The bridging procedure used for mortality numerators is similar to the procedure used to bridge multiracial population estimates (see “Population denominators”) (19,20). Multiracial decedents are imputed to a single race (white, black, AIAN, or API) according to the combination of races, Hispanic origin, sex, and age indicated on the death certificate. The imputation procedure is described in detail at http://www.cdc.gov/nchs/data/dvs/Multiple_race_docu_5-10-04.pdf.

Because race and Hispanic origin are reported separately on the death certificate, data shown by race include persons of Hispanic or non-Hispanic origin, and data for Hispanic origin include persons of any race. In this report, unless otherwise specified, deaths of persons of Hispanic origin are included in the totals for each race group—white, black, AIAN, and API—according to the decedent’s race as reported on the death certificate. Data shown for Hispanic persons include all persons of Hispanic origin of any race. Mortality data on the population of Hispanic origin are based on deaths in all states. Death rates for Hispanic, AIAN, and API persons should be interpreted with caution because of inconsistencies in reporting race on death certificates compared with such reporting on censuses, surveys, and birth certificates. Studies have shown underreporting on death certificates of AIAN, API, and Hispanic decedents, as well as undercounts of these groups in censuses (22–24).

Injury at work

Information on deaths attributed to injuries at work is derived from a separate item on the death certificate that asks the medical certifier whether the death resulted from an injury sustained at work. The item is on the death certificate of all states. Number of deaths, crude death rates, and age-adjusted death rates for injury at work for those aged 15 and over, excluding age not stated, are shown in [Table 2](#). Age-adjusted death rates presented in this report for injury at work were computed using age-specific death rates and the U.S. standard population based on year 2000 standard for ages 15 and over, excluding age not stated ([Table V](#)). If the estimated “employed” population aged 15 and over had been used instead in the denominator, higher death rates would have resulted, especially for population groups with lower employment rates (see “Computing rates and percentages”).

Cause-of-death classification

Mortality statistics are compiled in accordance with World Health Organization (WHO) regulations specifying that member nations classify and code causes of death in accordance with the current revision of the *International Statistical Classification of Diseases and Related Health Problems* (ICD). The ICD provides the basic guidance used in virtually all countries to code and classify causes of death. It provides not only disease, injury, and poisoning categories but also the rules used to select the single underlying cause of death for tabulation from the several diagnoses that may be reported on a

single death certificate, as well as definitions, tabulation lists, the format of the death certificate, and regulations on the use of the classification. Causes of death for data presented in this report were coded according to ICD guidelines, which are described in annual issues of part 2a of the “NCHS Instruction Manual” (33).

Effective with deaths occurring in 1999, the United States began using the Tenth Revision of the ICD (ICD–10) (34). In 2004, the Second Edition of ICD–10 was adopted (35). During 1979–1998, causes of death were coded and classified according to the Ninth Revision (ICD–9) (36). The change from ICD–9 to ICD–10 resulted in discontinuities for selected cause-of-death trends. These discontinuities are measured using comparability ratios derived from a comparability study (37).

Beginning with data for 2001, NCHS introduced categories *U01–*U03 for classifying and coding deaths due to acts of terrorism. The asterisks before the category codes indicate that they are not part of ICD–10. Deaths classified to the terrorism categories are included in the categories Assault (homicide) and Intentional self-harm (suicide) for the 113 causes-of-death list ([Table 2](#)) and Assault (homicide) in the 130 causes-of-infant death list ([Table 5](#)). Additional information on the new terrorism categories can be found at http://www.cdc.gov/nchs/icd/terrorism_code.htm. No deaths occurring in 2007 and 2008 were classified to the terrorism categories.

Enterocolitis due to Clostridium difficile (*C. difficile*)—Due to growing concerns about the increasing number of deaths from Enterocolitis due to *Clostridium difficile* (ICD–10 code A04.7), beginning in 2006, *C. difficile* deaths are included separately as a rankable cause of death in tables showing data for 113 selected causes of death ([Table 2](#)). The number of deaths due to this cause increased by 17.4 percent from 6,372 in 2007 to 7,483 in 2008. The age-adjusted death rate for this cause increased significantly from 2.0 per 100,000 standard population in 2007 to 2.3 in 2008.

Codes for drug-induced deaths—The list of codes included in drug-induced causes was expanded in data year 2003 to be more comprehensive. The following 37 ICD–10 codes were added to the previous list of drug-induced codes: D52.1, D59.0, D59.2, D61.1, D64.2, E06.4, E16.0, E23.1, E24.2, E27.3, E66.1, G21.1, G24.0, G25.1, G25.4, G25.6, G44.4, G62.0, G72.0, I95.2, J70.2, J70.3, J70.4, L10.5, L27.0, L27.1, M10.2, M32.0, M80.4, M81.4, M83.5, M87.1, R78.1, R78.2, R78.3, R78.4, and R78.5. In addition to expansion of the list in 2003, ICD codes K85.3 (drug-induced acute pancreatitis) and R50.2 (drug-induced fever) were added to the list of drug-induced codes in 2006. No deaths were classified to these two new codes in 2007 and two deaths were assigned to K85.3 in 2008.

Codes for alcohol-induced deaths—The list of codes included in alcohol-induced causes was expanded in data year 2003 to be more comprehensive. Three ICD–10 codes were added to the previous list of alcohol-induced codes: E24.4, G72.1, and K86.0. Additionally, K85.2 (alcohol-induced acute pancreatitis) was added to the list in 2006. In 2007, 400 deaths were classified to K85.2; in 2008, 395 deaths were classified to K85.2.

Recently added codes—Beginning with data for 2007, NCHS introduced category J09 for classifying and coding deaths for influenza due to identified avian influenza virus; U04.9 for Severe Acute Respiratory Syndrome (SARS); X59.0 for exposure to unspecified factor causing fracture; and X59.9 for exposure to unspecified factor causing other and unspecified injury. Deaths classified to code J09 are included in the category Influenza and pneumonia. Deaths classified to code

U04.9 are included in the category Other acute lower respiratory infections. Deaths for codes X59.0 and X59.9 are included in Other and unspecified nontransport accidents and their sequelae in the list of 113 selected causes of death (Tables 2 and 7) and the list of 130 selected causes of infant death (Table 5). Additional information on these new categories is available from: <http://www.cdc.gov/nchs/data/dvs/Part9InstructionManual2007.pdf> (38).

Changes to classification rules affecting respiratory conditions— Rules governing coding and classifying the causes of death sometimes require modifications as recommended by WHO. This often happens when evidence suggests that the changes will improve the quality of cause-of-death data or better describe the condition under consideration. These changes, however, may affect comparability of data between years for selected causes of death. In 2008, the implementation of changes in coding rules had an impact on Chronic lower respiratory diseases (ICD–10 codes J40–J47) and, therefore, changes in mortality statistics for this condition must be interpreted with caution. Chronic obstructive pulmonary disease with acute lower respiratory infection (ICD–10 code J44.0) is a component condition of the larger category Chronic lower respiratory diseases (CLRD). This component condition J44.0 increased as a proportion of all deaths from Chronic lower respiratory diseases between 2007 and 2008 (from 0.2 percent of CLRD deaths in 2007 to 17.3 percent of CLRD deaths in 2008). A yet-unspecified portion of this increase is due to a change in the rules that govern coding and classifying to this condition. Specifically, many deaths that would have previously been assigned to Chronic obstructive pulmonary disease, unspecified (ICD–10 code J44.9); pneumonia (J12–J16, J18); and Other acute lower respiratory infections (ICD–10 codes J20–J22) are now being classified to J44.0. The Mortality Statistics Branch plans to conduct a thorough analysis on this change and its effect on CLRD once the final file for 2008 is produced. The actual change to the coding and classification rules can be seen on page E-240 of the instruction manual *ICD–10 ACME Decision Tables for Classifying Underlying Causes of Death, 2008*, available from <http://www.cdc.gov/nchs/data/dvs/2008Final2C.pdf> (39). (Modifications relative to earlier rules can be identified by asterisks placed to the right of table entries.)

Nonsampling error

Causes of death in this report are subject to nonsampling error. This is because the preliminary file is processed before a full year's worth of data is available. The file is thus subject to the seasonality of certain causes of death that may not be equally distributed throughout the year. It is known, for example, that external causes such as unintentional injuries occur disproportionately during the summer months and that fatal respiratory conditions are more prevalent during the winter months. Accordingly, the truncated nature of the preliminary file may systematically overemphasize or underemphasize causes with pronounced seasonality, particularly when these deaths cluster at the end of the year. However, in years where the preliminary file completeness is more than 90 percent, it is unlikely that seasonality is a major factor.

Furthermore, for some deaths, especially those subject to medicolegal investigation such as unintentional injuries, homicides, suicides, and sudden infant death syndrome (SIDS), the final cause may not be available at the time the preliminary file is processed. In those cases,

the causes of death may be reported in the preliminary file as unknown or pending investigation and coded to the category Other ill-defined and unspecified causes of mortality (ICD–10 code R99), a subcategory of Symptoms, signs, and abnormal clinical and laboratory findings, not elsewhere classified (ICD–10 codes R00–R99). In the final data, some of the deaths of unknown cause in the preliminary file will be reassigned to specific causes if further, more specific cause-of-death information is provided.

A quantitative assessment of the degree of the nonsampling error can be made by comparing final data and preliminary data for the same year. A comparison of such data for the selected 113 causes of death for the total U.S. population for 2005–2007 indicates that preliminary estimates for some causes of death are sometimes underestimated and sometimes overestimated in the preliminary file (Table II). Thus, the number of deaths for unintentional injuries was underestimated in the preliminary file by 2.5 percent in 2005, 3.2 percent in 2006, and 5.4 percent in 2007. Similar undercounts occurred for suicide, with preliminary underestimates of 2.7 percent in 2005, 3.4 percent in 2006, and 4.1 percent in 2007. Likewise, homicide showed a 2.4 percent underestimate in the preliminary file in 2005, 2.9 percent in 2006, and 4.6 percent in 2007. Chronic lower respiratory disease was estimated with high accuracy in 2005 and 2006 but was overestimated in the 2007 preliminary file by 1.1 percent (Table II).

Comparisons showing nonsampling error in preliminary estimates for causes of infant death are shown in Table III, where Disorders related to short gestation and low birth weight, not elsewhere classified (P07) was underestimated 0.1 percent in 2005, estimated accurately in 2006, and underestimated 3.7 percent in 2007. Unintentional injuries (V01–X59) and SIDS (R95) were underestimated in the preliminary data for each of the three years of 2005–2007, with unintentional injuries being underestimated between 1.3 percent and 3.7 percent, and SIDS between 5.5 percent and 13.7 percent (Table III).

Tabulation lists and cause-of-death ranking

Tabulation lists for ICD–10 are published in the “NCHS Instruction Manual, Part 9, ICD–10 Cause-of-Death Lists for Tabulating Mortality Statistics, Updated October 2007” (38). For this report, two tabulation lists are used: the List of 113 Selected Causes of Death used for deaths of all ages, and the List of 130 Selected Causes of Infant Death used for infants. Modifications in the lists reflecting changes in ICD codes are footnoted in the report tables. These lists are also used to rank leading causes of death for the two population groups (40). For the List of 113 Selected Causes of Death, the group titles Major cardiovascular diseases (ICD–10 codes I00–I78) and Symptoms, signs, and abnormal clinical and laboratory findings, not elsewhere classified (ICD–10 codes R00–R99) are not ranked. In addition, category titles that begin with the words “Other” and “All other” are not ranked to determine the leading causes of death. When one of the titles that represents a subtotal is ranked [for example, Tuberculosis (ICD–10 codes A16–A19)], its component parts are not ranked [in this case, Respiratory tuberculosis (ICD–10 code A16) and Other tuberculosis (ICD–10 codes A17–A19)]. For the List of 130 Selected Causes of Infant Death, the same ranking procedures are used, except that the category Major cardiovascular diseases is not on the list.

Table II. Ratios of preliminary to final reported numbers of deaths from 113 selected causes: United States, 2005–2007

[For explanation of asterisk preceding cause-of-death codes, see "Technical Notes"]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Preliminary number of deaths 2007	Final number of deaths 2007	Ratio of preliminary to final 2007	Preliminary number of deaths 2006	Final number of deaths 2006	Ratio of preliminary to final 2006	Preliminary number of deaths 2005	Final number of deaths 2005	Ratio of preliminary to final 2005
All causes	2,424,059	2,423,712	1.0001	2,425,901	2,426,264	0.9999	2,447,910	2,448,017	1.0000
Salmonella infections (A01–A02)	30	30	1.0000	33	34	0.9706	30	30	1.0000
Shigellosis and amebiasis (A03,A06)	4	4	1.0000	4	6	0.6667	9	10	0.9000
Certain other intestinal infections (A04,A07–A09)	6,822	6,758	1.0095	6,671	6,639	1.0048	5,694	5,667	1.0048
Tuberculosis (A16–A19)	541	554	0.9765	644	652	0.9877	646	648	0.9969
Respiratory tuberculosis (A16)	410	424	0.9670	485	490	0.9898	477	480	0.9938
Other tuberculosis (A17–A19)	131	130	1.0077	159	162	0.9815	169	168	1.0060
Whooping cough (A37)	11	9	1.2222	9	9	1.0000	32	31	1.0323
Scarlet fever and erysipelas (A38,A46)	3	3	1.0000	2	2	1.0000	3	3	1.0000
Meningococcal infection (A39)	73	87	0.8391	103	105	0.9810	119	123	0.9675
Septicemia (A40–A41)	34,851	34,828	1.0007	34,031	34,234	0.9941	34,142	34,136	1.0002
Syphilis (A50–A53)	50	42	1.1905	35	38	0.9211	46	47	0.9787
Acute poliomyelitis (A80)	2	–	...	–	–	...	4	–	...
Arthropod-borne viral encephalitis (A83–A84,A85.2)	2	3	0.6667	5	5	1.0000	6	6	1.0000
Measles (B05)	–	–	...	–	–	...	1	1	1.0000
Viral hepatitis (B15–B19)	7,313	7,407	0.9873	6,021	7,250	0.8305	5,524	5,529	0.9991
Human immunodeficiency virus (HIV) disease (B20–B24)	11,061	11,295	0.9793	12,045	12,113	0.9944	12,456	12,543	0.9931
Malaria (B50–B54)	4	5	0.8000	9	9	1.0000	6	6	1.0000
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05, A20–A36,A42–A44,A48–A49,A54–A79,A81–A82,A85.0–A85.1,A85.8,A86–B04,B06–B09, B25–B49,B55–B99)	5,774	5,825	0.9912	7,110	5,897	1.2057	7,695	7,727	0.9959
Malignant neoplasms (C00–C97)	560,187	562,875	0.9952	560,102	559,888	1.0004	559,300	559,312	1.0000
Malignant neoplasms of lip, oral cavity and pharynx (C00–C14)	7,950	8,067	0.9855	7,727	7,720	1.0009	7,775	7,773	1.0003
Malignant neoplasm of esophagus (C15)	13,488	13,592	0.9923	13,674	13,686	0.9991	13,512	13,499	1.0010
Malignant neoplasm of stomach (C16)	11,308	11,388	0.9930	11,354	11,345	1.0008	11,473	11,514	0.9964
Malignant neoplasms of colon, rectum and anus (C18–C21)	53,100	53,586	0.9909	53,465	53,549	0.9984	53,228	53,252	0.9995
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	17,033	17,146	0.9934	16,447	16,525	0.9953	16,049	16,076	0.9983
Malignant neoplasm of pancreas (C25)	34,032	34,117	0.9975	33,437	33,454	0.9995	32,741	32,760	0.9994
Malignant neoplasm of larynx (C32)	3,680	3,634	1.0127	3,824	3,821	1.0008	3,790	3,797	0.9982
Malignant neoplasms of trachea, bronchus and lung (C33–C34)	158,258	158,760	0.9968	158,525	158,664	0.9991	159,415	159,292	1.0008
Malignant melanoma of skin (C43)	8,499	8,461	1.0045	8,487	8,441	1.0054	8,368	8,345	1.0028
Malignant neoplasm of breast (C50)	40,514	40,970	0.9889	41,223	41,210	1.0003	41,471	41,491	0.9995
Malignant neoplasm of cervix uteri (C53)	3,942	4,021	0.9804	3,926	3,976	0.9874	3,914	3,924	0.9975
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54–C55)	7,319	7,456	0.9816	7,374	7,384	0.9986	7,070	7,096	0.9963
Malignant neoplasm of ovary (C56)	14,535	14,621	0.9941	14,906	14,857	1.0033	14,770	14,787	0.9989
Malignant neoplasm of prostate (C61)	28,823	29,093	0.9907	28,331	28,372	0.9986	28,916	28,905	1.0004
Malignant neoplasms of kidney and renal pelvis (C64–C65)	12,569	12,703	0.9895	12,376	12,379	0.9998	12,513	12,517	0.9997
Malignant neoplasm of bladder (C67)	13,827	13,843	0.9988	13,492	13,474	1.0013	13,258	13,253	1.0004
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70–C72)	13,172	13,234	0.9953	12,853	12,886	0.9974	13,149	13,152	0.9998
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81–C96)	54,950	54,991	0.9993	55,241	55,045	1.0036	55,023	55,028	0.9999
Hodgkin's disease (C81)	1,251	1,271	0.9843	1,331	1,327	1.0030	1,272	1,272	1.0000
Non-Hodgkin's lymphoma (C82–C85)	20,537	20,528	1.0004	20,663	20,594	1.0034	20,863	20,873	0.9995
Leukemia (C91–C95)	21,696	21,825	0.9941	22,022	21,944	1.0036	21,616	21,623	0.9997
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	11,420	11,307	1.0100	11,153	11,111	1.0038	11,213	11,200	1.0012

See footnotes at end of table.

Table II. Ratios of preliminary to final reported numbers of deaths from 113 selected causes: United States, 2005–2007—Con.

[For explanation of asterisk preceding cause-of-death codes, see “Technical Notes”]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Preliminary number of deaths 2007	Final number of deaths 2007	Ratio of preliminary to final 2007	Preliminary number of deaths 2006	Final number of deaths 2006	Ratio of preliminary to final 2006	Preliminary number of deaths 2005	Final number of deaths 2005	Ratio of preliminary to final 2005
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	47	60	0.7833	71	69	1.0290	60	60	1.0000
All other and unspecified malignant neoplasms (C17,C23–C24,C26–C31,C37–C41,C44–C49,C51–C52,C57–C60,C62–C63,C66,C68–C69,C73–C80,C97)	63,186	63,192	0.9999	63,441	63,100	1.0054	62,865	62,851	1.0002
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	14,151	14,204	0.9963	14,101	14,122	0.9985	13,732	13,710	1.0016
Anemias (D50–D64)	4,848	4,829	1.0039	4,007	3,996	1.0028	4,611	4,624	0.9972
Diabetes mellitus (E10–E14)	70,905	71,382	0.9933	72,507	72,449	1.0008	74,817	75,119	0.9960
Nutritional deficiencies (E40–E64)	2,810	2,852	0.9853	2,568	2,556	1.0047	3,185	3,183	1.0006
Malnutrition (E40–E46)	2,600	2,644	0.9834	2,383	2,377	1.0025	3,004	3,003	1.0003
Other nutritional deficiencies (E50–E64)	210	208	1.0096	185	179	1.0335	181	180	1.0056
Meningitis (G00,G03)	626	655	0.9557	632	634	0.9968	652	669	0.9746
Parkinson’s disease (G20–G21)	20,136	20,058	1.0039	19,660	19,566	1.0048	19,547	19,544	1.0002
Alzheimer’s disease (G30)	74,944	74,632	1.0042	72,914	72,432	1.0067	71,696	71,599	1.0014
Major cardiovascular diseases (I00–I78)	803,504	806,156	0.9967	821,494	823,746	0.9973	853,188	856,030	0.9967
Diseases of heart (I00–I09,I11,I13,I20–I51)	615,651	616,067	0.9993	629,191	631,636	0.9961	649,399	652,091	0.9959
Acute rheumatic fever and chronic rheumatic heart diseases (I00–I09)	3,188	3,201	0.9959	3,257	3,257	1.0000	3,359	3,365	0.9982
Hypertensive heart disease (I11)	30,354	30,780	0.9862	29,217	29,788	0.9808	28,902	29,282	0.9870
Hypertensive heart and renal disease (I13)	2,954	2,987	0.9890	2,919	2,918	1.0003	3,148	3,172	0.9924
Ischemic heart diseases (I20–I25)	403,741	406,351	0.9936	424,892	425,425	0.9987	443,891	445,687	0.9960
Acute myocardial infarction (I21–I22)	132,841	132,968	0.9990	141,965	141,462	1.0036	150,852	151,004	0.9990
Other acute ischemic heart diseases (I24)	4,046	4,092	0.9888	3,938	3,932	1.0015	3,556	3,565	0.9975
Other forms of chronic ischemic heart disease (I20,I25)	266,854	269,291	0.9910	278,988	280,031	0.9963	289,483	291,118	0.9944
Atherosclerotic cardiovascular disease, so described (I25.0)	57,639	59,051	0.9761	59,734	61,030	0.9788	61,864	62,799	0.9851
All other forms of chronic ischemic heart disease (I20,I25.1–I25.9)	209,215	210,240	0.9951	219,254	219,001	1.0012	227,619	228,319	0.9969
Other heart diseases (I26–I51)	175,413	172,748	1.0154	168,906	170,248	0.9921	170,099	170,585	0.9972
Acute and subacute endocarditis (I33)	1,206	1,225	0.9845	1,209	1,216	0.9942	1,203	1,209	0.9950
Diseases of pericardium and acute myocarditis (I30–I31,I40)	843	867	0.9723	784	816	0.9608	836	864	0.9676
Heart failure (I50)	57,235	56,565	1.0118	60,315	60,337	0.9996	59,001	58,933	1.0012
All other forms of heart disease (I26–I28,I34–I38,I42–I49,I51)	116,129	114,091	1.0179	106,598	107,879	0.9881	109,059	109,579	0.9953
Essential hypertension and hypertensive renal disease (I10,I12,I15) ¹	23,769	23,965	0.9918	23,985	23,855	1.0054	24,865	24,902	0.9985
Cerebrovascular diseases (I60–I69)	133,990	135,952	0.9856	137,265	137,119	1.0011	143,497	143,579	0.9994
Atherosclerosis (I70)	8,223	8,232	0.9989	8,619	8,652	0.9962	11,833	11,841	0.9993
Other diseases of circulatory system (I71–I78)	21,872	21,940	0.9969	22,435	22,484	0.9978	23,594	23,617	0.9990
Aortic aneurysm and dissection (I71)	12,887	12,986	0.9924	13,178	13,238	0.9955	13,811	13,843	0.9977
Other diseases of arteries, arterioles and capillaries (I72–I78)	8,985	8,954	1.0035	9,258	9,246	1.0013	9,783	9,774	1.0009
Other disorders of circulatory system (I80–I99)	3,981	4,101	0.9707	3,941	3,995	0.9865	4,777	4,813	0.9925
Influenza and pneumonia (J09–J18) ²	52,847	52,717	1.0025	56,247	56,326	0.9986	62,804	63,001	0.9969
Influenza (J09–J11) ²	457	411	1.1119	860	849	1.0130	1,806	1,812	0.9967
Pneumonia (J12–J18)	52,389	52,306	1.0016	55,387	55,477	0.9984	60,998	61,189	0.9969
Other acute lower respiratory infections (J20–J22,U04) ³	268	255	1.0510	289	297	0.9731	403	404	0.9975
Acute bronchitis and bronchiolitis (J20–J21)	225	213	1.0563	203	214	0.9486	281	283	0.9929
Other and unspecified acute lower respiratory infections (J22,U04) ^{3,4}	43	42	1.0238	86	83	1.0361	121	121	1.0000
Chronic lower respiratory diseases (J40–J47)	129,311	127,924	1.0108	124,614	124,583	1.0002	130,957	130,933	1.0002
Bronchitis, chronic and unspecified (J40–J42)	704	667	1.0555	740	740	1.0000	860	866	0.9931
Emphysema (J43)	12,963	12,790	1.0135	12,570	12,551	1.0015	13,982	14,002	0.9986
Asthma (J45–J46)	3,355	3,447	0.9733	3,563	3,613	0.9862	3,857	3,884	0.9930

See footnotes at end of table.

Table II. Ratios of preliminary to final reported numbers of deaths from 113 selected causes: United States, 2005–2007—Con.

[For explanation of asterisk preceding cause-of-death codes, see "Technical Notes"]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Preliminary number of deaths 2007	Final number of deaths 2007	Ratio of preliminary to final 2007	Preliminary number of deaths 2006	Final number of deaths 2006	Ratio of preliminary to final 2006	Preliminary number of deaths 2005	Final number of deaths 2005	Ratio of preliminary to final 2005
Other chronic lower respiratory diseases (J44,J47)	112,289	111,020	1.0114	107,741	107,679	1.0006	112,259	112,181	1.0007
Pneumoconioses and chemical effects (J60–J66,J68)	907	915	0.9913	923	924	0.9989	999	1,007	0.9921
Pneumonitis due to solids and liquids (J69)	17,302	16,988	1.0185	16,961	16,887	1.0044	17,351	17,279	1.0042
Other diseases of respiratory system (J00–J06,J30–J39,J67,J70–J98)	28,773	28,508	1.0093	27,676	27,644	1.0012	27,065	27,056	1.0003
Peptic ulcer (K25–K28)	3,000	3,045	0.9852	3,286	3,323	0.9889	3,453	3,478	0.9928
Diseases of appendix (K35–K38)	413	426	0.9695	429	424	1.0118	434	439	0.9886
Hernia (K40–K46)	1,663	1,698	0.9794	1,738	1,744	0.9966	1,629	1,639	0.9939
Chronic liver disease and cirrhosis (K70,K73–K74)	28,504	29,165	0.9773	27,299	27,555	0.9907	27,393	27,530	0.9950
Alcoholic liver disease (K70)	13,891	14,406	0.9643	12,925	13,050	0.9904	12,865	12,928	0.9951
Other chronic liver disease and cirrhosis (K73–K74)	14,613	14,759	0.9901	14,374	14,505	0.9910	14,529	14,602	0.9950
Cholelithiasis and other disorders of gallbladder (K80–K82)	3,178	3,237	0.9818	3,123	3,114	1.0029	3,062	3,072	0.9967
Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	46,095	46,448	0.9924	44,791	45,344	0.9878	43,679	43,901	0.9949
Acute and rapidly progressive nephritic and nephrotic syndrome (N00–N01,N04)	191	206	0.9272	135	138	0.9783	137	137	1.0000
Chronic glomerulonephritis, nephrosis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02–N03,N05–N07,N26)	2,821	2,958	0.9537	1,365	1,841	0.7414	588	867	0.6782
Renal failure (N17–N19)	43,064	43,263	0.9954	43,270	43,344	0.9983	42,925	42,868	1.0013
Other disorders of kidney (N25,N27)	19	21	0.9048	20	21	0.9524	29	29	1.0000
Infections of kidney (N10–N12,N13.6,N15.1)	612	628	0.9745	661	673	0.9822	767	767	1.0000
Hyperplasia of prostate (N40)	498	491	1.0143	518	514	1.0078	527	525	1.0038
Inflammatory diseases of female pelvic organs (N70–N76)	100	116	0.8621	113	112	1.0089	120	120	1.0000
Pregnancy, childbirth and the puerperium (O00–O99)	762	769	0.9909	787	760	1.0355	678	760	0.8921
Pregnancy with abortive outcome (O00–O07)	28	31	0.9032	21	26	0.8077	32	33	0.9697
Other complications of pregnancy, childbirth and the puerperium (O10–O99)	734	738	0.9946	765	734	1.0422	646	727	0.8886
Certain conditions originating in the perinatal period (P00–P96)	14,293	14,599	0.9790	14,384	14,442	0.9960	14,539	14,549	0.9993
Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	10,277	10,421	0.9862	10,434	10,489	0.9948	10,349	10,410	0.9941
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99)	49,960	33,500	1.4913	40,759	31,725	1.2848	40,962	31,999	1.2801
All other diseases (residual)	237,037	238,192	0.9952	236,614	237,421	0.9966	217,020	217,632	0.9972
Accidents (unintentional injuries) (V01–X59,Y85–Y86)	117,075	123,706	0.9464	117,748	121,599	0.9683	114,876	117,809	0.9751
Transport accidents (V01–V99,Y85)	45,832	46,844	0.9784	47,601	48,412	0.9832	48,140	48,441	0.9938
Motor vehicle accidents (V02–V04,V09.0,V09.2,V12–V14,V19.0–V19.2,V19.4–V19.6,V20–V79,V80.3–V80.5,V81.0–V81.1,V82.0–V82.1,V83–V86,V87.0–V87.8,V88.0–V88.8,V89.0,V89.2)	43,098	43,945	0.9807	44,572	45,316	0.9836	45,053	45,343	0.9936
Other land transport accidents (V01,V05–V06,V09.1,V09.3–V09.9,V10–V11,V15–V18,V19.3,V19.8–V19.9,V80.0–V80.2,V80.6–V80.9,V81.2–V81.9,V82.2–V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)	1,017	1,083	0.9391	1,177	1,181	0.9966	1,251	1,241	1.0081
Water, air and space, and other and unspecified transport accidents and their sequelae (V90–V99,Y85)	1,716	1,816	0.9449	1,852	1,915	0.9671	1,837	1,857	0.9892
Nontransport accidents (W00–X59,Y86)	71,244	76,862	0.9269	70,147	73,187	0.9585	66,736	69,368	0.9621
Falls (W00–W19)	22,736	22,631	1.0046	20,533	20,823	0.9861	19,488	19,656	0.9915
Accidental discharge of firearms (W32–W34)	721	613	1.1762	777	642	1.2103	810	789	1.0266
Accidental drowning and submersion (W65–W74)	3,237	3,443	0.9402	3,483	3,579	0.9732	3,468	3,582	0.9682
Accidental exposure to smoke, fire and flames (X00–X09)	3,276	3,286	0.9970	3,066	3,109	0.9862	3,144	3,197	0.9834
Accidental poisoning and exposure to noxious substances (X40–X49)	24,313	29,846	0.8146	24,702	27,531	0.8972	21,416	23,618	0.9068
Other and unspecified nontransport accidents and their sequelae (W20–W31,W35–W64,W75–W99,X10–X39,X50–X59,Y86)	16,961	17,043	0.9952	17,586	17,503	1.0047	18,409	18,526	0.9937

See footnotes at end of table.

Table II. Ratios of preliminary to final reported numbers of deaths from 113 selected causes: United States, 2005–2007—Con.

[For explanation of asterisk preceding cause-of-death codes, see “Technical Notes”]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Preliminary number of deaths 2007	Final number of deaths 2007	Ratio of preliminary to final 2007	Preliminary number of deaths 2006	Final number of deaths 2006	Ratio of preliminary to final 2006	Preliminary number of deaths 2005	Final number of deaths 2005	Ratio of preliminary to final 2005
Intentional self-harm (suicide) (*U03,X60–X84,Y87.0)	33,185	34,598	0.9592	32,185	33,300	0.9665	31,769	32,637	0.9734
Intentional self-harm (suicide) by discharge of firearms (X72–X74)	17,348	17,352	0.9998	16,650	16,883	0.9862	16,932	17,002	0.9959
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60–X71,X75–X84,Y87.0)	15,837	17,246	0.9183	15,535	16,417	0.9463	14,838	15,635	0.9490
Assault (homicide) (*U01–*U02,X85–Y09,Y87.1)	17,520	18,361	0.9542	18,029	18,573	0.9707	17,694	18,124	0.9763
Assault (homicide) by discharge of firearms (*U01.4,X93–X95)	12,129	12,632	0.9602	12,509	12,791	0.9780	12,121	12,352	0.9813
Assault (homicide) by other and unspecified means and their sequelae (*U01.0–*U01.3,*U01.5–*U01.9,*U02,X85–X92,X96–Y09,Y87.1)	5,391	5,729	0.9410	5,520	5,782	0.9547	5,573	5,772	0.9655
Legal intervention (Y35,Y89.0)	371	412	0.9005	411	434	0.9470	377	414	0.9106
Events of undetermined intent (Y10–Y34,Y87.2,Y89.9)	4,888	5,381	0.9084	4,706	5,131	0.9172	4,423	4,742	0.9327
Discharge of firearms, undetermined intent (Y22–Y24)	256	276	0.9275	214	220	0.9727	215	221	0.9729
Other and unspecified events of undetermined intent and their sequelae (Y10–Y21,Y25–Y34,Y87.2,Y89.9)	4,632	5,105	0.9073	4,492	4,911	0.9147	4,209	4,521	0.9310
Operations of war and their sequelae (Y36,Y89.1)	19	21	0.9048	30	28	1.0714	27	27	1.0000
Complications of medical and surgical care (Y40–Y84,Y88)	2,566	2,597	0.9881	2,492	2,521	0.9885	2,630	2,653	0.9913

– Quantity zero.

. . . Category not applicable.

¹Cause-of-death title was changed in 2006 to reflect the addition of Secondary hypertension (ICD–10 code I15).

²New ICD–10 code J09 (Influenza due to identified avian influenza virus) was added to the category in 2007.

³New ICD–10 code U04 (Severe acute respiratory syndrome [SARS], unspecified) was added to the category in 2007.

⁴Cause-of-death title was changed in 2007 to reflect the addition of SARS (ICD–10 Code U04).

SOURCE: CDC/NCHS, National Vital Statistics System, preliminary and final data.

Table III. Ratios of preliminary to final reported numbers of deaths from 130 selected causes of infant death: United States, 2005–2007

[For explanation of asterisk preceding cause-of-death codes, see "Technical Notes"]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Preliminary number of deaths 2007	Final number of deaths 2007	Ratio of preliminary to final 2007	Preliminary number of deaths 2006	Final number of deaths 2006	Ratio of preliminary to final 2006	Preliminary number of deaths 2005	Final number of deaths 2005	Ratio of preliminary to final 2005
All causes	29,241	29,138	1.0035	28,609	28,527	1.0029	28,534	28,440	1.0033
Certain infectious and parasitic diseases (A00–B99)	484	482	1.0041	500	479	1.0438	528	530	0.9962
Certain intestinal infectious diseases (A00–A08)	20	13	1.5385	15	16	0.9375	8	9	0.8889
Diarrhea and gastroenteritis of infectious origin (A09)	—	—	...	2	1	2.0000	1	1	1.0000
Tuberculosis (A16–A19)	2	2	1.0000	1	—	...	2	2	1.0000
Tetanus (A33,A35)	—	—	...	—	—	...	—	—	...
Diphtheria (A36)	—	—	...	—	—	...	—	—	...
Whooping cough (A37)	8	8	1.0000	8	8	1.0000	28	28	1.0000
Meningococcal infection (A39)	9	11	0.8182	10	11	0.9091	17	17	1.0000
Septicemia (A40–A41)	271	283	0.9576	293	269	1.0892	301	302	0.9967
Congenital syphilis (A50)	4	5	0.8000	—	—	...	—	—	...
Gonococcal infection (A54)	—	—	...	—	—	...	—	—	...
Viral diseases (A80–B34)	112	115	0.9739	116	120	0.9667	113	119	0.9496
Acute poliomyelitis (A80)	—	—	...	—	—	...	—	—	...
Varicella (chickenpox) (B01)	—	—	...	—	—	...	—	—	...
Measles (B05)	—	—	...	—	—	...	—	—	...
Human immunodeficiency virus (HIV) disease (B20–B24)	7	5	1.4000	7	7	1.0000	2	2	1.0000
Mumps (B26)	—	—	...	—	—	...	—	—	...
Other and unspecified viral diseases (A81–B00,B02–B04,B06–B19,B25,B27–B34)	105	110	0.9545	109	113	0.9646	111	117	0.9487
Candidiasis (B37)	15	13	1.1538	15	15	1.0000	21	20	1.0500
Malaria (B50–B54)	—	—	...	—	—	...	—	—	...
Pneumocystosis (B59)	1	1	1.0000	—	1	...	2	2	1.0000
All other and unspecified infectious and parasitic diseases (A20–A32, A38, A42–A49,A51–A53,A55–A79,B35–B36,B38–B49,B55–B58,B60–B99)	42	31	1.3548	38	38	1.0000	34	30	1.1333
Neoplasms (C00–D48)	149	131	1.1374	139	141	0.9858	141	134	1.0522
Malignant neoplasms (C00–C97)	92	72	1.2778	77	76	1.0132	78	75	1.0400
Hodgkin's disease and non-Hodgkin's lymphomas (C81–C85)	6	2	3.0000	1	1	1.0000	1	1	1.0000
Leukemia (C91–C95)	20	21	0.9524	31	31	1.0000	22	22	1.0000
Other and unspecified malignant neoplasms (C00–C80,C88–C90,C96–C97)	66	49	1.3469	45	44	1.0227	54	52	1.0385
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	57	59	0.9661	62	65	0.9538	63	59	1.0678
Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50–D89)	108	116	0.9310	104	102	1.0196	94	94	1.0000
Anemias (D50–D64)	16	17	0.9412	10	11	0.9091	19	19	1.0000
Hemorrhagic conditions and other diseases of blood and blood-forming organs (D65–D76)	72	77	0.9351	63	62	1.0161	60	60	1.0000
Certain disorders involving the immune mechanism (D80–D89)	20	22	0.9091	31	29	1.0690	15	15	1.0000
Endocrine, nutritional and metabolic diseases (E00–E88)	255	252	1.0119	200	207	0.9662	222	226	0.9823
Short stature, not elsewhere classified (E34.3)	3	5	0.6000	8	11	0.7273	8	8	1.0000
Nutritional deficiencies (E40–E64)	4	7	0.5714	7	9	0.7778	5	5	1.0000
Cystic fibrosis (E84)	11	11	1.0000	10	11	0.9091	5	5	1.0000
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86–E87)	62	60	1.0333	56	53	1.0566	65	63	1.0317
All other endocrine, nutritional and metabolic diseases (E00–E32,E34.0–E34.2, E34.4–E34.9,E65–E83,E85,E88)	175	169	1.0355	118	123	0.9593	139	145	0.9586
Diseases of the nervous system (G00–G98)	424	413	1.0266	361	373	0.9678	348	354	0.9831

See footnotes at end of table.

Table III. Ratios of preliminary to final reported numbers of deaths from 130 selected causes of infant death: United States, 2005–2007—Con.

[For explanation of asterisk preceding cause-of-death codes, see "Technical Notes"]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Preliminary number of deaths 2007	Final number of deaths 2007	Ratio of preliminary to final 2007	Preliminary number of deaths 2006	Final number of deaths 2006	Ratio of preliminary to final 2006	Preliminary number of deaths 2005	Final number of deaths 2005	Ratio of preliminary to final 2005
Meningitis (G00,G03)	88	82	1.0732	57	61	0.9344	56	57	0.9825
Infantile spinal muscular atrophy, type I (Werdnig–Hoffman) (G12.0)	15	13	1.1538	9	8	1.1250	15	15	1.0000
Infantile cerebral palsy (G80)	11	11	1.0000	8	7	1.1429	7	8	0.8750
Anoxic brain damage, not elsewhere classified (G93.1)	53	64	0.8281	55	60	0.9167	40	42	0.9524
Other diseases of nervous system (G04,G06–G11,G12.1–G12.9,G20–G72,G81–G92,G93.0,G93.2–G93.9,G95–G98)	257	243	1.0576	232	237	0.9789	229	232	0.9871
Diseases of the ear and mastoid process (H60–H93)	3	3	1.0000	2	3	0.6667	7	7	1.0000
Diseases of the circulatory system (I00–I99)	612	624	0.9808	539	543	0.9926	523	529	0.9887
Pulmonary heart disease and diseases of pulmonary circulation (I26–I28)	96	100	0.9600	81	81	1.0000	98	98	1.0000
Pericarditis, endocarditis and myocarditis (I30,I33,I40)	17	21	0.8095	7	9	0.7778	13	13	1.0000
Cardiomyopathy (I42)	117	120	0.9750	109	115	0.9478	82	82	1.0000
Cardiac arrest (I46)	34	29	1.1724	14	15	0.9333	23	24	0.9583
Cerebrovascular diseases (I60–I69)	136	132	1.0303	145	142	1.0211	127	126	1.0079
All other diseases of circulatory system (I00–I25,I31,I34–I38,I44–I45,I47–I51,I70–I99)	212	222	0.9550	181	181	1.0000	181	186	0.9731
Diseases of the respiratory system (J00–J98,U04) ¹	641	640	1.0016	644	692	0.9306	650	669	0.9716
Acute upper respiratory infections (J00–J06)	15	14	1.0714	11	11	1.0000	10	10	1.0000
Influenza and pneumonia (J09–J18) ²	218	222	0.9820	244	263	0.9278	255	265	0.9623
Influenza (J09–J11) ²	9	13	0.6923	17	18	0.9444	17	19	0.8947
Pneumonia (J12–J18)	209	209	1.0000	228	245	0.9306	238	246	0.9675
Acute bronchitis and acute bronchiolitis (J20–J21)	44	45	0.9778	45	52	0.8654	48	50	0.9600
Bronchitis, chronic and unspecified (J40–J42)	23	24	0.9583	19	19	1.0000	22	25	0.8800
Asthma (J45–J46)	5	4	1.2500	5	6	0.8333	3	4	0.7500
Pneumonitis due to solids and liquids (J69)	11	10	1.1000	9	11	0.8182	17	17	1.0000
Other and unspecified diseases of respiratory system (J22,J30–J39,J43–J44,J47–J68,J70–J98,U04) ¹	324	321	1.0093	311	330	0.9424	294	298	0.9866
Diseases of the digestive system (K00–K92)	650	677	0.9601	588	582	1.0103	630	626	1.0064
Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50–K55)	394	413	0.9540	326	323	1.0093	344	341	1.0088
Hernia of abdominal cavity and intestinal obstruction without hernia (K40–K46,K56)	68	68	1.0000	70	67	1.0448	76	77	0.9870
All other and unspecified diseases of digestive system (K00–K28,K30–K38,K57–K92)	189	196	0.9643	192	192	1.0000	209	208	1.0048
Diseases of the genitourinary system (N00–N98)	169	169	1.0000	181	180	1.0056	181	180	1.0056
Renal failure and other disorders of kidney (N17–N19,N25,N27)	137	138	0.9928	159	154	1.0325	152	151	1.0066
Other and unspecified diseases of genitourinary system (N00–N15,N20–N23,N26,N28–N98)	32	31	1.0323	22	26	0.8462	29	29	1.0000
Certain conditions originating in the perinatal period (P00–P96)	14,141	14,466	0.9775	14,223	14,321	0.9932	14,405	14,423	0.9988
Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00–P04)	3,280	3,274	1.0018	3,125	3,150	0.9921	3,230	3,228	1.0006
Newborn affected by maternal hypertensive disorders (P00.0)	96	89	1.0787	85	84	1.0119	89	88	1.0114
Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1–P00.9)	99	91	1.0879	68	80	0.8500	69	70	0.9857
Newborn affected by maternal complications of pregnancy (P01)	1,770	1,769	1.0006	1,694	1,683	1.0065	1,786	1,776	1.0056
Newborn affected by incompetent cervix (P01.0)	505	488	1.0348	444	444	1.0000	498	496	1.0040
Newborn affected by premature rupture of membranes (P01.1)	852	851	1.0012	830	824	1.0073	842	837	1.0060
Newborn affected by multiple pregnancy (P01.5)	222	238	0.9328	213	214	0.9953	256	255	1.0039
Newborn affected by other maternal complications of pregnancy (P01.2–P01.4,P01.6–P01.9)	191	192	0.9948	207	201	1.0299	189	188	1.0053

See footnotes at end of table.

Table III. Ratios of preliminary to final reported numbers of deaths from 130 selected causes of infant death: United States, 2005–2007—Con.

[For explanation of asterisk preceding cause-of-death codes, see "Technical Notes"]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Preliminary number of deaths 2007	Final number of deaths 2007	Ratio of preliminary to final 2007	Preliminary number of deaths 2006	Final number of deaths 2006	Ratio of preliminary to final 2006	Preliminary number of deaths 2005	Final number of deaths 2005	Ratio of preliminary to final 2005
Newborn affected by complications of placenta, cord and membranes (P02)	1,139	1,135	1.0035	1,123	1,140	0.9851	1,111	1,110	1.0009
Newborn affected by complications involving placenta (P02.0–P02.3)	586	579	1.0121	561	563	0.9964	586	585	1.0017
Newborn affected by complications involving cord (P02.4–P02.6)	46	43	1.0698	54	54	1.0000	50	50	1.0000
Newborn affected by chorioamnionitis (P02.7)	503	511	0.9843	507	522	0.9713	471	471	1.0000
Newborn affected by other and unspecified abnormalities of membranes (P02.8–P02.9)	4	2	2.0000	1	1	1.0000	4	4	1.0000
Newborn affected by other complications of labor and delivery (P03)	120	127	0.9449	97	102	0.9510	131	134	0.9776
Newborn affected by noxious influences transmitted via placenta or breast milk (P04)	55	63	0.8730	58	61	0.9508	44	50	0.8800
Disorders related to length of gestation and fetal malnutrition (P05–P08)	4,782	4,961	0.9639	4,943	4,940	1.0006	4,793	4,798	0.9990
Slow fetal growth and fetal malnutrition (P05)	105	104	1.0096	102	99	1.0303	84	83	1.0120
Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	4,678	4,857	0.9631	4,841	4,841	1.0000	4,709	4,714	0.9989
Extremely low birth weight or extreme immaturity (P07.0,P07.2)	3,573	3,706	0.9641	3,678	3,683	0.9986	3,644	3,645	0.9997
Other low birth weight or preterm (P07.1,P07.3)	1,104	1,151	0.9592	1,163	1,158	1.0043	1,064	1,069	0.9953
Disorders related to long gestation and high birth weight (P08)	–	–	–	–	–	–	1	1	–
Birth trauma (P10–P15)	14	12	1.1667	23	22	1.0455	25	26	0.9615
Intrauterine hypoxia and birth asphyxia (P20–P21)	349	356	0.9803	344	344	1.0000	522	529	0.9868
Intrauterine hypoxia (P20)	108	106	1.0189	109	109	1.0000	116	119	0.9748
Birth asphyxia (P21)	241	250	0.9640	236	235	1.0043	406	410	0.9902
Respiratory distress of newborn (P22)	735	789	0.9316	801	825	0.9709	861	860	1.0012
Other respiratory conditions originating in the perinatal period (P23–P28)	1,077	1,117	0.9642	1,199	1,207	0.9934	1,163	1,160	1.0026
Congenital pneumonia (P23)	102	103	0.9903	89	104	0.8558	103	104	0.9904
Neonatal aspiration syndromes (P24)	50	51	0.9804	49	50	0.9800	45	46	0.9783
Interstitial emphysema and related conditions originating in the perinatal period (P25)	131	124	1.0565	155	150	1.0333	122	121	1.0083
Pulmonary hemorrhage originating in the perinatal period (P26)	154	161	0.9565	179	178	1.0056	183	181	1.0110
Chronic respiratory disease originating in the perinatal period (P27)	225	243	0.9259	262	258	1.0155	271	270	1.0037
Atelectasis (P28.0–P28.1)	354	366	0.9672	396	398	0.9950	379	377	1.0053
All other respiratory conditions originating in the perinatal period (P28.2–P28.9)	61	69	0.8841	69	69	1.0000	61	61	1.0000
Infections specific to the perinatal period (P35–P39)	1,039	1,057	0.9830	979	998	0.9810	1,033	1,039	0.9942
Bacterial sepsis of newborn (P36)	790	820	0.9634	786	807	0.9740	834	834	1.0000
Omphalitis of newborn with or without mild hemorrhage (P38)	5	4	1.2500	–	–	–	6	6	1.0000
All other infections specific to the perinatal period (P35,P37,P39)	244	233	1.0472	193	191	1.0105	193	199	0.9698
Hemorrhagic and hematological disorders of newborn (P50–P61)	723	711	1.0169	708	725	0.9766	782	782	1.0000
Neonatal hemorrhage (P50–P52,P54)	614	597	1.0285	598	618	0.9676	664	665	0.9985
Hemorrhagic disease of newborn (P53)	–	–	–	1	1	–	–	–	–
Hemolytic disease of newborn due to isoimmunization and other perinatal jaundice (P55–P59)	20	15	1.3333	10	13	0.7692	16	16	1.0000
Hematological disorders (P60–P61)	89	99	0.8990	99	93	1.0645	102	101	1.0099
Syndrome of infant of a diabetic mother and neonatal diabetes mellitus (P70.0–P70.2)	12	14	0.8571	16	12	1.3333	19	19	1.0000
Necrotizing enterocolitis of newborn (P77)	529	554	0.9549	528	530	0.9962	549	546	1.0055
Hydrops fetalis not due to hemolytic disease (P83.2)	195	177	1.1017	171	168	1.0179	168	165	1.0182
Other perinatal conditions (P29,P70.3–P76,P78–P81,P83.0–P83.1,P83.3–P96)	1,405	1,444	0.9730	1,385	1,400	0.9893	1,258	1,271	0.9898
Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	5,769	5,785	0.9972	5,827	5,819	1.0014	5,562	5,552	1.0018
Anencephaly and similar malformations (Q00)	306	321	0.9533	332	336	0.9881	313	313	1.0000

See footnotes at end of table.

Table III. Ratios of preliminary to final reported numbers of deaths from 130 selected causes of infant death: United States, 2005–2007—Con.

[For explanation of asterisk preceding cause-of-death codes, see "Technical Notes"]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Preliminary number of deaths 2007	Final number of deaths 2007	Ratio of preliminary to final 2007	Preliminary number of deaths 2006	Final number of deaths 2006	Ratio of preliminary to final 2006	Preliminary number of deaths 2005	Final number of deaths 2005	Ratio of preliminary to final 2005
Congenital hydrocephalus (Q03)	92	93	0.9892	82	88	0.9318	92	90	1.0222
Spina bifida (Q05)	21	19	1.1053	24	23	1.0435	22	22	1.0000
Other congenital malformations of nervous system (Q01–Q02,Q04,Q06–Q07)	408	393	1.0382	389	390	0.9974	318	314	1.0127
Congenital malformations of heart (Q20–Q24)	1,345	1,363	0.9868	1,399	1,396	1.0021	1,378	1,377	1.0007
Other congenital malformations of circulatory system (Q25–Q28)	256	254	1.0079	232	236	0.9831	246	245	1.0041
Congenital malformations of respiratory system (Q30–Q34)	393	410	0.9585	449	437	1.0275	599	598	1.0017
Congenital malformations of digestive system (Q35–Q45)	129	132	0.9773	109	108	1.0093	103	105	0.9810
Congenital malformations of genitourinary system (Q50–Q64)	495	514	0.9630	523	518	1.0097	377	375	1.0053
Congenital malformations and deformations of musculoskeletal system, limbs and integument (Q65–Q85)	608	623	0.9759	627	619	1.0129	552	558	0.9892
Down's syndrome (Q90)	78	82	0.9512	93	97	0.9588	123	123	1.0000
Edward's syndrome (Q91.0–Q91.3)	547	525	1.0419	518	509	1.0177	413	405	1.0198
Patau's syndrome (Q91.4–Q91.7)	302	295	1.0237	327	322	1.0155	314	310	1.0129
Other congenital malformations and deformations (Q10–Q18,Q86–Q89)	575	552	1.0417	523	538	0.9721	519	526	0.9867
Other chromosomal abnormalities, not elsewhere classified (Q92–Q99)	216	209	1.0335	200	202	0.9901	190	191	0.9948
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99)	4,162	3,617	1.1507	3,749	3,462	1.0829	3,760	3,589	1.0476
Sudden infant death syndrome (R95)	2,118	2,453	0.8634	2,145	2,323	0.9234	2,107	2,230	0.9448
Other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R53,R55–R94,R96–R99)	2,044	1,164	1.7560	1,604	1,139	1.4083	1,653	1,359	1.2163
All other diseases (residual)	27	16	1.6875	42	25	1.6800	19	15	1.2667
External causes of mortality (*U01,V01–Y84)	1,646	1,747	0.9422	1,510	1,598	0.9449	1,463	1,512	0.9676
Accidents (unintentional injuries) (V01–X59)	1,238	1,285	0.9634	1,119	1,147	0.9756	1,069	1,083	0.9871
Transport accidents (V01–V99)	136	127	1.0709	136	142	0.9577	148	147	1.0068
Motor vehicle accidents (V02–V04,V09.0,V09.2,V12–V14,V19.0–V19.2,V19.4–V19.6,V20–V79,V80.3–V80.5,V81.0–V81.1,V82.0–V82.1,V83–V86,V87.0–V87.8,V88.0–V88.8,V89.0,V89.2)	133	124	1.0726	133	140	0.9500	147	146	1.0068
Other and unspecified transport accidents (V01,V05–V06,V09.1,V09.3–V09.9,V10–V11,V15–V18,V19.3,V19.8–V19.9,V80.0–V80.2,V80.6–V80.9,V81.2–V81.9,V82.2–V82.9,V87.9,V88.9,V89.1,V89.3,V89.9,V90–V99)	4	3	1.3333	2	2	1.0000	1	1	1.0000
Falls (W00–W19)	33	24	1.3750	22	23	0.9565	15	16	0.9375
Accidental discharge of firearms (W32–W34)	2	1	2.0000	–	–	–	1	1	1.0000
Accidental drowning and submersion (W65–W74)	54	57	0.9474	51	51	1.0000	64	64	1.0000
Accidental suffocation and strangulation in bed (W75)	628	669	0.9387	559	588	0.9507	504	514	0.9805
Other accidental suffocation and strangulation (W76–W77,W81–W84)	203	220	0.9227	193	193	1.0000	181	186	0.9731
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78–W80)	62	70	0.8857	64	62	1.0323	45	48	0.9375
Accidents caused by exposure to smoke, fire and flames (X00–X09)	38	38	1.0000	27	27	1.0000	34	34	1.0000
Accidental poisoning and exposure to noxious substances (X40–X49)	16	19	0.8421	15	16	0.9375	20	20	1.0000
Other and unspecified accidents (W20–W31,W35–W64,W85–W99,X10–X39,X50–X59)	66	60	1.1000	51	45	1.1333	56	53	1.0566

See footnotes at end of table.

Table III. Ratios of preliminary to final reported numbers of deaths from 130 selected causes of infant death: United States, 2005–2007—Con.

[For explanation of asterisk preceding cause-of-death codes, see "Technical Notes"]

Cause of death (based on <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>)	Preliminary number of deaths 2007	Final number of deaths 2007	Ratio of preliminary to final 2007	Preliminary number of deaths 2006	Final number of deaths 2006	Ratio of preliminary to final 2006	Preliminary number of deaths 2005	Final number of deaths 2005	Ratio of preliminary to final 2005
Assault (homicide)(*U01,X85–Y09)	322	352	0.9148	292	336	0.8690	281	306	0.9183
Assault (homicide) by hanging, strangulation and suffocation (X91)	32	30	1.0667	23	34	0.6765	24	27	0.8889
Assault (homicide) by discharge of firearms(*U01.4,X93–X95)	13	15	0.8667	6	6	1.0000	7	6	1.1667
Neglect, abandonment and other maltreatment syndromes (Y06–Y07)	74	86	0.8605	67	75	0.8933	93	99	0.9394
Assault (homicide) by other and unspecified means(*U01.0–*U01.3, *U01.5–*U01.9,X85–X90,X92,X96–X99,Y00–Y05,Y08–Y09)	203	221	0.9186	195	221	0.8824	157	174	0.9023
Complications of medical and surgical care (Y40–Y84)	24	22	1.0909	21	23	0.9130	20	19	1.0526
Other external causes and their sequelae (Y10–Y36)	62	88	0.7045	78	92	0.8478	94	104	0.9038

– Quantity zero.

. . . Category not applicable.

¹New ICD–10 code U04 (Severe acute respiratory syndrome [SARS], unspecified) was added to the category in 2007.

²New ICD–10 code J09 (Influenza due to identified avian influenza virus) was added to the category in 2007.

SOURCE: CDC/NCHS, National Vital Statistics System, preliminary and final data.

Infant mortality

The infant mortality rate is the most commonly used index for measuring the risk of dying during the first year of life. The rates presented in this report are calculated by dividing the preliminary number of infant deaths that occurred during 2008 by the number of live births for the same period and are presented as rates per 1,000 or per 100,000 live births. For preliminary birth figures used in the denominator for infant mortality rates, see “Births: Preliminary Data for 2008” (30). In contrast to infant mortality rates based on live births, infant death rates are based on the estimated population under age 1 year (Table 1). Infant death rates that appear in tabulations of age-specific death rates in this report are calculated by dividing the number of infant deaths in 2008 by the estimated population of persons under age 1 on July 1, 2008, and are presented as rates per 100,000 population in this age group. Because of differences in their denominators, infant death rates may differ from infant mortality rates. Information on infant deaths can also be obtained from a file where the infant’s death certificate is linked to the birth certificate. The linked birth/infant death data set (linked file) is a better source of data for infant deaths and mortality rates by race and ethnicity because the race of the mother as reported by the mother on the birth certificate is used in both the numerator and denominator of the infant mortality rate. In contrast, for infant deaths and mortality rates in this report, race information for the denominator is the race of the mother as reported on the birth certificate, but the race information for the numerator is the race of the infant decedent as reported on the death certificate. Race information reported on the birth certificate is considered to be more accurate than that on the death certificate, because the race of each parent is usually reported on the birth certificate by the mother at the time of delivery, whereas on the death certificate, the race of the deceased infant is reported by the funeral director based on information provided by an informant or by observation. This difference in the method of reporting race data has a larger impact for races other than white and black and can lead to differences in race-specific infant mortality rates between the two data sources (29).

Life tables

The period life table provides a measure of the effect of current mortality on life expectancy. It is composed of sets of values showing the mortality experience of a hypothetical group of infants born at the same time and subject throughout their lifetime to the age-specific death rates of a particular time period, usually a given year. Beginning with final data reported for 1997, the life table methodology was changed from previous annual reports. Previously, U.S. life tables were abridged and constructed by reference to a standard table (41). In addition, the age range for these life tables was limited to 5-year age groups ending with the age group 85 years and over.

For 1997–1999 mortality data, a revised life table methodology was used to construct complete life tables by single years of age that extend to age 100 (42) using a methodology similar to that of the decennial life tables (43). The advantages of the new methodology over the previous methodology are its comparability with decennial life table methodology, greater accuracy, and greater age detail. A comparison of the two methods shows small differences in resulting values for life expectancy (42). Although the new method produces complete life

tables, that is, life tables by single years of age, life table data shown in this report are summarized in 5-year age groupings. To calculate the probability of dying at each age, the revised methodology uses vital statistics death rates for ages under 85 and mortality data from the Medicare program for ages 85 and over. Medicare data were used to model the probability of dying at ages 85 and over because the data on decedents’ age are shown to be significantly more reliable than vital statistics data at the oldest ages (44).

The methodology for constructing life tables was revised once more starting with mortality data for the year 2000. Life table data shown in this report for 2007–2008 (Table 6) are based on this revised methodology. Complete life tables by single years of age that extend to age 100 were constructed using a methodology similar to that developed for the 1999–2001 decennial life tables (45). To calculate the probability of dying at each age, two major changes were made to the methodology: 1) Probabilities for ages 66–100 were based on blended vital statistics and Medicare probabilities of dying (probabilities of death for ages 65 and under were based on vital statistics data as before); and 2) smoothing and extrapolation of death rates for ages 66–100 was performed using a mathematical model (45). In the previous method, Medicare probabilities were modeled for ages 85 and over; no blending or smoothing was done. The newly revised methodology, along with a more comprehensive description of the methodology, was published in *United State Life Tables, 2005* (46).

Population denominators

The rates in this report use population estimates based on the 2000 census and are estimated as of July 1, 2007, and July 1, 2008. These population estimates are available on the NCHS website (14,47).

The population estimates have been produced under a collaborative arrangement with the U.S. Census Bureau and are based on the 2000 census counts. Reflecting the new standards issued in 1997 by the OMB, the 2000 census included an option for persons to report more than one race as appropriate for themselves and household members (21). In addition, the 1997 OMB standards called for reporting of Asian persons separately from NHOPi. In the 1977 OMB standards, data for API persons were collected as a single group (17). Death certificates for 16 states currently collect only one race in the same categories as specified in the 1977 OMB standards (see “2003 revision of U.S. Standard Certificate of Death”). In addition, those death certificate data do not report Asians separately from NHOPi. Thus, for nearly one-third of the states, the death certificate data by race (numerators for death rates) are incompatible with population data collected in the 2000 census (the denominators for the rates).

In order to produce national death rates for 2007 and 2008, the reported population data for multiple-race persons had to be “bridged” back to single race categories. In addition, the census counts were modified to be consistent with the 1977 OMB racial categories; that is, to report the data for Asian persons and NHOPi as one combined category, API, and to reflect age as of the census reference date. The procedures used to produce the bridged populations are described in separate publications (19,20). Bridged data are anticipated to be used over the next few years for computing population-based rates. As more states collect data on race according to the 1997 OMB standards (21), use of the bridged populations is expected to be discontinued.

Computing rates and percentages

Death rates are on an annual basis per 100,000 estimated population residing in the specified area. Infant mortality rates are per 1,000 or per 100,000 live births.

Age-adjusted death rates (R') are used to compare relative mortality risks among groups and over time. However, they should be viewed as relative indexes rather than as actual measures of mortality risk. They were computed by the direct method; that is, by applying age-specific death rates (R_i) to the U.S. standard population (relative age distribution of year 2000 projected population of the United States); see the following formula for age-adjusted death rate, and the table of U.S. standard population (Table IV):

$$R' = \sum_i \frac{P_{si}}{P_s} R_i$$

where

P_{si} = standard population for age group i

P_s = total U.S. standard population [all ages combined (Table IV)]

Age-adjusted death rates for injury at work were computed by applying the age-specific death rates to the U.S. standard population for ages 15 and over. The year 2000 standard population used for computing age-adjusted rates and standard errors for injury at work is shown in Table V.

Age-adjusted rates for Puerto Rico, Virgin Islands, Guam, American Samoa, and Northern Marianas were computed by applying age-specific death rates to the U.S. standard population. Age groups for 75 years and over were combined because population counts were unavailable by age group for ages over 79. The year 2000 standard

population used for computing age-adjusted rates and standard errors for the territories is shown in Table VI.

Effective with 1999 data, the standard population was changed from 1940 to the year 2000 population in accordance with the new statistical policy promulgated by the Secretary of Health and Human Services in August 1998 (48). The new population standard affects levels of mortality and to some extent trends and group comparisons. Of particular note are the effects on race comparison of mortality; see "Age Standardization of Death Rates: Implementation of the Year 2000 Standard" (48). Beginning with 2003 data, the traditional standard million population along with corresponding standard weights to six decimal places were replaced by the projected year 2000 population age distribution. The effect of the change is negligible and does not significantly affect comparability with age-adjusted rates calculated using the previous method.

Death rates for the Hispanic population are based only on events to persons reported as Hispanic. Rates for non-Hispanic white persons are based on the sum of all events to white decedents reported as non-Hispanic and white decedents with origin not stated. Likewise, rates for non-Hispanic black persons are based on the sum of all events to black decedents reported as non-Hispanic and black decedents with origin not stated. Hispanic origin is not imputed if it is not reported. For calculating death rates, deaths with age not stated are not distributed. The number of deaths with age not stated in 2008 was 166, less than 0.01 percent of all deaths.

For statistics shown on tables throughout this report, an asterisk (*) indicates that the figure does not meet standards of reliability or precision. In this report, two sets of criteria determine whether a figure meets these standards:

- Reporting for any particular variable is at least 80 percent complete. In this report, no data were suppressed based on this criterion.
- A rate or percentage is based on at least 20 deaths. Rates based on fewer than 20 deaths have a relative standard error (RSE) of about 23 percent or more and, therefore, are considered highly variable. For age-adjusted death rates, this criterion is applied to the sum of the age-specific deaths. However, some death rates (based on data files that are less than 100 percent complete and on 20–31 deaths) may have RSEs of 23 percent or more but are still shown instead of asterisks. As a result, caution should be exercised in analyzing rates based on 20–31 events. Additional information on random variation in numbers of events, rates, ratios, and percentages may be found in "Reliability of estimates."

Table IV. U.S. year 2000 standard population

Age	Population
All ages	274,633,642
Under 1 year	3,794,901
1–4 years	15,191,619
5–14 years	39,976,619
15–24 years	38,076,743
25–34 years	37,233,437
35–44 years	44,659,185
45–54 years	37,030,152
55–64 years	23,961,506
65–74 years	18,135,514
75–84 years	12,314,793
85 years and over	4,259,173

Table V. U.S. year 2000 standard population for those aged 15 years and over

Age	Population
15 years and over	215,670,503
15–24 years	38,076,743
25–34 years	37,233,437
35–44 years	44,659,185
45–54 years	37,030,152
55–64 years	23,961,506
65 years and over	34,709,480

Table VI. U.S. year 2000 standard population for the territories

Age	Population
All ages	274,633,642
Under 1 year	3,794,901
1–4 years	15,191,619
5–14 years	39,976,619
15–24 years	38,076,743
25–34 years	37,233,437
35–44 years	44,659,185
45–54 years	37,030,152
55–64 years	23,961,506
65–74 years	18,135,514
75 years and over	16,573,966

Reliability of estimates

Because the preliminary estimates of deaths in this report are based on files that may not be complete, they are subject to sampling variability. This concept is reflected in the fact that record weights are used to adjust record counts to independent control totals. The lack of completeness of the vital statistics files is due to delays in receiving and processing the death records. Although the proportion of records making up the preliminary file does not constitute a veritable random sample, for the sake of convenience the variability associated with this error (sampling variability) is treated as if it were from a random sample.

Even where the number of vital events in this report is 100 percent complete and not subject to sampling variability, it might be affected by random variation. Thus, when the number of events is small and the probability of such an event is small, considerable caution must be observed in interpreting the data. Such infrequent events may be assumed to follow a Poisson probability distribution. The first column of Table VII shows the estimated RSEs of a file that is nearly 100 percent complete.

Data in a file that is less than 100 percent complete are affected by sampling variation as well as by random variation. The estimated RSEs in columns 2–6 of Table VII for various levels of file completeness are measures of the sampling errors and the random errors of the estimates. They do not include nonsampling error.

The estimated RSEs in Table VII were computed using this formula:

$$1. RSE = 100 \sqrt{\frac{1 - (1 - f)(N - X)}{X + fX(N - (1/f))}}$$

where

f = the sampling fraction or the percent of file completeness/100 from Table I. For mortality data based on deaths under 1 year of age, use f for “infant deaths” for either the demographic or medical file as appropriate. For mortality data based on all ages combined or any age group that is 1 year and over, use f for “deaths 1 year of age and over” for either the demographic or medical files as appropriate.

X = the estimated number of infant deaths or deaths.

N = total count of infant deaths or deaths for the United States or any state. (NOTE: The RSEs shown in Table VII are based on $N = 4,000,000$. If N is smaller, the RSEs may be slightly smaller than those shown.)

RSEs may be used to compute 95 percent confidence intervals for the number of events (X), for a rate (R), or for a percentage (P) and to compute statistical tests concerning the equality of two rates (R_1 and R_2) or two percentages (P_1 and P_2).

Table VII. Relative standard errors for preliminary number of deaths, by percent of file completeness

[Relative standard errors are expressed as a percentage of the estimate]

Estimated number of deaths	Percent of file completeness					
	100	95	90	80	70	60
	Relative standard error (percent)					
1	100.0	102.6	105.4	111.8	119.5	129.1
5	44.7	45.9	47.1	50.0	53.5	57.7
10	31.6	32.4	33.3	35.4	37.8	40.8
20	22.4	22.9	23.6	25.0	26.7	28.9
30	18.3	18.7	19.2	20.4	21.8	23.6
40	15.8	16.2	16.7	17.7	18.9	20.4
50	14.1	14.5	14.9	15.8	16.9	18.3
60	12.9	13.2	13.6	14.4	15.4	16.7
70	12.0	12.3	12.6	13.4	14.3	15.4
80	11.2	11.5	11.8	12.5	13.4	14.4
90	10.5	10.8	11.1	11.8	12.6	13.6
100	10.0	10.3	10.5	11.2	12.0	12.9
200	7.1	7.3	7.5	7.9	8.5	9.1
300	5.8	5.9	6.1	6.5	6.9	7.5
400	5.0	5.1	5.3	5.6	6.0	6.5
500	4.5	4.6	4.7	5.0	5.3	5.8
600	4.1	4.2	4.3	4.6	4.9	5.3
700	3.8	3.9	4.0	4.2	4.5	4.9
800	3.5	3.6	3.7	4.0	4.2	4.6
900	3.3	3.4	3.5	3.7	4.0	4.3
1,000	3.2	3.2	3.3	3.5	3.8	4.1
2,000	2.2	2.3	2.4	2.5	2.7	2.9
5,000	1.4	1.5	1.5	1.6	1.7	1.8
10,000	1.0	1.0	1.1	1.1	1.2	1.3
20,000	0.7	0.7	0.7	0.8	0.8	0.9
50,000	0.4	0.5	0.5	0.5	0.5	0.6
100,000	0.3	0.3	0.3	0.4	0.4	0.4
200,000	0.2	0.2	0.2	0.2	0.3	0.3
500,000	0.1	0.1	0.1	0.2	0.2	0.2
1,000,000	0.1	0.1	0.1	0.1	0.1	0.1
2,000,000	0.1	0.1	0.1	0.1	0.1	0.1
4,000,000	0.1	0.1	0.1	0.1	0.1	0.1

For the number of deaths, the 95 percent confidence interval may be computed as:

2. Lower limit: $X_1 - \left(1.96 \cdot X_1 \cdot \frac{RSE(X_1)}{100} \right)$

3. Upper limit: $X_1 + \left(1.96 \cdot X_1 \cdot \frac{RSE(X_1)}{100} \right)$

As a hypothetical example, assume the number of deaths, X_1 , is 70 from a file with 80 percent completeness. Then

Lower limit: $70 - \left(1.96 \cdot 70 \cdot \frac{13.4}{100} \right) = 51.6$

Upper limit: $70 + \left(1.96 \cdot 70 \cdot \frac{13.4}{100} \right) = 88.4$

This means that the chances are 95 times out of 100 that the confidence interval (51.6–88.4) will cover the “true” number of deaths.

For rates based on population estimates in the denominator, the 95 percent confidence interval may be computed as:

4. Lower limit: $R_1 - \left(1.96 \cdot R_1 \cdot \frac{RSE(R_1)}{100} \right)$

5. Upper limit: $R_1 + \left(1.96 \cdot R_1 \cdot \frac{RSE(R_1)}{100} \right)$

As a hypothetical example, assume the death rate, R_1 , is 20.0, which is based on 70 deaths from a file with 80 percent completeness. Then

Lower limit: $20.0 - \left(1.96 \cdot 20.0 \cdot \frac{13.4}{100} \right) = 14.7$

Upper limit: $20.0 + \left(1.96 \cdot 20.0 \cdot \frac{13.4}{100} \right) = 25.3$

This means that the chances are 95 times out of 100 that the confidence interval (14.7–25.3) will cover the “true” rate.

For age-adjusted death rates, R' , the 95 percent confidence interval may be computed as:

6. Lower limit: $R' - \left(1.96 \cdot R' \cdot \frac{RSE(R')}{100} \right)$

7. Upper limit: $R' + \left(1.96 \cdot R' \cdot \frac{RSE(R')}{100} \right)$

where

8. $RSE(R') = 100 \sqrt{\frac{\sum_i \left(w_i^2 R_i^2 \left[\frac{1}{X_i} + \frac{(1-f_i)(N_i - X_i)}{f_i X_i \left(N_i - \frac{1}{f_i} \right)} \right] \right)}{R'}}$

where

i = each age group where $i = 1$ for infant deaths, $i = 2$ for 1–4 years, $i = 3$ for 5–14 years, . . . and $i = 11$ for 85 years and over.

R_i = age-specific rate for the i th age group.

w_i = i th age-specific U.S. standard population such that $\sum w_i = 1.000000$ (see “Computing rates and percentages”).

X_i = the estimated number of deaths for the i th age group.

N_i = total count of deaths from Table I for each i th age group (for infant deaths, use the count of records as shown; for all age groups 1–4 through 85 years and over, use the count of records as shown for deaths 1 year and over).

f_i = percentage of file completeness/100 from Table I (for infant deaths, use the percent completeness for the demographic or medical file as appropriate for deaths under age 1 year; for all age groups 1–4 through 85 and over, use the percent completeness for the demographic or medical file as appropriate for deaths at ages 1 year and over).

For testing the equality of two rates, R_1 and R_2 , the following z-test may be used to define a significance test statistic:

9. $z = \frac{R_1 - R_2}{\sqrt{R_1^2 \left(\frac{RSE(R_1)}{100} \right)^2 + R_2^2 \left(\frac{RSE(R_2)}{100} \right)^2}}$

The two-tailed 0.95 critical value for a z statistic is 1.96. Therefore, if $|z| \geq 1.96$, the difference is significant at the 0.05 level. If $|z| < 1.96$, then the difference would be considered not statistically significant at the 0.05 level.

As a hypothetical example, assume R_1 is the same as the above example for the current 12-month period and that R_2 , 15.0, is based on 50 deaths occurring in the prior 12-month period (which implies that the file is approximately 100 percent complete for R_2). The z-test may be determined as follows:

$z = \frac{20.0 - 15.0}{\sqrt{(20.0)^2 \left(\frac{13.4}{100} \right)^2 + (15.0)^2 \left(\frac{14.1}{100} \right)^2}} = 1.46$

Because $|z| < 1.96$, there is no statistically significant difference between the two rates at the 0.05 level of significance.

For rates or percentages based on denominators having random variation only or random and sampling variation, the RSE must take into account the variation in both the numerator and denominator. For example, for a rate, R_3 , based on the number of live births in the denominator, the RSE is computed as:

10. $RSE(R_3) = 100 \sqrt{\left(\frac{RSE(D)}{100} \right)^2 + \left(\frac{RSE(B)}{100} \right)^2}$

where

$RSE(D)$ = RSE of the number of deaths, D

$RSE(B)$ = RSE of the number of births, B

The 95 percent confidence interval of R_3 may be computed as follows:

$$11. \text{ Lower limit: } R_3 - \left(1.96 \cdot R_3 \cdot \frac{\text{RSE}(R_3)}{100} \right)$$

$$12. \text{ Upper limit: } R_3 + \left(1.96 \cdot R_3 \cdot \frac{\text{RSE}(R_3)}{100} \right)$$

As a hypothetical example, assume the infant mortality rate, R_3 , is 15.0, which is based on 30 infant deaths (D) from a file with 70 percent completeness and 2,000 live births (B) from a file with 80 percent completeness. Then

$$\text{RSE}(R_3) = 100 \sqrt{\left(\frac{21.8}{100} \right)^2 + \left(\frac{2.5}{100} \right)^2} = 21.9$$

$$\text{Lower limit: } 15.0 - \left(1.96 \cdot 15.0 \cdot \frac{21.9}{100} \right) = 8.6$$

$$\text{Upper limit: } 15.0 + \left(1.96 \cdot 15.0 \cdot \frac{21.9}{100} \right) = 21.4$$

This means that the chances are 95 times out of 100 that the confidence interval (8.6–21.4) will cover the “true” rate. The same formulas are applicable to a percentage (P_1) that has variation in both the numerator and denominator. To compare the equality of two infant mortality rates or two percentages that have variation in the numerator and denominator, the above-mentioned z-test may be used.

Contents

Abstract	1
Introduction	1
Data Sources and Methods	1
Results	2
Trends in numbers and rates	2
Causes of death	4
Infant mortality	6
References	7
List of Detailed Tables	8
Technical Notes	34

Acknowledgments

This report was prepared in the Division of Vital Statistics (DVS) under the direction of Charles J. Rothwell, Director, DVS; Robert N. Anderson, Chief, Mortality Statistics Branch (MSB); and Nicholas F. Pace, Chief, Systems, Programming, and Statistical Resources Branch (SPSRB). Elizabeth Arias of MSB provided content related to life expectancy. David W. Justice of the Data Acquisition and Evaluation Branch (DAEB) contributed to the “[Technical Notes](#).” Jordan Sacks, Annie Liu, Candace Cosgrove, Jaleh Mousavi, and John Birken of SPSRB provided computer programming support and produced statistical tables. Jaleh Mousavi of SPSRB prepared the mortality file. Staff of the Mortality Medical Classification Branch processed the cause-of-death data for individual records. Registration Methods staff and staff of DAEB provided consultation to state vital statistics offices regarding collection of the death certificate data on which this report is based. This report was edited and produced by CDC/OSELS/NCHS/OD/Office of Information Services, Information Design and Publishing Staff. Jane Sudol edited the report, typesetting was done by Jacqueline M. Davis, and graphics were produced by Michael W. Jones (contractor).

Copyright information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

Suggested citation

Miniño AM, Xu JQ, Kochanek KD. Deaths: Preliminary data for 2008. National Vital Statistics Reports; vol 59 no 2. Hyattsville, MD: National Center for Health Statistics. 2010.

National Center for Health Statistics

Edward J. Sondik, Ph.D., *Director*
Jennifer H. Madans, Ph.D., *Associate Director for Science*

Division of Vital Statistics

Charles J. Rothwell, M.S., *Director*
