

A Literature Review of Zika Virus

Anna R. Plourde, Evan M. Bloch

Zika virus is a mosquito-borne flavivirus that is the focus of an ongoing pandemic and public health emergency. Previously limited to sporadic cases in Africa and Asia, the emergence of Zika virus in Brazil in 2015 heralded rapid spread throughout the Americas. Although most Zika virus infections are characterized by subclinical or mild influenza-like illness, severe manifestations have been described, including Guillain-Barre syndrome in adults and microcephaly in babies born to infected mothers. Neither an effective treatment nor a vaccine is available for Zika virus; therefore, the public health response primarily focuses on preventing infection, particularly in pregnant women. Despite growing knowledge about this virus, questions remain regarding the virus's vectors and reservoirs, pathogenesis, genetic diversity, and potential synergistic effects of co-infection with other circulating viruses. These questions highlight the need for research to optimize surveillance, patient management, and public health intervention in the current Zika virus epidemic.

Zika virus is a flavivirus that was first isolated in 1947 from a febrile rhesus macaque monkey in the Zika Forest of Uganda and later identified in *Aedes africanus* mosquitoes from the same forest (1). In 1954, the first 3 cases of human infection were reported in Nigeria (2). Serosurveillance studies in humans suggest that Zika virus is widespread throughout Africa, Asia, and Oceania (online Technical Appendix Table 1, <http://wwwnc.cdc.gov/EID/article/22/7/15-1990-Techapp1.pdf>). However, these studies may overestimate the virus's true prevalence, given serologic overlap between Zika virus and related flaviviruses, such as dengue virus (DENV) and West Nile virus (WNV) (3,4).

Historically, symptomatic Zika virus infections were limited to sporadic cases or small clusters of patients (online Technical Appendix Table 2). This pattern changed in 2007, when the first major outbreak of Zika virus infection occurred in Yap (Federated States of Micronesia), where ≈73% of the population were infected and symptomatic disease developed in ≈18% of infected persons (5). Since then, Zika virus infection has spread rapidly. Outbreaks have occurred in French Polynesia (6), Cook Islands (6), Easter Island (7), New Caledonia (8), and, most recently,

the Americas (9), with sporadic exportations to Europe (Figures 1–3; online Technical Appendix Table 2).

Zika virus was first reported in May 2015 in continental South America in Brazil, where ≈440,000–1,300,000 persons have subsequently been infected (as of February 16, 2016). Furthermore, 29 other countries in the Americas have reported autochthonous Zika virus transmission, including Puerto Rico and US Virgin Islands (Figure 3; online Technical Appendix Table 2) (13). Except for 2 sexually acquired cases, Zika virus in the United States, Canada, and Europe has been restricted to travelers from affected areas (Figure 1; online Technical Appendix Table 2); a patient who delivered an infant with microcephaly in Hawaii had spent part of her pregnancy in Brazil (14).

Given the wealth of new information about Zika virus, we conducted a literature review to summarize the published findings. This review contextualizes the ongoing Zika virus epidemic in the Americas and identifies knowledge gaps that must be addressed to combat Zika virus successfully.

The Review

Search Strategy and Selection Criteria

Using the keywords “Zika,” “ZIKV,” “ZIKAV,” and “Zika virus,” we searched Google, PubMed, Web of Science, Scopus, and ProMed Mail. We reviewed all literature published through February 16, 2016, including peer-reviewed journal articles, infectious disease reporting system broadcasts, and public health agency information (e.g., US Centers for Disease Control and Prevention [CDC] and European Centre for Disease Prevention and Control [ECDC]). To ensure the capture of all information, we cross-referenced the bibliographies of reviewed articles. The search included English-language and foreign-language articles, which were computer translated.

Virology and Pathogenesis

Zika virus is a positive-sense single-stranded RNA virus in the family *Flaviviridae*, which includes several other mosquito-borne viruses of clinical importance (e.g., DENV, WNV, and yellow fever virus [YFV]) (15). Its closest relative is Spondweni virus, the only other member of its clade (15,16). The Zika virus genome contains 10,794 nt encoding 3,419 aa (16). Like other flaviviruses, Zika virus is composed of 2 noncoding regions (5' and 3') that flank an open reading frame (16), which encodes a polyprotein

Author affiliations: University of California, San Francisco, California, USA (A.R. Plourde); The Johns Hopkins University, Baltimore, Maryland, USA (E.M. Bloch); Blood Systems Research Institute, San Francisco (E.M. Bloch)

DOI: <http://dx.doi.org/10.3201/eid2207.151990>




Figure 1. Cases of laboratory-confirmed, imported Zika virus infections in the United States, by state, January 1, 2015–February 10, 2016 (10). All cases are imported, with the exception of 2 sexually acquired autochthonous cases (11,12).

cleaved into the capsid, precursor of membrane, envelope, and 7 nonstructural proteins (16).

Phylogenetic analysis shows that Zika virus can be classified into distinct African and Asian lineages; both emerged from East Africa during the late 1800s or early 1900s (17). The Asian lineage originated during the virus's migration from Africa to Southeast Asia, where it was first detected in Malaysia. From there, Zika virus spread to the Pacific Islands, separately to Yap and French Polynesia, and then to New Caledonia, Cook Islands, Easter Island, and the Americas (17).

A study of Zika virus's molecular evolution, based on viral strains collected from 4 countries in West Africa during 1947–2007, identified several sites within the Zika viral genome that were under strong negative selection pressure. This finding suggests frequent purging of deleterious polymorphisms in functionally important genes and the possibility of recombination, which occurs rarely among flaviviruses (18). The implications of this finding require further evaluation with respect to viral spread, zoonotic maintenance, and epidemiologic potential.

After mosquito inoculation of a human host, cellular entry likely resembles that of other flaviviruses, whereby the

virus enters skin cells through cellular receptors, enabling migration to the lymph nodes and bloodstream. Few studies have investigated the pathogenesis of Zika virus infection. One study showed that human skin fibroblasts, keratinocytes, and immature dendritic cells allow entry of Zika virus (19). Several entry and adhesion factors (e.g., AXL receptor tyrosine kinase) facilitate infection, and cellular autophagy, needed for flaviviral replication, enhances Zika virus replication in skin fibroblasts (19). After cellular entry, flaviviruses typically replicate within endoplasmic reticulum-derived vesicles. However, Zika virus antigens were found exclusively in the nuclei of infected cells; this finding suggests a location for replication that differs from that of other flaviviruses and merits further investigation (20).

Transmission

Zika virus, like other flaviviruses, is transmitted by mosquitoes, primarily of the *Aedes* (*Stegomyia*) genus. Several *Aedes* spp. have been implicated, including *Ae. aegypti*, *Ae. africanus*, *Ae. hensilli*, and *Ae. albopictus* (1,21–23). The *Ae. aegypti* mosquito appears to be the major vector in Asia (24) and was the suspected primary vector for the French Polynesia outbreak (25). Zika virus has been detected in wild-caught



Figure 2. All countries and regions reporting laboratory-confirmed autochthonous Zika virus cases, January 1, 2015–February 10, 2016 (online Technical Appendix Table 2, <http://wwwnc.cdc.gov/EID/article/22/7/15-1990-Techapp1.pdf>). Data represent outbreaks and case reports for all reported autochthonous laboratory-confirmed cases of Zika virus infection, including those reported in the peer-reviewed literature; public health agency Web sites, bulletins, and broadcasts; and media reports for selected dates.

Ae. aegypti mosquitoes, which laboratory experiments have shown to be capable of transmitting Zika virus (26,27). *Ae. hensilli* mosquitoes were implicated in the Yap outbreak, yet Zika virus has never been isolated from these mosquitoes (28,29). In Africa, the predominant *Aedes* species vector has not been definitively identified, although viral isolation studies suggest that *Ae. albopictus* was the likely vector in a 2007 Zika virus outbreak in Gabon (23).

Aedes mosquitoes are widely distributed globally, and native habitats of most species are warm tropical and subtropical regions (29–31). Some species show a limited distribution (e.g., *Ae. luteocephalus* in Africa and *Ae. hensilli* in the Pacific Islands); others have a broad geographic span (e.g., *Ae. aegypti* and *Ae. albopictus*) (29–31). *Ae. albopictus* does not yet appear to be a major vector of Zika virus. However, its role in the 2007 Gabon outbreak, its wide distribution throughout the United States, and Zika virus's lack of restriction to a specific *Aedes* sp. indicate that this species could serve as a vector in the United States (9).

Mosquito acquisition of the virus likely occurs during a blood meal; after uptake, the virus replicates and is transmitted to a reservoir animal at the next blood meal

(32). Isolation of the virus or of anti-Zika virus antibodies from various nonhuman primates and other wild and domestic animals suggests multiple animal reservoirs (33). One study examined the kinetics of Zika virus infectivity in *Ae. aegypti* mosquitoes by using blood-feeding membranes (27); viral content was high on the day of feeding (inoculation), decreased to undetectable levels through day 10, increased by day 15, and remained high on days 20–60. These findings suggest an incubation period in mosquitoes of ≈ 10 days.

Other nonvector modes of Zika virus transmission include congenital (34), perinatal (35), and sexual (11,36). Possible transmission by blood transfusion (37,38), animal bite (39), and laboratory exposure (40; online Technical Appendix reference 41) has been described; however, confounding by contemporaneous vectorborne transmission in these instances cannot be excluded. For example, the patient who became infected with Zika virus after a monkey bite had concomitant exposure to mosquitoes, a more plausible route of acquisition (39). Similarly, 1 of 2 patients with potentially laboratory-acquired infection (40; online Technical Appendix reference 41) reported recent



Figure 3. South America, Central America, and Caribbean countries and regions reporting laboratory-confirmed autochthonous Zika virus disease cases during January 1, 2015–February 10, 2016 (online Technical Appendix Table 2, <http://wwwnc.cdc.gov/EID/article/22/7/15-1990-Techapp1.pdf>). Data represent outbreaks and case reports for all reported autochthonous laboratory-confirmed cases of Zika virus infection in these countries and regions during January 1, 2015–February 10, 2016, including those reported in peer-reviewed literature; public health agency Web sites, bulletins, and broadcasts; and media reports.

exposure to mosquitoes (40); no definitive mechanism for transmission was described for either patient.

Intrauterine transmission is supported by the finding of Zika virus RNA by reverse transcription PCR (RT-PCR) in amniotic fluid of 2 mothers with symptoms of Zika virus infection during pregnancy; both delivered babies with microcephaly (34). Zika virus RNA has also been identified in tissue of fetuses from women infected during pregnancy and in brains of 2 live-born infants with microcephaly who died <20 hours after birth (online Technical Appendix references 42–45). Probable intrapartum transmission has also been described: 2 newborns were found to be viremic with Zika virus \leq 4 days after being born to infected mothers (35). Viral RNA, but not culturable virus, has been detected in breast milk (35), but transmission by breast-feeding has not been reported.

Two cases of possible transfusion-transmitted Zika virus were reported in Brazil (38). Furthermore, during the French Polynesia outbreak, a study found that 42 (2.8%) of 1,505 asymptomatic blood donors were positive for Zika virus by RT-PCR; 11 donors described a Zika fever-like syndrome 3–10 days after donation (37).

Clinical Manifestations

In humans, the incubation period from mosquito bite to symptom onset is \approx 3–12 days. Infection is likely asymptomatic in \approx 80% of cases (5,32). All ages are susceptible (4 days–76 years), with a slight preponderance of cases in females (online Technical Appendix Table 3). When symptoms occur, they are typically mild, self-limiting, and non-specific (online Technical Appendix Table 3); similarity to other arbovirus infections (e.g., DENV and chikungunya virus [CHIKV]) may confound the diagnosis (online Technical Appendix reference 46). Commonly reported symptoms include rash, fever, arthralgia, myalgia, fatigue, headache, and conjunctivitis (online Technical Appendix Table 3). Rash, a prominent feature, is maculopapular and pruritic in most cases; it begins proximally and spreads to the extremities with spontaneous resolution within 1–4 days of onset (40). Fever is typically low grade (37.4°C – 38.0°C) (8,36,40). Symptoms resolve within 2 weeks; accounts of longer persistence are rare (25; online Technical Appendix reference 47).

More severe clinical sequelae have increasingly been associated with Zika virus. During the ongoing outbreak

in Brazil, reports of infants born with microcephaly have markedly increased (>3,800 cases; 20 cases/10,000 live births vs. 0.5/10,000 live births in previous years) (online Technical Appendix reference 48). However, concern exists that these findings may in part be artifactual, resulting from previous underreporting of cases and confounding by other risk factors for microcephaly (online Technical Appendix reference 49). Because systematic surveillance for microcephaly was not previously undertaken, the baseline rate of microcephaly in Brazil is unknown, and subsequent reports suggest that a substantial proportion of infants that reportedly have microcephaly do not actually have the condition (online Technical Appendix reference 50).

Health officials in French Polynesia have reported an apparent increase in congenital central nervous system (CNS) malformations, coinciding with the outbreak occurring during 2013–2014 (online Technical Appendix reference 51). However, this finding should be cautiously interpreted; reports included only 17 cases, and none were laboratory-confirmed Zika virus cases. In addition, the true baseline rate of such malformations before the outbreak is unknown (online Technical Appendix reference 51).

A plausible neuropathologic link between Zika virus and CNS anomalies is supported by research showing viral neurotropism in intraperitoneally infected mice (online Technical Appendix reference 52) and progression of disease in directly infected mouse brains (online Technical Appendix reference 53). One hypothesis for Zika virus's role in CNS malformations pertains to the virus's hijacking of autophagy during viral replication (online Technical Appendix reference 54). Some cellular proteins have a dual role in autophagy and centrosome stability; a normal number of centrosomes is important for brain development (online Technical Appendix reference 54). An increase in centrosomes in mice has been shown to result in microcephaly (online Technical Appendix reference 54). Therefore, Zika virus's interference in autophagy has been hypothesized to lead to an increase in centrosome number and microcephaly; this potential role in malformations merits further investigation.

Severe neurologic sequelae have also been described in adults, including meningitis, meningoencephalitis, and Guillain-Barre syndrome (online Technical Appendix reference 55). A surge in Guillain-Barre syndrome cases has been observed in Brazil, Colombia, El Salvador, Suriname, Venezuela, and French Polynesia during outbreaks; however, Zika virus has been laboratory confirmed in only some of these cases (online Technical Appendix reference 55).

Nonneurologic sequelae include transient hearing loss, hypotension, and genitourinary symptoms (11,36; online Technical Appendix references 56,57). Hematospermia was reported in 2 cases (11,36). A 44-year-old man in Tahiti in whom hematospermia developed 2 weeks after symptoms of Zika virus infection was found to have replicative

cultured Zika virus particles in his semen (36). In addition, a 36-year-old man from the United States contracted Zika virus infection while in Senegal, and subsequently, his wife was infected in the United States; this case supports sexual transmission (11). A second sexually acquired case was reported in Texas (online Technical Appendix reference 58).

Rare deaths have been described in patients infected with Zika virus (online Technical Appendix reference 44). Besides 1 infant death, 3 other fatalities were reported (2 from Brazil and 1 from Colombia): 1 man with lupus erythematosus, chronic corticosteroid use, rheumatoid arthritis, and alcoholism; and 2 girls 16 years of age, 1 with sickle cell disease (online Technical Appendix reference 59). (Medical history was not reported for the other girl [online Technical Appendix reference 44].)

General Laboratory Findings

Information on laboratory findings for Zika virus infection is limited. Complete blood count is often normal; even if blood count is abnormal, changes may be nonspecific (e.g., mild lymphopenia, mild neutropenia, mild-to-moderate thrombocytopenia) (8; online Technical Appendix references 46,60–62). Mild elevations in inflammatory markers (C-reactive protein, fibrinogen, and ferritin), serum lactate dehydrogenase, or liver enzymes have been described (8,25; online Technical Appendix reference 57). These findings are observed in many other viral infections, including the co-circulating viruses DENV and CHIKV, so none of these laboratory alterations reliably distinguish among these infections.

Diagnosis

Clinical evaluation alone is unreliable for a diagnosis of Zika virus infection. Because of clinical overlap with other arboviruses, diagnosis relies on laboratory testing. Evaluation for Zika virus, CHIKV, and DENV should be undertaken concurrently for all patients who have acute fever, rash, myalgia, or arthralgia after recent (previous 2 weeks) travel to an area of ongoing Zika virus transmission (online Technical Appendix reference 63). Commercial assays have been developed, including a PCR-based assay that has been approved by the Communauté Européenne (RealStar Zika Virus RT-PCR Kit 1.0, Altona Diagnostics, Hamburg, Germany) and a serologic assay that has been approved by the US Food and Drug Administration for restricted use in emergency situations (online Technical Appendix reference 64). Testing has typically been performed by large reference laboratories (e.g., US CDC and US state laboratories) and universities. CDC's typical turnaround time is 4–14 days. Appropriate tests are selected by the laboratory on the basis of clinical information provided by the requesting healthcare provider (online Technical Appendix reference 65). To coordinate sample collection, providers should contact local public health agencies before testing.

Molecular amplification (e.g., RT-PCR) on serum samples remains the most specific diagnostic approach and is the preferred testing method for Zika virus during the acute phase of illness (<7 days from symptom onset) (online Technical Appendix reference 63). In contrast, serologic testing is not recommended during the acute phase, when Zika virus IgM may be undetectable (22). However, molecular testing must be performed during the viremic period (15). Several case reports of negative RT-PCR results but positive IgM results for patients whose samples were tested at ≥ 5 days after symptom onset indicate a possible viremic period as brief as 5 days (25,36; online Technical Appendix reference 61). Consequently, testing algorithms are based on sampling relative to symptom onset, and serologic testing should be considered if samples are negative for Zika virus by RT-PCR (online Technical Appendix reference 63).

Serologic testing has limitations. Zika virus IgM and IgG are notoriously cross-reactive with those against other flaviviruses (particularly DENV), limiting specificity (5,15; online Technical Appendix reference 46). Therefore, positive serologic test results should be confirmed with testing that uses an alternative platform such as a seroneutralization assay (e.g., plaque-reduction neutralization test) (22). However, flaviviral cross-reactivity can also pose problems in confirmatory assays, especially for patients immunized (e.g., against YFV or Japanese encephalitis virus) or infected with another flavivirus (e.g., WNV or St. Louis encephalitis virus); presence of antibodies confounds diagnosis (online Technical Appendix reference 63).

The type of sample can also affect the probability of detection. Although diagnostic testing is performed primarily on serum or cerebrospinal fluid, the diagnostic utility of other specimen types (e.g., urine, saliva, amniotic fluid, and tissue) is being evaluated (online Technical Appendix reference 63). Urine and saliva may offer alternatives, particularly when blood collection is difficult (e.g., in children or remote locations). Viruria may persist longer than viremia. One study reported that Zika virus RNA was detected in urine up to 20 days after viremia had become undetectable (online Technical Appendix reference 62); therefore, RT-PCR testing of urine should be considered when Zika virus is clinically suspected, despite negative serum testing (22,33,35,36; online Technical Appendix reference 62). Similarly, RT-PCR conducted with saliva has been shown to increase the detection rate during the acute phase of infection but does not extend the window of detection of Zika virus RNA; consequently, blood remains the preferred sample (online Technical Appendix reference 66).

Management and Prevention

No specific treatment or vaccine is available for Zika virus infection. Management is supportive and includes rest, fluids,

antipyretics, and analgesics. Aspirin and other nonsteroidal antiinflammatory drugs should be avoided until dengue is excluded because of the risk for hemorrhage among dengue patients (online Technical Appendix reference 67).

Other general measures focus on prevention of mosquito bites, including individual protection (e.g., long pants, light-colored clothing, insect repellants, bed nets), particularly during known *Ae. aegypti* peak biting times (early morning and late afternoon) (online Technical Appendix reference 68). Community-level strategies target mosquito breeding through elimination of potential egg-laying sites (e.g., potted plant saucers, water storage units, used tires) by drying wet environments or using insecticide treatment (online Technical Appendix reference 68). Pregnant women residing in countries that are not Zika virus–endemic are advised against travel to affected countries (online Technical Appendix reference 69). Testing should be offered to all pregnant women who have traveled to areas with ongoing Zika virus transmission (online Technical Appendix reference 70). Serial fetal ultrasounds should be considered to monitor fetal anatomy and growth every 3–4 weeks in pregnant women with positive or inconclusive Zika virus test results, and the infant should be tested at birth (online Technical Appendix reference 70). Men who reside in or have traveled to an area of active Zika virus transmission and who have a pregnant partner should abstain from sexual activity or use condoms during sex; similar guidelines apply for men with a nonpregnant female sex partner who is concerned about sexual transmission of Zika virus (online Technical Appendix reference 58).

Discussion

Zika virus has been declared a public health emergency. As many as 1.3 million persons have been affected in Brazil alone (online Technical Appendix Table 2), and 20 countries or territories have reported local transmission of the virus during 2016 (Figures 2,3). Because of the ease of air travel and international trade, further spread into regions where the virus is not endemic is likely, and transmission is probable in locations with competent mosquito vectors. A robust, multifaceted response is underway that involves public health authorities, government agencies, the biomedical industry, medical practitioners, and researchers. However, uncertainty remains regarding aspects of the virus's vectors, epidemiology, and pathogenesis. As the epidemic unfolds, evaluating incoming data critically will be necessary to separate fact from speculation.

Foremost, diagnosis remains suboptimal. Diagnostic guidelines are contingent on laboratory testing that is not widely available. Although commercial tests for Zika virus are limited in number and availability, more are in development, including prototype multiplex molecular assays that concurrently test for Zika virus, CHIKV, and DENV

(M.P. Busch, pers. comm.). However, although not unique to Zika virus, laboratory infrastructure and testing capability is lacking in resource-constrained settings where Zika virus is most prevalent.

Prevention measures (specifically, vector control) are a current priority, pending advances in diagnostics; the World Health Organization and the Pan American Health Organization have issued recommendations (online Technical Appendix reference 44). In the United States, multiple factors guard against the explosive epidemic occurring throughout Central and South America. Specifically, lower rates of human crowding in urban areas, wider access to air conditioning and mosquito repellants, and waste management limit mosquito-borne transmission, which has been the case for DENV (online Technical Appendix reference 71). Nonetheless, further entomologic research is needed to define the range of Zika virus vectors and identify new areas where autochthonous transmission could take place to enable early intervention. Investment is also needed in durable control measures such as adaptable vaccine platforms for arboviruses; currently, no Zika virus vaccines are in advanced development (9).

Aspects of Zika virus pathogenesis remain unclear. Zika virus's association with neurologic sequelae, including potential neuropathophysiologic mechanisms, is being actively investigated. Continued epidemiologic study, combined with research involving animal models, will offer increased insight, which could spur novel prevention strategies (9). If confirmed, insights into the timing of infection relative to gestational outcomes will guide policy. In the interim, new cases of Zika virus infection should be monitored for complications, particularly in babies born to mothers residing in Zika virus-affected areas. The effects of Zika virus in other vulnerable clinical subsets (e.g., those who have concurrent conditions or are immunocompromised) also merit further investigation, as does co-infection or sequential infection by co-circulating viruses.

Given reports of possible transfusion-transmitted Zika virus, the pandemic also has implications for the blood supply within Zika virus-endemic and nonendemic regions. The US Food and Drug Administration recommends 28-day deferral for blood donors with confirmed or suspected Zika virus infection (38). Donor screening by nucleic acid testing is being considered but will be challenging to implement because of high costs and regulatory considerations. Pathogen-reduction technology has shown efficacy for treatment of plasma (online Technical Appendix reference 72); however, absence of a licensed pathogen reduction technology for use in red cells, high incremental cost, and technical barriers render such technology an unlikely short-term solution.

Zika virus has the propensity to infect large numbers of persons with severe consequences in some cases.

The epidemic has serious medical, ethical, and economic ramifications, particularly in countries where the resources for early diagnosis are lacking and potential intervention measures (e.g., contraception or termination of pregnancy) are discouraged or illegal (online Technical Appendix reference 73). Although autochthonous transmission in the United States is unlikely to match the scale of the epidemic in Central and South America, much about Zika virus and the way the pandemic will evolve are unknown. Continued vigilance is warranted, along with a concerted effort toward improving our understanding, management, and prevention of this emerging pathogen.

Acknowledgments

We thank Matthew L. Brandwein for his technical expertise in generating the figures.

Dr. Plourde is a resident physician in the Departments of Pathology and Laboratory Medicine at the University of California, San Francisco Medical Center. Her interests include public health microbiology and emerging infectious diseases.

Dr. Bloch is an assistant professor in the Department of Pathology and an associate director of the Transfusion Division at The Johns Hopkins University School of Medicine. His interests include neglected tropical diseases and transfusion-transmissible infections.

References

1. Dick GW, Kitchen SF, Haddow AJ. Zika virus. I. Isolations and serological specificity. *Trans R Soc Trop Med Hyg.* 1952;46:509–20. [http://dx.doi.org/10.1016/0035-9203\(52\)90042-4](http://dx.doi.org/10.1016/0035-9203(52)90042-4)
2. MacNamara FN. Zika virus: a report on three cases of human infection during an epidemic of jaundice in Nigeria. *Trans R Soc Trop Med Hyg.* 1954;48:139–45. [http://dx.doi.org/10.1016/0035-9203\(54\)90006-1](http://dx.doi.org/10.1016/0035-9203(54)90006-1)
3. Korhonen EM, Huhtamo E, Smura T, Kallio-Kokko H, Raassina M, Vapalahti O. Zika virus infection in a traveller returning from the Maldives, June 2015. *Euro Surveill.* 2016; 21:30107. <http://dx.doi.org/10.2807/1560-7917.ES.2016.21.2.30107>
4. Baba SS, Fagbami AH, Ojeh CK. Preliminary studies on the use of solid-phase immunosorbent techniques for the rapid detection of Wesselsbron virus (WSLV) IgM by haemagglutination-inhibition. *Comp Immunol Microbiol Infect Dis.* 1999;22:71–9. [http://dx.doi.org/10.1016/S0147-9571\(98\)00003-4](http://dx.doi.org/10.1016/S0147-9571(98)00003-4)
5. Duffy MR, Chen TH, Hancock WT, Powers AM, Kool JL, Lanciotti RS, et al. Zika virus outbreak on Yap Island, Federated States of Micronesia. *N Engl J Med.* 2009;360:2536–43. <http://dx.doi.org/10.1056/NEJMoa0805715>
6. Roth A, Mercier A, Lepers C, Hoy D, Duituturaga S, Benyon E, et al. Concurrent outbreaks of dengue, chikungunya and Zika virus infections - an unprecedented epidemic wave of mosquito-borne viruses in the Pacific 2012–2014. *Euro Surveill.* 2014;19:20929. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.41.20929>
7. Tognarelli J, Ulloa S, Villagra E, Lagos J, Aguayo C, Fasce R, et al. A report on the outbreak of Zika virus on Easter Island, South Pacific, 2014. *Arch Virol.* 2016;161:665–8. <http://dx.doi.org/10.1007/s00705-015-2695-5>
8. Dupont-Rouzeyrol M, O'Connor O, Calvez E, Daurès M, John M, Grangeon JP, et al. Co-infection with Zika and dengue viruses in 2

- patients, New Caledonia, 2014. *Emerg Infect Dis.* 2015;21:381–2. <http://dx.doi.org/10.3201/eid2102.141553>
9. Fauci AS, Morens DM. Zika virus in the Americas—yet another arbovirus threat. *N Engl J Med.* 2016;374:601–4. <http://dx.doi.org/10.1056/NEJMp1600297>
 10. Centers for Disease Control and Prevention. Zika virus disease in the United States, 2015–2016. 2016 Feb 10 [cited 2016 Feb 17]. <http://www.cdc.gov/zika/geo/united-states.html>
 11. Foy BD, Kobylinski KC, Chilson Foy JL, Blitvich BJ, Travassos da Rosa A, Haddock AD, et al. Probable non-vector-borne transmission of Zika virus, Colorado, USA. *Emerg Infect Dis.* 2011;17:880–2. <http://dx.doi.org/10.3201/eid1705.101939>
 12. Dallas County Health and Human Services. DCHHS reports first Zika virus case in Dallas County acquired through sexual transmission. 2016 Feb 2 [cited 2016 Feb 8]. <http://www.dallascounty.org/department/hhs/press/documents/PR2-2-16DCHHSReportsFirstCaseofZikaVirusThroughSexualTransmission.pdf>
 13. Hennessey M, Fischer M, Staples JE. Zika virus spreads to new areas—region of the Americas, May 2015–January 2016. *MMWR Morb Mortal Wkly Rep.* 2016;65:55–8. <http://dx.doi.org/10.15585/mmwr.mm6503e1>
 14. European Centre for Disease Prevention and Control. Zika virus disease epidemic: potential association with microcephaly and Guillain-Barré syndrome (first update). 2016 Jan 21 [cited 2016 Feb 3]. <http://ecdc.europa.eu/en/publications/Publications/rapid-risk-assessment-zika-virus-first-update-jan-2016.pdf>
 15. Lanciotti RS, Kosoy OL, Laven JJ, Velez JO, Lambert AJ, Johnson AJ, et al. Genetic and serologic properties of Zika virus associated with an epidemic, Yap State, Micronesia, 2007. *Emerg Infect Dis.* 2008;14:1232–9. <http://dx.doi.org/10.3201/eid1408.080287>
 16. Kuno G, Chang GJ. Full-length sequencing and genomic characterization of Bagaza, Kedougou, and Zika viruses. *Arch Virol.* 2007;152:687–96. <http://dx.doi.org/10.1007/s00705-006-0903-z>
 17. Gatherer D, Kohl A. Zika virus: a previously slow pandemic spreads rapidly through the Americas. *J Gen Virol.* 2016;97:269–73. <http://dx.doi.org/10.1099/jgv.0.000381>
 18. Faye O, Freire CC, Iamarino A, Faye O, de Oliveira JV, Diallo M, et al. Molecular evolution of Zika virus during its emergence in the 20(th) century. *PLoS Negl Trop Dis.* 2014;8:e2636. <http://dx.doi.org/10.1371/journal.pntd.0002636>
 19. Hamel R, Dejarnac O, Wicht S, Ekchariyawat P, Neyret A, Luplertlop N, et al. Biology of Zika virus infection in human skin cells. *J Virol.* 2015;89:8880–96. <http://dx.doi.org/10.1128/JVI.00354-15>
 20. Buckley A, Gould EA. Detection of virus-specific antigen in the nuclei or nucleoli of cells infected with Zika or Langkat virus. *J Gen Virol.* 1988;69:1913–20. <http://dx.doi.org/10.1099/0022-1317-69-8-1913>
 21. Marchette NJ, Garcia R, Rudnick A. Isolation of Zika virus from *Aedes aegypti* mosquitoes in Malaysia. *Am J Trop Med Hyg.* 1969;18:411–5.
 22. Hayes EB. Zika virus outside Africa. *Emerg Infect Dis.* 2009;15:1347–50. <http://dx.doi.org/10.3201/eid1509.090442>
 23. Grard G, Caron M, Mombou IM, Nkoghe D, Mboui Ondo S, Jiolle D, et al. Zika virus in Gabon (Central Africa)—2007: a new threat from *Aedes albopictus*? *PLoS Negl Trop Dis.* 2014;8:e2681. <http://dx.doi.org/10.1371/journal.pntd.0002681>
 24. Olson JG, Ksiazek TG, Suhandiman, Triwibowo. Zika virus, a cause of fever in Central Java, Indonesia. *Trans R Soc Trop Med Hyg.* 1981;75:389–93. [http://dx.doi.org/10.1016/0035-9203\(81\)90100-0](http://dx.doi.org/10.1016/0035-9203(81)90100-0)
 25. Oehler E, Watrin L, Larre P, Leparc-Goffart I, Lastere S, Valour F, et al. Zika virus infection complicated by Guillain-Barre syndrome—case report, French Polynesia, December 2013. *Euro Surveill.* 2014;19:20720. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.9.20720>
 26. Diallo D, Sall AA, Diagne CT, Faye O, Faye O, Ba Y, et al. Zika virus emergence in mosquitoes in southeastern Senegal, 2011. *PLoS One.* 2014;9:e109442. <http://dx.doi.org/10.1371/journal.pone.0109442>
 27. Boorman JP, Porterfield JS. A simple technique for infection of mosquitoes with viruses; transmission of Zika virus. *Trans R Soc Trop Med Hyg.* 1956;50:238–42. [http://dx.doi.org/10.1016/0035-9203\(56\)90029-3](http://dx.doi.org/10.1016/0035-9203(56)90029-3)
 28. Musso D, Nilles EJ, Cao-Lorreau VM. Rapid spread of emerging Zika virus in the Pacific area. *Clin Microbiol Infect.* 2014;20:O595–6. <http://dx.doi.org/10.1111/1469-0691.12707>
 29. Ledermann JP, Guillaumot L, Yug L, Saweyog SC, Tided M, Machieng P, et al. *Aedes hensilli* as a potential vector of Chikungunya and Zika viruses. *PLoS Negl Trop Dis.* 2014;8:e3188. <http://dx.doi.org/10.1371/journal.pntd.0003188>
 30. Kraemer MU, Sinka ME, Duda KA, Mylne A, Shearer FM, Brady OJ, et al. The global compendium of *Aedes aegypti* and *Ae. albopictus* occurrence. *Sci Data.* 2015;2:150035. <http://dx.doi.org/10.1038/sdata.2015.35>
 31. Gaffigan TVWR, Pecor JE, Stoffer JA, Anderson T. Systematic catalog of culicidae. Suitland (MD): Smithsonian Institution; Silver Spring (MD): Walter Reed Army Institute of Research; 2016 [cited 2016 Feb 10]. <http://mosquitocatalog.org/>
 32. Ios S, Mallet HP, Leparc Goffart I, Gauthier V, Cardoso T, Herida M. Current Zika virus epidemiology and recent epidemics. *Med Mal Infect.* 2014;44:302–7. <http://dx.doi.org/10.1016/j.medmal.2014.04.008>
 33. Haddock AD, Schuh AJ, Yasuda CY, Kasper MR, Heang V, Huy R, et al. Genetic characterization of Zika virus strains: geographic expansion of the Asian lineage. *PLoS Negl Trop Dis.* 2012;6:e1477. <http://dx.doi.org/10.1371/journal.pntd.0001477>
 34. Oliveira Melo AS, Malinger G, Ximenes R, Szejnfeld PO, Alves Sampaio S, Bispo de Filippis AM. Zika virus intrauterine infection causes fetal brain abnormality and microcephaly: tip of the iceberg? *Ultrasound Obstet Gynecol.* 2016;47:6–7. <http://dx.doi.org/10.1002/uog.1583135>
 - Besnard M, Lastere S, Teissier A, Cao-Lorreau V, Musso D. Evidence of perinatal transmission of Zika virus, French Polynesia, December 2013 and February 2014. *Euro Surveill.* 2014;19:20751. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.13.20751>
 36. Musso D, Roche C, Robin E, Nhan T, Teissier A, Cao-Lorreau VM. Potential sexual transmission of Zika virus. *Emerg Infect Dis.* 2015;21:359–61. <http://dx.doi.org/10.3201/eid2102.141363>
 37. Musso D, Nhan T, Robin E, Roche C, Bierlaire D, Zisou K, et al. Potential for Zika virus transmission through blood transfusion demonstrated during an outbreak in French Polynesia, November 2013 to February 2014. *Euro Surveill.* 2014;19:20761. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.14.20761>
 38. US Food and Drug Administration. Recommendations for donor screening, deferral, and product management to reduce the risk of transfusion transmission of Zika virus, recommendations for industry. 2016 Feb 16 [cited 2016 Feb 16]. <http://www.fda.gov/downloads/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/Blood/UCM486360.pdf>
 39. Leung GH, Baird RW, Druce J, Anstey NM. Zika virus infection in Australia following a monkey bite in Indonesia. *Southeast Asian J Trop Med Public Health.* 2015;46:460–4.
 40. Simpson DI. Zika virus infection in man. *Trans R Soc Trop Med Hyg.* 1964;58:335–8. [http://dx.doi.org/10.1016/0035-9203\(64\)90201-9](http://dx.doi.org/10.1016/0035-9203(64)90201-9)

Address for correspondence: Anna R. Plourde, University of California San Francisco, Departments of Pathology and Laboratory Medicine, 505 Parnassus Ave, San Francisco, CA 94143, USA; email: anna.plourde@ucsf.edu

A Literature Review of Zika Virus

Technical Appendix

Technical Appendix Table 1. Seroprevalence surveys of Zika virus infection by location, January 1, 1952–February 10, 2016*

Continent/country	% Seropositivity (details)	Years survey performed	Laboratory assessment method	Ref
Africa				
Kenya	17.6 (476/2,698 serum samples)	1966–1968	HI only	(74)
Nigeria†	a) 31.2 (59/189 by HI; 40.3 (121/300 by neutralization) b) 56.2 (150/267) c) 11.5 (15/131 by HI, 1965 and 1970); 12.2 (18/147 by HI and neutralization, 1970–1971)	a) 1971–1975 (b) 1980 c) 1965, 1970–1971	a) HI and/or neutralization b) HA and HI only c) HI or HI followed by neutralization	a) (75) b) (76) c) (77)
Côte d'Ivoire	47.6 (20/42)	1999	IgG by ELISA only	(78)
Senegal	6.7 (57/852)	1988, 1990	IgM by ELISA only	(79)
The Gambia	86.0 (37/43 by HI); 23.8 (10/28 by CF)	1979	HI or CF	(80)
Uganda‡	a) 12 (261 patients tested)‡ b) 6.1 (8/132) c) 6.1 (6/99)	(a) 1945–1948 (b) 1984 (c) Not specified	(a) Viral neutralization (b) HI only (c) Not specified§	a) (81) b) (82) c) (52)
Tanzania	17 (36 patients tested)‡	1945–1948	Viral neutralization	(81)
Zambia	6.0 (217/3,625)	Unspecified	Unspecified¶	(83)
Asia				
Borneo	44.1 (49/111)	1996–1997	“Plasma samples were tested for neutralizing antibody...sera were screened at a 1:10 dilution. Endpoint titrations were conducted for samples that tested positive (i.e., titer >10).”	(84)
Pakistan	2.3 (1/43)	1979, 1980	Initial screen for antibodies (not further specified), followed by CF	(85)
Oceania				
Yap Island, Micronesia	74.3 (414/557)	2007	IgM by ELISA only	(5)

*Summary data of all seroprevalence surveys of Zika virus antibodies in humans, reported in peer-reviewed literature, 1952–2016. Publications for which the abstract only was accessible or that did not provide complete data were excluded. CF, complement fixation; HA, hemagglutination assay; HI, hemagglutination-inhibition; Ref, Technical Appendix reference number.

† Multiple studies were found for Nigeria and Uganda; these studies are indicated separately with a, b, c.

‡ Study for Uganda and United Republic of Tanzania reported only numbers of patients tested and whole percentages of seropositive patients (81).

§Methodology information provided in primary source: “Human sera tested for presence of Zika virus antibody...Only those sera which were found to neutralize 100 or more LD50 have been considered as positive” (52).

¶Methodology information provided in primary source: “Serum was assessed for IgG and IgM antibodies” (83).

Technical Appendix Table 2. Outbreaks and case reports of laboratory-confirmed Zika virus infection by location, January 1, 1952–February 10, 2016*

Continent/country	No. cases	Years	Ref	Comments
Africa				
Cape Verde	17	2015	(86)	15 IgM positive only; 2 positive by reverse transcription PCR
Gabon	5	2007–2010	(23)	
Nigeria	5	1952, 1971–1975	(2,75)	
Uganda	1	Not specified	(40)	
Asia				
Cambodia	1	2010	(56)	
Indonesia	7	1977–1978	(24)	
Japan	3	2013–2014	(87,88)	Imported cases (ex-Thailand, ex-French Polynesia)
Thailand	7	2012–2014	(89)	
Philippines	1	2012	(67)	
Caribbean				
Dominican Republic	10	2016	(90)	8 local cases; 2 imported cases (ex-El Salvador)
Guadeloupe	10	2016	(91)	
Jamaica	1	2016	(92)	
St. Croix, US Virgin Islands	1	2016	(93)	
Central America				
Costa Rica	2	2016	(94)	1 case imported (ex-Colombia); other case not specified
Mexico	3	2015	(95)	2 local cases; 1 imported case (ex-Colombia)
Nicaragua	2	2016	(91,96)	
Europe				
Denmark	1	2016	(91,97)	Imported case (ex-Mexico and Brazil)
Finland	1	2015	(3)	Imported case (ex-Maldives)
France	1	2013	(98)	Imported case (ex-French Polynesia)
Germany	4	2013, 2014, 2016	(57,99,100)	Imported cases (ex-Haiti, ex-Malaysian Borneo, ex-Thailand)
Italy	3	2013–2014, 2015	(60,101)	
Netherlands	13	2015	(102,103)	Imported cases (ex-Suriname [1]; country of import unspecified [12])
Norway	1	2013	(61)	Imported case (ex-Tahiti)
Portugal	5	2016	(91,104)	Imported cases (ex-Brazil)
Spain	7	2016	(105)	Imported cases (country of import unspecified)
Sweden	1	2015	(91,106)	Imported case (country of import unspecified)
United Kingdom	6	2014–2016	(107)	Imported cases (ex-Colombia [3], ex-Cook-Islands [1], ex-Guyana/Suriname [1], and ex-Mexico/Venezuela [1])
North America				
Canada	2	2013, 2016	(14,46)	Imported cases (ex-Thailand, ex-El Salvador)
United States	56	2008, 2013, 2016	(10,12,36,68)	Cases reported in Arkansas (1), California (2), Colorado (3), Delaware (1), District of Columbia (3), Florida (16), Georgia (1), Hawaii (4), Illinois (3), Massachusetts (2), Minnesota (1), Nebraska (2), New Jersey (1), New York (1), Ohio (1), Oregon (1), Pennsylvania (2), Virginia (1), Texas (10). All were imported cases, except for 2 autochthonous sexually-acquired cases, including 1 in Colorado and 1 in Texas. Reported locations of import include Senegal, French Polynesia, South America, Easter Island, and Hawaii.
Oceania				
Australia	3	2014 (1 case); year unspecified (2 cases)	(39,108,109)	Imported cases (ex-Indonesia [2], ex-Cook Islands [1])
Cook Islands	50	2014	(6)	932 additional suspected cases
Easter Island	53	2014	(7,110,111)	One imported case (ex-Tahiti); 40 additional suspected cases

Continent/country	No. cases	Years	Ref	Comments
Fiji	1	2015	(112)	
French Polynesia	383	2013–2014	(14)	32,000 additional suspected cases
New Caledonia	1,385	2014	(8)	
New Zealand	67	2002–2015, 2016	(113)	Imported cases (ex-Cook Islands [27], ex-Vanuatu [1], ex-Tonga [4], ex-Samoa [4], unspecified country of import [1])
Samoa	1 (at least)	2016	(91,114)	
Solomon Islands	4	2015	(115)	
Tonga	1	2016	(113)	
Vanuatu	2	2015	(115)	
Yap, Federated States of Micronesia	108	2007	(5)	919 additional suspected cases
South America				
Barbados	3	2016	(116)	
Bolivia	1	2016	(117)	
Brazil	69	2015–2016	(14,54,118–121)	440,000–1,300,000 estimated total cases
Chile (Mainland)	3	2016	(122)	Imported cases (ex-Venezuela [1], ex-Colombia [1], ex-Brazil [1])
Colombia	1,050	2015	(123)	
Curacao	1	2016	(91)	
Ecuador	6	2016	(116)	2 local cases; 4 imported cases (ex-Colombia [3], ex-Venezuela [1]).
El Salvador	3	2015–2016	(124,125)	3,836 additional suspected cases
French Guiana	59	2016	(91)	
Guatemala	1	2015	(124)	
Guyana	164	2016	(91)	
Haiti	5	2016	(100)	
Honduras	2	2015	(126)	
Martinique	2	2016	(91)	
Panama	4	2015	(127)	
Paraguay	6	2015	(128)	
Puerto Rico	9	2016	(10)	1 imported case; 8 local cases
Saint Martin	1	2016	(129)	
Suriname	6	2015	(130)	
Venezuela	4	2015	(131)	

*Data represent outbreaks and case reports for all reported laboratory-confirmed cases of Zika virus infection (confirmed by serology and/or RT-PCR, or mode of laboratory testing not specified), including those reported in peer-reviewed literature; public health agency Web sites, bulletins, and broadcasts; and media reports, January 1, 1952–February 10, 2016. Ref, Technical Appendix reference number; ex-, exported from.

Technical Appendix Table 3. Symptoms and sequelae of 195 symptomatic patients with confirmed Zika virus infection reported in Peer-reviewed literature for 1964–2016*

Symptom	No. of patients	% Total patients
Rash	131	67.2
Fever	124	63.6
Arthralgia	56	28.7
Myalgia	46	23.6
Headache	42	21.5
Conjunctivitis	40	20.5
Retroorbital pain	22	11.3
Edema	19	9.7
Pruritus	15	7.7
Fatigue/asthenia	14	7.2

*Data represent reports that included clinical symptoms and sequelae for all symptomatic patients with laboratory-confirmed Zika virus infection (confirmed by serologic testing or RT-PCR, or mode of laboratory testing not specified), with a complete and detailed account of symptoms reported in peer-reviewed literature for January 1, 1964–February 3, 2016 (age range of patients was 4 days–76 years). Reports only for this period included symptoms and therefore met criteria for inclusion in the table. Patients may have experienced ≥ 1 symptom. Less prevalent reported symptoms and sequelae (<5% of patients with reported symptoms) include microcephaly, cerebral calcifications, Guillain-Barré syndrome, prostatitis, hematospermia, hematuria, hypotension, gingival bleeding, hearing loss, pruritus, malaise, diarrhea, nausea/vomiting, constipation, edema, sore throat, cough, stomach pain, anorexia, lightheadedness, dizziness, chills, oral aphthous ulcers, photophobia, lymphadenopathy, rhinorrhea, and burning sensation of palms/soles.

References: (3,7,8,11,23–25,34,36,37,39–41,46,47,56,57,60–62,68,75,87,88,99,101,108,109).

References

1. Dick GW, Kitchen SF, Haddow AJ. Zika virus. I. Isolations and serological specificity. *Trans R Soc Trop Med Hyg.* 1952;46:509–20. [http://dx.doi.org/10.1016/0035-9203\(52\)90042-4](http://dx.doi.org/10.1016/0035-9203(52)90042-4)
2. MacNamara FN. Zika virus: a report on three cases of human infection during an epidemic of jaundice in Nigeria. *Trans R Soc Trop Med Hyg.* 1954;48:139–45. [http://dx.doi.org/10.1016/0035-9203\(54\)90006-1](http://dx.doi.org/10.1016/0035-9203(54)90006-1)
3. Korhonen EM, Huhtamo E, Smura T, Kallio-Kokko H, Raassina M, Vapalahti O. Zika virus infection in a traveller returning from the Maldives, June 2015. *Euro Surveill.* 2016;21:30107. <http://dx.doi.org/10.2807/1560-7917.ES.2016.21.2.30107>
4. Baba SS, Fagbami AH, Ojeh CK. Preliminary studies on the use of solid-phase immunosorbent techniques for the rapid detection of Wesselsbron virus (WSLV) IgM by haemagglutination-inhibition. *Comp Immunol Microbiol Infect Dis.* 1999;22:71–9. [http://dx.doi.org/10.1016/S0147-9571\(98\)00003-4](http://dx.doi.org/10.1016/S0147-9571(98)00003-4)
5. Duffy MR, Chen TH, Hancock WT, Powers AM, Kool JL, Lanciotti RS, et al. Zika virus outbreak on Yap Island, Federated States of Micronesia. *N Engl J Med.* 2009;360:2536–43. <http://dx.doi.org/10.1056/NEJMoa0805715>
6. Roth A, Mercier A, Lepers C, Hoy D, Duituturaga S, Benyon E, et al. Concurrent outbreaks of dengue, chikungunya and Zika virus infections - an unprecedented epidemic wave of mosquito-borne viruses in the Pacific 2012-2014. *Euro Surveill.* 2014;19:20929. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.41.20929>
7. Tognarelli J, Ulloa S, Villagra E, Lagos J, Aguayo C, Fasce R, et al. A report on the outbreak of Zika virus on Easter Island, South Pacific, 2014. *Arch Virol.* 2016;161:665–8. <http://dx.doi.org/10.1007/s00705-015-2695-5>
8. Dupont-Rouzeyrol M, O'Connor O, Calvez E, Daurès M, John M, Grangeon JP, et al. Co-infection with Zika and dengue viruses in 2 patients, New Caledonia, 2014. *Emerg Infect Dis.* 2015;21:381–2. <http://dx.doi.org/10.3201/eid2102.141553>
9. Fauci AS, Morens DM. Zika virus in the Americas—yet another arbovirus threat. *N Engl J Med.* 2016;374:601–4. <http://dx.doi.org/10.1056/NEJMp1600297>
10. Centers for Disease Control and Prevention. Zika virus disease in the United States, 2015–2016. 2016 Feb 10 [cited 2016 Feb 17]. <http://www.cdc.gov/zika/geo/united-states.html>

11. Foy BD, Kobylinski KC, Chilson Foy JL, Blitvich BJ, Travassos da Rosa A, Haddow AD, et al. Probable non-vector-borne transmission of Zika virus, Colorado, USA. *Emerg Infect Dis*. 2011;17:880–2. <http://dx.doi.org/10.3201/eid1705.101939>
12. Dallas County Health and Human Services. DCHHS reports first Zika virus case in Dallas County acquired through sexual transmission. 2016 Feb 2 [cited 2016 Feb 8]. <http://www.dallascounty.org/department/hhs/press/documents/PR2-2-16DCHHSReportsFirstCaseofZikaVirusThroughSexualTransmission.pdf>
13. Hennessey M, Fischer M, Staples JE. Zika virus spreads to new areas—region of the Americas, May 2015–January 2016. *MMWR Morb Mortal Wkly Rep*. 2016;65:55–8. <http://dx.doi.org/10.15585/mmwr.mm6503e1>
14. European Centre for Disease Prevention and Control. Zika virus disease epidemic: potential association with microcephaly and Guillain-Barré syndrome (first update). 2016 Jan 21 [cited 2016 Feb 3]. <http://ecdc.europa.eu/en/publications/Publications/rapid-risk-assessment-zika-virus-first-update-jan-2016.pdf>
15. Lanciotti RS, Kosoy OL, Laven JJ, Velez JO, Lambert AJ, Johnson AJ, et al. Genetic and serologic properties of Zika virus associated with an epidemic, Yap State, Micronesia, 2007. *Emerg Infect Dis*. 2008;14:1232–9. <http://dx.doi.org/10.3201/eid1408.080287>
16. Kuno G, Chang GJ. Full-length sequencing and genomic characterization of Bagaza, Kedougou, and Zika viruses. *Arch Virol*. 2007;152:687–96. <http://dx.doi.org/10.1007/s00705-006-0903-z>
17. Gatherer D, Kohl A. Zika virus: a previously slow pandemic spreads rapidly through the Americas. *J Gen Virol*. 2016;97:269–73. <http://dx.doi.org/10.1099/jgv.0.000381>
18. Faye O, Freire CC, Iamarino A, Faye O, de Oliveira JV, Diallo M, et al. Molecular evolution of Zika virus during its emergence in the 20(th) century. *PLoS Negl Trop Dis*. 2014;8:e2636. <http://dx.doi.org/10.1371/journal.pntd.0002636>
19. Hamel R, Dejarnac O, Wichit S, Ekchariyawat P, Neyret A, Luplertlop N, et al. Biology of Zika virus infection in human skin cells. *J Virol*. 2015;89:8880–96. <http://dx.doi.org/10.1128/JVI.00354-15>
20. Buckley A, Gould EA. Detection of virus-specific antigen in the nuclei or nucleoli of cells infected with Zika or Langkat virus. *J Gen Virol*. 1988;69:1913–20. <http://dx.doi.org/10.1099/0022-1317-69-8-1913>
21. Marchette NJ, Garcia R, Rudnick A. Isolation of Zika virus from *Aedes aegypti* mosquitoes in Malaysia. *Am J Trop Med Hyg*. 1969;18:411–5.

22. Hayes EB. Zika virus outside Africa. *Emerg Infect Dis*. 2009;15:1347–50.
<http://dx.doi.org/10.3201/eid1509.090442>
23. Grard G, Caron M, Mombo IM, Nkoghe D, Mboui Ondo S, Jiolle D, et al. Zika virus in Gabon (Central Africa)—2007: a new threat from *Aedes albopictus*? *PLoS Negl Trop Dis*. 2014;8:e2681. <http://dx.doi.org/10.1371/journal.pntd.0002681>
24. Olson JG, Ksiazek TG, Suhandiman, Triwibowo. Zika virus, a cause of fever in Central Java, Indonesia. *Trans R Soc Trop Med Hyg*. 1981;75:389–93. [http://dx.doi.org/10.1016/0035-9203\(81\)90100-0](http://dx.doi.org/10.1016/0035-9203(81)90100-0)
25. Oehler E, Watrin L, Larre P, Leparc-Goffart I, Lastere S, Valour F, et al. Zika virus infection complicated by Guillain-Barre syndrome—case report, French Polynesia, December 2013. *Euro Surveill*. 2014;19:20720. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.9.20720>
26. Diallo D, Sall AA, Diagne CT, Faye O, Faye O, Ba Y, et al. Zika virus emergence in mosquitoes in southeastern Senegal, 2011. *PLoS One*. 2014;9:e109442.
<http://dx.doi.org/10.1371/journal.pone.0109442>
27. Boorman JP, Porterfield JS. A simple technique for infection of mosquitoes with viruses; transmission of Zika virus. *Trans R Soc Trop Med Hyg*. 1956;50:238–42. [http://dx.doi.org/10.1016/0035-9203\(56\)90029-3](http://dx.doi.org/10.1016/0035-9203(56)90029-3)
28. Musso D, Nilles EJ, Cao-Lormeau VM. Rapid spread of emerging Zika virus in the Pacific area. *Clin Microbiol Infect*. 2014;20:O595–6. <http://dx.doi.org/10.1111/1469-0691.12707>
29. Ledermann JP, Guillaumot L, Yug L, Saweyog SC, Tided M, Machieng P, et al. *Aedes hensilli* as a potential vector of Chikungunya and Zika viruses. *PLoS Negl Trop Dis*. 2014;8:e3188.
<http://dx.doi.org/10.1371/journal.pntd.0003188>
30. Kraemer MU, Sinka ME, Duda KA, Mylne A, Shearer FM, Brady OJ, et al. The global compendium of *Aedes aegypti* and *Ae. albopictus* occurrence. *Sci Data*. 2015;2:150035.
<http://dx.doi.org/10.1038/sdata.2015.35>
31. Gaffigan TVWR, Pecor JE, Stoffer JA, Anderson T. Systematic catalog of culicidae. Suitland (MD): Smithsonian Institution; Silver Spring (MD): Walter Reed Army Institute of Research; 2016 [cited 2016 Feb 10]. <http://mosquitocatalog.org/>
32. Ioos S, Mallet HP, Leparc Goffart I, Gauthier V, Cardoso T, Herida M. Current Zika virus epidemiology and recent epidemics. *Med Mal Infect*. 2014;44:302–7.
<http://dx.doi.org/10.1016/j.medmal.2014.04.008>

33. Haddow AD, Schuh AJ, Yasuda CY, Kasper MR, Heang V, Huy R, et al. Genetic characterization of Zika virus strains: geographic expansion of the Asian lineage. *PLoS Negl Trop Dis*. 2012;6:e1477. <http://dx.doi.org/10.1371/journal.pntd.0001477>
34. Oliveira Melo AS, Malinge G, Ximenes R, Szejnfeld PO, Alves Sampaio S, Bispo de Filippis AM. Zika virus intrauterine infection causes fetal brain abnormality and microcephaly: tip of the iceberg? *Ultrasound Obstet Gynecol*. 2016;47:6–7. <http://dx.doi.org/10.1002/uog.15831>
35. Besnard M, Lastere S, Teissier A, Cao-Lormeau V, Musso D. Evidence of perinatal transmission of Zika virus, French Polynesia, December 2013 and February 2014. *Euro Surveill*. 2014;19:20751. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.13.20751>
36. Musso D, Roche C, Robin E, Nhan T, Teissier A, Cao-Lormeau VM. Potential sexual transmission of Zika virus. *Emerg Infect Dis*. 2015;21:359–61. <http://dx.doi.org/10.3201/eid2102.141363>
37. Musso D, Nhan T, Robin E, Roche C, Bierlaire D, Zisou K, et al. Potential for Zika virus transmission through blood transfusion demonstrated during an outbreak in French Polynesia, November 2013 to February 2014. *Euro Surveill*. 2014;19:20761. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.14.20761>
38. US Food and Drug Administration. Recommendations for donor screening, deferral, and product management to reduce the risk of transfusion transmission of Zika virus, recommendations for industry. 2016 Feb 16 [cited 2016 Feb 16]. <http://www.fda.gov/downloads/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/Blood/UCM486360.pdf>
39. Leung GH, Baird RW, Druce J, Anstey NM. Zika virus infection in Australia following a monkey bite in Indonesia. *Southeast Asian J Trop Med Public Health*. 2015;46:460–4.
40. Simpson DI. Zika virus infection in man. *Trans R Soc Trop Med Hyg*. 1964;58:335–8. [http://dx.doi.org/10.1016/0035-9203\(64\)90201-9](http://dx.doi.org/10.1016/0035-9203(64)90201-9)
41. Filipe AR, Martins CM, Rocha H. Laboratory infection with Zika virus after vaccination against yellow fever. *Arch Gesamte Virusforsch*. 1973;43:315–9. <http://dx.doi.org/10.1007/BF01556147>
42. Martines RB, Bhatnagar J, Keating MK, Silva-Flannery L, Muehlenbachs A, Gary J, et al. Notes from the field: evidence of Zika virus infection in brain and placental tissues from two congenitally infected newborns and two fetal losses—Brazil, 2015. *MMWR Morb Mortal Wkly Rep*. 2016;65:159–60. <http://dx.doi.org/10.15585/mmwr.mm6506e1>

43. European Centre for Disease Prevention and Control. Rapid risk assessment. Zika virus epidemic in the Americas: potential association with microcephaly and Guillain-Barré syndrome. 2015 Dec 10. [cited 2016 Feb 3]. <http://ecdc.europa.eu/en/publications/Publications/zika-virus-americas-association-with-microcephaly-rapid-risk-assessment.pdf>
44. Pan American Health Organization; World Health Organization. Epidemiological alert. Neurological syndrome, congenital malformations, and Zika virus infection. Implications for public health in the Americas. 2015 Dec 1 [cited 2016 Feb 10]. 2015-dec-1-cha-epi-alert-zika-neuro-syndrome.pdf
45. Mlakar J, Korva M, Tul N, Popović M, Poljšak-Prijatelj M, Mraz J, et al. Zika virus associated with microcephaly. *N Engl J Med*. 2016;374:951–8. <http://dx.doi.org/10.1056/NEJMoa1600651>
46. Fonseca K, Meatherall B, Zarra D, Drebot M, MacDonald J, Pabbaraju K, et al. First case of Zika virus infection in a returning Canadian traveler. *Am J Trop Med Hyg*. 2014;91:1035–8. <http://dx.doi.org/10.4269/ajtmh.14-0151>
47. Zanluca C, Melo VC, Mosimann AL, Santos GI, Santos CN, Luz K. First report of autochthonous transmission of Zika virus in Brazil. *Mem Inst Oswaldo Cruz*. 2015;110:569–72. <http://dx.doi.org/10.1590/0074-02760150192>
48. Schuler-Faccini L, Ribeiro EM, Feitosa IM, Horovitz DD, Cavalcanti DP, Pessoa A, et al.; Brazilian Medical Genetics Society–Zika Embryopathy Task Force. Possible association between Zika virus infection and microcephaly—Brazil, 2015. *MMWR Morb Mortal Wkly Rep*. 2016;65:59–62. <http://dx.doi.org/10.15585/mmwr.mm6503e2>
49. Simmins CH Jr. Establishing base levels of microcephaly in Brazil prior to the arrival of Zika viral illnesses. *Bull World Health Organ*. 2016 [cited 2016 Feb 10]. Epub 2016 Feb 8.
50. World Health Organization. Zika situation report: Zika and potential complications, 12 February 2016. 2016 Feb 12 [cited 2016 Feb 12]. <http://www.who.int/emergencies/zika-virus/situation-report/who-zika-situation-report-12-02-2016.pdf?ua=1>
51. European Centre for Disease Prevention and Control. Rapid risk assessment: microcephaly in Brazil potentially linked to the Zika virus epidemic. 2015 Nov 24 [cited 2016 Feb 3]. <http://ecdc.europa.eu/en/publications/Publications/zika-microcephaly-Brazil-rapid-risk-assessment-Nov-2015.pdf>
52. Dick GW. Zika virus. II. Pathogenicity and physical properties. *Trans R Soc Trop Med Hyg*. 1952;46:521–34. [http://dx.doi.org/10.1016/0035-9203\(52\)90043-6](http://dx.doi.org/10.1016/0035-9203(52)90043-6)

53. Bell TM, Field EJ, Narang HK. Zika virus infection of the central nervous system of mice. *Arch Gesamte Virusforsch.* 1971;35:183–93. <http://dx.doi.org/10.1007/BF01249709>
54. Tetro JA. Zika and microcephaly: causation, correlation, or coincidence? *Microbes Infect.* 2016;18:167–8. <http://dx.doi.org/10.1016/j.micinf.2015.12.010>
55. World Health Organization. Zika situation report: neurological syndrome and congenital anomalies. 2016 Feb 5 [cited 2016 Feb 8]. <http://www.who.int/emergencies/zika-virus/situation-report/5-february-2016/en/>
56. Heang V, Yasuda CY, Sovann L, Haddow AD, Travassos da Rosa AP, Tesh RB, et al. Zika virus infection, Cambodia, 2010. *Emerg Infect Dis.* 2012;18:349–51. <http://dx.doi.org/10.3201/eid1802.111224>
57. Tappe D, Nachtigall S, Kapaun A, Schnitzler P, Günther S, Schmidt-Chanasit J. Acute Zika virus infection after travel to Malaysian Borneo, September 2014. *Emerg Infect Dis.* 2015;21:911–3. <http://dx.doi.org/10.3201/eid2105.141960>
58. Oster AM, Brooks JT, Stryker JE, Kachur RE, Mead P, Pesik NT, et al. Interim guidelines for prevention of sexual transmission of Zika Virus—United States, 2016. *MMWR Morb Mortal Wkly Rep.* 2016;65:120–1. <http://dx.doi.org/10.15585/mmwr.mm6505e1>
59. Arzuza-Ortega LPA, Pérez-Tatis G, López-García H, Parra E, Pardo-Herrera LC, Rico-Turca AM, et al. Fatal Zika virus infection in girl with sickle cell disease, Colombia [letter]. *Emerg Infect Dis.* 2016 May [date cited]. <http://dx.doi.org/10.3201/eid2205.151934>
60. Zammarchi L, Stella G, Mantella A, Bartolozzi D, Tappe D, Günther S, et al. Zika virus infections imported to Italy: clinical, immunological and virological findings, and public health implications. *J Clin Virol.* 2015;63:32–5. <http://dx.doi.org/10.1016/j.jcv.2014.12.005>
61. Wæhre T, Maagard A, Tappe D, Cadar D, Schmidt-Chanasit J. Zika virus infection after travel to Tahiti, December 2013. *Emerg Infect Dis.* 2014;20:1412–4. <http://dx.doi.org/10.3201/eid2008.140302>
62. Gourinat AC, O'Connor O, Calvez E, Goarant C, Dupont-Rouzeyrol M. Detection of Zika virus in urine. *Emerg Infect Dis.* 2015;21:84–6. <http://dx.doi.org/10.3201/eid2101.140894>
63. Centers for Disease Control and Prevention. Updated diagnostic testing for Zika, chikungunya, and dengue viruses in US public health laboratories. 2016 Feb 7 [cited 2016 Feb 8]. <http://www.cdc.gov/zika/pdfs/denvchikvzika-testing-algorithm.pdf>

64. Centers for Disease Control and Prevention. New CDC laboratory test for Zika virus authorized for emergency use by FDA. 2016 Feb 26 [cited 2016 Feb 26].
<http://www.cdc.gov/media/releases/2016/s0226-laboratory-test-for-zika-virus.html>
65. Centers for Disease Control and Prevention. Zika virus: diagnostic testing. 2016 Feb 11 [cited 2016 Feb 11]. <http://www.cdc.gov/zika/hc-providers/diagnostic.html>
66. Musso D, Roche C, Nhan TX, Robin E, Teissier A, Cao-Lormeau VM. Detection of Zika virus in saliva. *J Clin Virol*. 2015;68:53–5. <http://dx.doi.org/10.1016/j.jcv.2015.04.021>
67. Alera MT, Hermann L, Tac-An IA, Klungthong C, Rutvisuttinunt W, Manasatienkij W, et al. Zika virus infection, Philippines, 2012. *Emerg Infect Dis*. 2015;21:722–4.
<http://dx.doi.org/10.3201/eid2104.141707>
68. Summers DJ, Acosta RW, Acosta AM. Zika virus in an American recreational traveler. *J Travel Med*. 2015;22:338–40. <http://dx.doi.org/10.1111/jtm.12208>
69. Petersen EE, Staples JE, Meaney-Delman D, Fischer M, Ellington SR, Callaghan WM, et al. Interim guidelines for pregnant women during a Zika virus outbreak—United States, 2016. *MMWR Morb Mortal Wkly Rep*. 2016;65:30–3. <http://dx.doi.org/10.15585/mmwr.mm6502e1>
70. Oduyebo T, Petersen EE, Rasmussen SA, Mead PS, Meaney-Delman D, Renquist CM, et al. Update: interim guidelines for health care providers caring for pregnant women and women of reproductive age with possible Zika virus exposure—United States, 2016. *MMWR Morb Mortal Wkly Rep*. 2016;65:122–7. <http://dx.doi.org/10.15585/mmwr.mm6505e2>
71. Reiter P, Lathrop S, Bunning M, Biggerstaff B, Singer D, Tiwari T, et al. Texas lifestyle limits transmission of dengue virus. *Emerg Infect Dis*. 2003;9:86–9.
<http://dx.doi.org/10.3201/eid0901.020220>
72. Aubry M, Richard V, Green J, Broult J, Musso D. Inactivation of Zika virus in plasma with amotosalen and ultraviolet A illumination. *Transfusion*. 2016;56:33–40.
<http://dx.doi.org/10.1111/trf.13271>
73. News BBC. Q&A: Abortion rules in Zika-affected countries. 2016 Jan 29 [cited 2016 Feb 10].
<http://www.bbc.com/news/world-latin-america-35438404>
74. Geser A, Henderson BE, Christensen S. A multipurpose serological survey in Kenya. 2. Results of arbovirus serological tests. *Bull World Health Organ*. 1970;43:539–52.
75. Fagbami AH. Zika virus infections in Nigeria: virological and seroepidemiological investigations in Oyo State. *J Hyg (Lond)*. 1979;83:213–9. <http://dx.doi.org/10.1017/S0022172400025997>

76. Adekolu-John EO, Fagbami AH. Arthropod-borne virus antibodies in sera of residents of Kainji Lake Basin, Nigeria 1980. *Trans R Soc Trop Med Hyg.* 1983;77:149–51.
[http://dx.doi.org/10.1016/0035-9203\(83\)90053-6](http://dx.doi.org/10.1016/0035-9203(83)90053-6)
77. Monath TP, Wilson DC, Casals J. The 1970 yellow fever epidemic in Okwoga District, Benue Plateau State, Nigeria. 3. Serological responses in persons with and without pre-existing heterologous group B immunity. *Bull World Health Organ.* 1973;49:235–44.
78. Akoua-Koffi C, Diarrassouba S, Béné VB, Ngbichi JM, Bozoua T, Bosson A, et al. [Investigation surrounding a fatal case of yellow fever in Côte d’Ivoire in 1999]. *Bull Soc Pathol Exot.* 2001;94:227–30.
79. Monlun E, Zeller H, Le Guenno B, Traoré-Lamizana M, Hervy JP, Adam F, et al. [Surveillance of the circulation of arbovirus of medical interest in the region of eastern Senegal]. *Bull Soc Pathol Exot.* 1993;86:21–8.
80. Monath TP, Craven RB, Muth DJ, Trautt CJ, Calisher CH, Fitzgerald SA. Limitations of the complement-fixation test for distinguishing naturally acquired from vaccine-induced yellow fever infection in flavivirus-hyperendemic areas. *Am J Trop Med Hyg.* 1980;29:624–34.
81. Smithburn KC. Neutralizing antibodies against certain recently isolated viruses in the sera of human beings residing in East Africa. *J Immunol.* 1952;69:223–34.
82. Rodhain F, Gonzalez JP, Mercier E, Helynck B, Larouze B, Hannoun C. Arbovirus infections and viral haemorrhagic fevers in Uganda: a serological survey in Karamoja district, 1984. *Trans R Soc Trop Med Hyg.* 1989;83:851–4. [http://dx.doi.org/10.1016/0035-9203\(89\)90352-0](http://dx.doi.org/10.1016/0035-9203(89)90352-0)
83. Babaniyi OA, Mwaba P, Mulenga D, Monze M, Songolo P, Mazaba-Liwewe ML, et al. Risk assessment for yellow Fever in Western and north-Western provinces of Zambia. *J Glob Infect Dis.* 2015;7:11–7. <http://dx.doi.org/10.4103/0974-777X.150884>
84. Wolfe ND, Kilbourn AM, Karesh WB, Rahman HA, Bosi EJ, Cropp BC, et al. Sylvatic transmission of arboviruses among Bornean orangutans. *Am J Trop Med Hyg.* 2001;64:310–6.
85. Darwish MA, Hoogstraal H, Roberts TJ, Ahmed IP, Omar F. A sero-epidemiological survey for certain arboviruses (Togaviridae) in Pakistan. *Trans R Soc Trop Med Hyg.* 1983;77:442–5.
[http://dx.doi.org/10.1016/0035-9203\(83\)90106-2](http://dx.doi.org/10.1016/0035-9203(83)90106-2)
86. World Health Organization. Zika virus infection—Cape Verde. 2015 Dec 21 [cited 2016 Feb 3].
<http://www.who.int/csr/don/21-december-2015-zika-cape-verde/en/>

87. Kutsuna S, Kato Y, Takasaki T, Moi M, Kotaki A, Uemura H, et al. Two cases of Zika fever imported from French Polynesia to Japan, December 2013 to January 2014 [corrected]. *Euro Surveill.* 2014;19:20683. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.4.20683>
88. Shinohara K, Kutsuna S, Takasaki T, Moi ML, Ikeda M, Kotaki A, et al. Zika fever imported from Thailand to Japan, and diagnosed by PCR in the urines. *J Travel Med.* 2016;23:tav011. <http://dx.doi.org/10.1093/jtm/tav011>
89. Buathong R, Hermann L, Thaisomboonsuk B, Rutvisuttinunt W, Klungthong C, Chinnawirotpisan P, et al. Detection of Zika Virus Infection in Thailand, 2012-2014. *Am J Trop Med Hyg.* 2015;93:380–3. <http://dx.doi.org/10.4269/ajtmh.15-0022>
90. World Health Organization. Zika virus infection—Dominican Republic. 2016 Jan 27 [cited 2016 Feb 3]. <http://www.who.int/csr/don/27-january-2016-zika-dominican-republic/en/>
91. European Centre for Disease Prevention and Control. Epidemiological update: outbreaks of Zika virus and complications potentially linked to the Zika virus infection. 2016 Feb 5 [cited 2016 Feb 5].
92. Jamaica Ministry of Health. Jamaica confirms first case of the Zika virus. 2016 Jan 30 [cited 2016 Feb 3]. <http://moh.gov.jm/jamaica-confirms-first-case-of-the-zika-virus/>
93. World Health Organization. Zika virus infection—United States of America—United States Virgin Islands. 2016 Jan 29 [cited 2016 Feb 3]. <http://www.who.int/csr/don/29-january-2016-zika-usa/en/>
94. Dyer Z. First Zika virus case reported in Costa Rica. *The Tico Times.* 2016 Jan 26 [cited 2016 Feb 10]. <http://www.ticotimes.net/2016/01/26/first-zika-virus-case-reported-costa-rica>
95. World Health Organization. Zika virus infection—Mexico. 2015 Dec 3 [cited 2016 Feb 3]. <http://www.who.int/csr/don/03-december-2015-zika-mexico/en/>
96. Decavele J. Honduras, Nicaragua report first cases of Zika in pregnant women. *Reuters.* 2016 Feb 4 [cited 2016 Feb 8]. <http://www.reuters.com/article/us-health-zika-nicaragua-idUSKCN0VD201>
97. Zander C. Denmark reports first case of Zika virus. Patient, currently hospitalized, had been traveling in Latin America. 2016 Jan 27 [cited 2016 Feb 10]. <http://www.wsj.com/articles/denmark-reports-first-case-of-zika-virus-1453895642>
98. Baronti C, Piorkowski G, Charrel RN, Boubis L, Leparç-Goffart I, de Lamballerie X. Complete coding sequence of Zika virus from a French Polynesia outbreak in 2013. *Genome Announc.* 2014;2:e00500–14. <http://dx.doi.org/10.1128/genomeA.00500-14>

99. Tappe D, Rissland J, Gabriel M, Emmerich P, Gunther S, Held G, et al. First case of laboratory-confirmed Zika virus infection imported into Europe, November 2013. *Euro Surveill.* 2014;19:20685. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.4.20685>
100. World Health Organization. Zika virus infection—Haiti. 2016 Jan 21 [cited 2016 Feb 3]. <http://www.who.int/csr/don/21-january-2016-zika-haiti/en/>
101. Zammarchi L, Tappe D, Fortuna C, Remoli ME, Günther S, Venturi G, et al. Zika virus infection in a traveller returning to Europe from Brazil, March 2015. *Euro Surveill.* 2015;20:21153. <http://dx.doi.org/10.2807/1560-7917.ES2015.20.23.21153>
102. European Centre for Disease Prevention and Control. Epidemiological update: outbreaks of Zika virus and complications potentially linked to the Zika virus infection. 2015 Dec 18 [cited 2016 Feb 3]. http://ecdc.europa.eu/en/press/news/_layouts/forms/News_DispatchForm.aspx?ID=1342&List=8db7286c-fe2d-476c-9133-18ff4cb1b568&Source=http%3A%2F%2Fecdc%2Eeuropa%2Eeu%2Fen%2Fpress%2Fepidemiological_updates%2FPages%2Fepidemiological_updates%2Easpx
103. Rijksinstituut voor Volksgezondheid en Milieu. Zika virus also in Curaçao. 2016 Jan 28 [cited 2016 Feb 8]. http://www.rivm.nl/Documenten_en_publicaties/Algemeen_Actueel/Nieuwsberichten/2016/Zika_virus_ook_op_Cura%C3%A7ao
104. Decavale J. Portugal has five mild cases of Zika virus, all after Brazil trips. Reuters. 2016 Jan 27 [cited 2016 Feb 8]. <http://www.reuters.com/article/health-zika-portugal-idUSKCN0V5256>
105. De Sanidad M, Sociales e Igualdad S. Weekly update of Zika cases confirmed by the National Center for Microbiology. 2016 Feb 4 [cited 2016 Feb 8]. <http://www.msssi.gob.es/gabinete/notasPrensa.do?id = 3895>
106. Huuhtanen M. Danish hospital says case of Zika virus discovered. Associated Press. 2016 Jan 27 [cited 2016 Feb 8]. <http://bigstory.ap.org/article/4491870e5b274776bf91780a19457b58/danish-hospital-says-case-zika-virus-discovered>
107. Public Health England. Zika virus. Clinical advice on Zika: assessing pregnant women following travel; symptoms, transmission (includes sexual transmission), epidemiology. GOV.UK. 2016 Feb 17 [cited 2016 Feb 17]. <https://www.gov.uk/guidance/zika-virus>

108. Kwong JC, Druce JD, Leder K. Zika virus infection acquired during brief travel to Indonesia. *Am J Trop Med Hyg.* 2013;89:516–7. <http://dx.doi.org/10.4269/ajtmh.13-0029>
109. Pyke AT, Daly MT, Cameron JN, Moore PR, Taylor CT, Hewitson GR, et al. Imported Zika virus infection from the Cook Islands into Australia, 2014. *PLoS Curr.* 2014;6:6.
110. Schwan K. Zika virus—Pacific (07): Chile (Easter Island), French Polynesia. *ProMed.* 2014 Mar 3 [cited 2016 Feb 10]. <http://www.promedmail.org/post/2322907>, archive no. 20140309.2322907.
111. Cooperativa (Chile). Zika virus—Pacific (15): Chile (Easter Island) ex Tahiti. *ProMed.* 2014 Sept 25 [cited 2016 Feb 10]. <http://www.promedmail.org/post/2804821>, archive no. 20140925.2804821.
112. World Health Organization Western Pacific Region. Pacific syndromic surveillance report—Week 33, ending 16 August, 2015. 2015 Aug 16 [cited 2016 Feb 3]. http://www.wpro.who.int/southpacific/programmes/communicable_diseases/disease_surveillance_response/PSS-16-August-2015/en/
113. New Zealand Ministry of Health. Zika virus update. The Ministry of Health has extended its Pacific travel advice around the Zika virus to include Tonga as well as Samoa as an area of active transmission. 2016 Jan 29 [cited 2016 Feb 3]. <http://www.health.govt.nz/news-media/media-releases/zika-virus-update>
114. Centers for Disease Control and Prevention. Zika virus in the Pacific Islands. 2016 Feb 12 [cited 2016 Feb 16]. <http://wwwnc.cdc.gov/travel/notices/alert/zika-virus-south-pacific>
115. World Health Organization Western Pacific Region. Pacific syndromic surveillance report, Week 17, ending 26 April, 2015. 2015 Apr 26 [cited 2016 Jan 29]. http://www.wpro.who.int/southpacific/programmes/communicable_diseases/disease_surveillance_response/PSS-26-April-2015/en/
116. World Health Organization. Zika virus infection—Guyana, Barbados and Ecuador. 2016 Jan 20 [cited 2016 Feb 8]. <http://www.who.int/csr/don/20-january-2016-zika-guyana-barbados-ecuador/en/>
117. World Health Organization. Zika virus infection—Bolivia. 2016 Jan 20 [cited 2016 Feb 8]. <http://www.who.int/csr/don/20-january-2016-zika-bolivia/en/>
118. Herriman R. Zika virus—Brazil (05). *ProMed.* 2015 Jun 12 [cited 2016 Feb 8]. <http://www.promedmail.org/post/3431148>, archive no. 20150612.3431148.
119. Castelo Branco B. Zika virus—Brazil (06): (Bahia). *ProMed.* 2015 Jun 19 [cited 2016 Feb 8]. <http://www.promedmail.org/post/3449500>, archive no. 20150619.3449500.

120. Zika virus—Brazil. (07). ProMed. 2015 Jun 30 [cited 2016 Feb 8].
<http://www.promedmail.org/post/3473420>, archive no. 20150630.3473420.
121. Zika virus—Brazil. (08). ProMed. 2015 July 16 [cited 2016 Feb 8].
<http://www.promedmail.org/post/3513770>, archive no. 20150716.3513770.
122. Associated Press. The latest: Chile reports its first 3 confirmed Zika cases. 2016 Feb 2 [cited 2016 Feb 8]. <http://bigstory.ap.org/article/52f57f3f5e0f48f5a9469acad6318faf/latest-4th-case-zika-identified-spain>
123. Instituto Nacional de Salud (National Institutes of Health Colombia). Epidemiologic week number 3 2016 (17 Jan–23 Jan). 2016 Jan 23 [cited 2016 Feb 8]. <http://www.ins.gov.co/boletin-epidemiologico/Boletin%20Epidemiologico/2016%20Boletin%20epidemiologico%20semana%203.pdf>
124. World Health Organization. Zika virus infection—Guatemala. 2015 Nov 27 [cited 2016 Feb 8].
<http://www.who.int/csr/don/27-november-2015-zika-guatemala/en/>
125. World Health Organization. Guillain-Barré syndrome—El Salvador. 2016 Jan 21 [cited 2016 Feb 8].
<http://www.who.int/csr/don/21-january-2016-gbs-el-salvador/en/>
126. World Health Organization. Zika virus infection—Honduras. 2015 Dec 21 [cited 2016 Feb 8].
<http://www.who.int/csr/don/21-december-2015-zika-honduras/en/>
127. World Health Organization. Zika virus infection—Panama. 2015 Dec 22 [cited 2016 Feb 3].
<http://www.who.int/csr/don/22-december-2015-zika-panama/en/>
128. World Health Organization. Zika virus infection—Paraguay. 2015 Dec 3 [cited 2016 Feb 3].
<http://www.who.int/csr/don/03-december-2015-zika-paraguay/en/>
129. World Health Organization. Zika virus infection—France—Saint Martin and Guadeloupe. 2016 Jan 21 [cited 2016 Feb 3]. <http://www.who.int/csr/don/21-january-2016-zika-france/en/>
130. Enfissi A, Codrington J, Roosblad J, Kazanji M, Rousset D. Zika virus genome from the Americas. *Lancet*. 2016;387:227–8. [http://dx.doi.org/10.1016/S0140-6736\(16\)00003-9](http://dx.doi.org/10.1016/S0140-6736(16)00003-9)
131. World Health Organization. Zika virus infection—Venezuela. 2015 Dec 3 [cited 2016 Feb 3].
<http://www.who.int/csr/don/03-december-2015-zika-venezuela/en/>