

Serologic Evidence of Fruit Bat Exposure to Filoviruses, Singapore, 2011–2016

Eric D. Laing,¹ Ian H. Mendenhall,¹ Martin Linster,
Dolyce H. W. Low, Yihui Chen, Lianying Yan,
Spencer L. Sterling, Sophie Borthwick,
Erica Sena Neves, Julia S. L. Lim, Maggie Skiles,
Benjamin P. Y. -H. Lee, Lin-Fa Wang,
Christopher C. Broder, Gavin J. D. Smith

To determine whether fruit bats in Singapore have been exposed to filoviruses, we screened 409 serum samples from bats of 3 species by using a multiplex assay that detects antibodies against filoviruses. Positive samples reacted with glycoproteins from Bundibugyo, Ebola, and Sudan viruses, indicating filovirus circulation among bats in Southeast Asia.

The genus *Ebolavirus* comprises 5 virus species: *Zaire ebolavirus* (EBOV), *Sudan ebolavirus* (SUDV), *Bundibugyo ebolavirus* (BDBV), *Tai Forest ebolavirus* (TAFV), and *Reston ebolavirus* (RESTV). The genus *Marburgvirus* comprises 1 species, *Marburg marburgvirus*, which includes 2 closely related virus strains: Marburg virus (MARV) and Ravn virus (RAVV). Viruses within the *Ebolavirus* and *Marburgvirus* genera are zoonotic; EBOV was the causative agent of the 2014–2016 Ebola virus disease epidemic in West Africa (1). *Rousettus* bats in Africa have been identified as *Marburgvirus* hosts (2), and viral nucleic acid and serologic evidence suggests that bats are also natural hosts of *Ebolavirus* spp. (3). Yet it remains unclear which species are the definitive reservoirs of filoviruses.

Ecologic models of *Ebolavirus* and *Marburgvirus* geographic distribution and habitat ranges of potential reservoir bat species suggest that both groups are distributed throughout Asia (3,4). Serologic evidence of filoviruses in frugivorous bats in Bangladesh, China, and the

Philippines has been reported (5–7), and RESTV nucleic acid was detected in an insectivorous bat in the Philippines, where RESTV is considered endemic (8). We examined pteropodid bats of 3 species: *Cynopterus brachyotis*, *Eonycteris spelaea*, and *Penthetor lucasi*, which are widely distributed across Southeast Asia and share ecologic niches (9).

The Study

During 2011–2016, we collected serum from bats of the 3 aforementioned species in Singapore and screened samples for evidence of exposure to filoviruses. Samples were collected with permission from the National University of Singapore Institutional Animal Care and Use Committee (B01/12) and the National Parks Board (NP/RP11–011–3a). We diluted venous blood 1:10 in phosphate-buffered saline and then centrifuged, recovered, and heat-inactivated the serum at 56°C for 30 minutes and stored it at –80°C.

We developed a Bio-Plex (Bio-Rad, Hercules, CA, USA) bead-based multiplex assay that simultaneously probes serum for immunoglobulins specific to the viral envelope glycoproteins (GPs) from representative strains of all described *Ebolavirus* and *Marburgvirus* species (Table 1). A human FreeStyle 293-F stable cell-line expression system was used to produce the *Ebolavirus* and *Marburgvirus* spp. GPs as a soluble GP consisting of the entire ectodomain, sGP_(1,2), which retains a native-like oligomeric conformation, as described previously with modifications (10). In brief, each GP_(1,2) coding sequence was truncated at the C-terminus to remove the predicted transmembrane domain and cytoplasmic tail, then appended with the GCN trimerization peptide sequence (10) together with a factor Xa protease cleave site and a Twin-Strep-tag sequence (IBA Lifesciences, Göttingen, Germany). The sGP_(1,2) proteins were produced in serum-free conditions and purified by Strep-Tactin XT technology (IBA Lifesciences). The Twin-Strep-tag was removed by factor Xa enzymatic cleavage; factor Xa was removed by Xarrest Agarose (Merck Millipore, Billerica, MA, USA); sGP_(1,2) was purified further by S-200 size exclusion chromatography, concentrated, and stored frozen. These sGP_(1,2)s were coupled to carboxylated beads (Bio-Rad). Screening was performed on a Bio-Rad Bio-Plex 200.

Author affiliations: Uniformed Services University, Bethesda, Maryland, USA (E.D. Laing, L. Yan, S.L. Sterling, C.C. Broder); Duke-National University of Singapore Medical School, Singapore, Singapore (I.H. Mendenhall, M. Linster, D.H.W. Low, Y. Chen, S. Borthwick, E.S. Neves, J.S.L. Lim, L.-F. Wang, G.J.D. Smith); North Carolina State University, Raleigh, North Carolina, USA (M. Skiles); National Parks Board, Singapore (B.P.Y.-H. Lee); Duke University, Durham, North Carolina, USA (L.-F. Wang, G.J.D. Smith)

DOI: <https://doi.org/10.3201/eid2401.170401>

¹These authors contributed equally to this article.

Table 1. *Ebolavirus* and *Marburgvirus* species soluble envelope glycoproteins conjugated Bio-Plex beads used in multiplex assay to detect antibodies against filoviruses*

Virus	Isolation host/location	Bio-Plex bead	
		no.	NCBI accession no.
Ebola virus/H.sapiens/COD/1976/Yambuku-Mayinga	Human/DRC	33	NC_002549.1
Bundibugyo virus/H. sapiens/UGA/2007	Human/Uganda	64	FJ217161.1
Tai Forest virus/H. sapiens/COV/1994/Pauleoula-CI	Human/Côte d'Ivoire	57	NC_014372
Sudan virus/H. sapiens/UGA/2000/Gulu-808892	Human/Uganda	77	NC_006432.1
Reston virus/M. fascicularis/USA/1989/Pennsylvania	Macaque/USA	85	AF522874.1
Reston virus/S. domesticus/PHL/2008/Reston08-A	Swine/Philippines	72	FJ621583.1
Marburg virus/H. sapiens/KEN/1980/Musoke	Human/Kenya	37	Z12132 S55429
Marburg virus/H. sapiens/AGO/2005/Ang0126	Human/Angola	28	DQ447656.1
Ravn virus/H. sapiens/KEN/1987/Kitum cave-810040	Human/Kenya	49	NC_024781.1

*Bio-Plex manufactured by Bio-Rad (Hercules, CA, USA). DRC, Democratic Republic of the Congo; NCBI, National Center for Biotechnology Information.

In the absence of confirmed filovirus-negative bat serum, we used methods developed by Peel et al. to establish a median fluorescence intensity (MFI) cutoff value (11). We confirmed a cutoff value of 200 MFI (online Technical Appendix, <https://wwwnc.cdc.gov/EID/article/24/1/17-0401-Techapp1.pdf>), as was previously used for *Eidolon helvum* bat serum in a Bio-Plex serologic assay (12). We screened 409 samples with our *Ebolavirus* and *Marburgvirus* spp. sGP_(1,2) Bio-Plex assay modified from that described by Bossart et al. (13). Samples were diluted 1:100 and tested in duplicate; the sGP_(1,2)-coupled beads were mixed with individual samples; and a 1:1 combination of recombinant biotinylated-protein A/protein G (1:500) (Pierce, Rockford, IL, USA) was added to the wells, followed by addition of streptavidin-phycoerythrin (1:1,000) (Bio-Rad) and determination of MFI.

Samples were positive for 17 (9.1%) of 186 *E. spelaea*, 13 (8.5%) of 153 *C. brachyotis*, and 3 (4.3%) of 70 *P. lucasi* bats (Figure 1). Positive samples reacted with EBOV, BDBV, SUDV, or TAFV sGP_(1,2). However, no samples were positive for RESTV, MARV, or RAVV sGP_(1,2). We further examined positive samples to determine cross-reactivity between the *Ebolavirus* spp. sGP_(1,2) (Table 2). Twelve (71%) samples from *E. spelaea* bats cross-reacted with ≥ 2 *Ebolavirus* spp. sGP_(1,2) (BDBV, EBOV, SUDV, or TAFV). In contrast, 8 (62%) *C. brachyotis* and 2 (66%) *P. lucasi* samples were positive for only 1 sGP_(1,2) (BDBV or SUDV).

To further determine the cross-reactivity of positive samples and to corroborate Bio-Plex assay results for a selected number of samples, we performed Western blot (WB) assays (Figure 2). The filovirus GP_(1,2) is a trimer of heterodimeric GP₁ and GP₂ subunits. The trimeric-like sGP_(1,2) is the antigen in the multiplex Bio-Plex assay, whereas linearized monomeric sGP₁ and sGP₂ subunits are the antigens in WBs. Reduced and denatured EBOV or BDBV unconjugated sGP_(1,2) was loaded on 8% sodium dodecyl sulfate–polyacrylamide electrophoresis gels, transferred to a polyvinylidene difluoride membrane, and probed with 1:100 dilutions of positive and negative bat serum, as previously determined by the Bio-Plex assay. All 3 *E. spelaea* bat samples and 2 of 3 *C. brachyotis* bat samples that were Bio-Plex positive were also positive by WB and displayed reactivity with EBOV and BDBV GP₁ and GP₂ antigens; no *P. lucasi* bat samples positive by Bio-Plex were positive by WB.

Conclusions

We present evidence of antibodies specific to filoviruses antigenically related to *Ebolavirus* spp. in 3 species of fruit bats widely distributed throughout Southeast Asia. We detected seroreactivity with *Ebolavirus* spp. but not *Marburgvirus* spp. GP. Despite the close relatedness of the viruses, we detected samples reacting with only SUDV, not RESTV, GP. This finding contrasts




Figure 1. Mean fluorescence intensity (MFI) values obtained from Bio-Plex assay (Bio-Rad, Hercules, CA, USA) screening of individual serum samples from bats of 3 species with soluble filovirus glycoproteins. Dashed line indicates the cutoff value at 200 MFI. 1, *Zaire ebolavirus*; 2, *Bundibugyo ebolavirus*; 3, *Tai Forest ebolavirus*; 4, *Sudan ebolavirus*; 5, *Reston ebolavirus*–pig; 6, *Reston ebolavirus*–monkey; 7, *Marburg virus*–Musoke; 8, *Marburg virus*–Angola; 9, *Ravn virus*.

Table 2. Bio-Plex median fluorescence intensity values for bat serum samples positive for ≥ 1 filovirus antigen*

Bat species, ID	EBOV	BDBV	TAFV	SUDV	RESTVm	RESTVp	MARV(Mus)	MARV(Ang)	RAVV
<i>Eonycteris spelaea</i> , n = 186									
0805149†	738			40	44	22	23	21	24
080814	86	318	105	258	26	12	17	16	20
082154	143	161	113	214	35	41	21	31	39
052313	284	408	177	285	89	72	29	23	30
052335	203	191	124	219	42	21	38	38	24
052339	357	306	141	293	54	31	26	26	42
071839	330	299	164	480	65	44	28	33	45
071842	446	327	202	362	65	49	42	38	57
110733	126	416	166	95	58	42	34	42	58
011603†	1151	130	91	69	36	32	51	35	39
011616	252	294	168	175	32	49	47	29	50
011656	306	386	204	394	89	73	18	39	37
012309†	579	659	315	69	35	31	27	33	35
021303	478	431	188	450	52	37	24	30	47
111903	469	384	276	113	52	57	37	69	54
111907	285	336	213	158	39	36	29	50	30
042722	260	262	174	167	75	31	54	24	42
<i>Cynopterus brachyotis</i> , n = 153									
051253	121	133	59	242	40	41	19	25	68
0516613	146	293	127	73	47	36	25	29	22
0516632	138	139	86	356	35	25	28	34	34
0726122†	119	501	100	60	40	46	25	19	29
1103241	84	141	128	241	50	47	66	38	34
100903	148	201	71	108	42	33	18	16	36
100914	74	228	70	55	39	38	30	27	26
100925	166	304	109	116	43	18	33	30	28
021357	201	299	179	264	65	44	25	55	47
050804	242	276	140	124	41	30	34	33	44
050818	383	374	198	332	60	55	29	26	68
040807†	297	597	194	192	40	38	122	95	32
042701†	339	547	222	417	60	78	54	25	62
<i>Penthetor lucasi</i> , n = 70									
062590†	34	496	93	39	36	18	23	17	23
070409†	95	238	129	89	62	27	34	36	37
112112†	251	352	148	235	51	29	23	23	29

*Bio-Plex manufactured by Bio-Rad (Hercules, CA, USA). Boldface indicates positive results. BDBV, Bundibugyo virus; EBOV, Ebola virus; ID, specimen identification number; MARV(Mus), Marburg virus–Musoke; MARV(Ang), Marburg virus–Angola; RESTVm, Reston virus–monkey; RESTVp, Reston virus–pig; SUDV, Sudan virus; RAVV, Ravn virus; TAFV, Tai Forest virus.

†Sample screened by Western blot and shown in Figure 2.

with previous reports of bat serum cross-reactivity with RESTV nucleoprotein (5,7,14). Possible explanations include 1) the fact that our customized Bio-Plex assay is based on conformational sGP_(1,2), which can differentiate antibody specificity better than the more sequence conserved nucleoprotein, and 2) the lack of evidence of RESTV GP positivity with *Cynopterus* and *Eonycteris* bat serum samples, which is in line with previous findings (both species were negative while only *Rousettus amplexicaudatus* bats were positive) (7). *E. spelaea* bats were previously predicted to be filovirus hosts (15), and sequences of novel filoviruses have been discovered in *E. spelaea* bat populations in Yunnan, China (14). Our data provide additional empirical evidence that populations of *C. brachyotis*, *E. spelaea*, and *P. lucasi* bats in Southeast Asia are hosts of filoviruses, which seem antigenically more closely related to EBOV, BDBV, and SUDV than to RESTV.

Examination of cross-reactivity of positive samples from *E. spelaea*, *C. brachyotis*, and *P. lucasi* bats revealed

no clear patterns of preferential reactivity with EBOV, BDBV, or SUDV GP. Factors that might contribute to the lack of *P. lucasi* positivity by WB include sensitivity differences between Bio-Plex and WB assays paired with the change in sGP_(1,2) conformation. Two Bio-Plex EBOV-positive samples (*E. spelaea* samples 0805149 and 011603) reacted with EBOV sGP₂ and BDBV sGP₁ in the WB. Bio-Plex and WB data strongly suggest the presence of yet-undetected batborne filoviruses, which are antigenically related to but distinct from BDBV, EBOV, and SUDV circulating in local bat populations. Reasons why these filoviruses have remained undetected include their inability to cross the species barrier, the rarity of spillovers into humans or domestic animals, or the fact that spillover events cause mild or no disease. We suggest that a yet-undescribed diversity of filoviruses exists in Southeast Asia bat populations, a hypothesis supported by the recent identification of filovirus sequences in *E. spelaea* and *R. leschenaulti* bats in China (14,16). Comprehensive surveillance including serology and detection of viral nucleic acid, along with virus




Figure 2. Western blot results of individual bat serum samples probed against *Zaire ebolavirus* and *Bundibugyo ebolavirus* glycoproteins 1 and 2 (GP₁, GP₂). Boldface indicates positivity by Western blot and underlining indicates positivity by Bio-Plex (Bio-Rad, Hercules, CA, USA). 1, soluble GP₁ and GP₂ blotted with control anti-Ebola virus nonhuman primate polyclonal serum that demonstrates cross-reactivity with *Bundibugyo ebolavirus* soluble GP. Other numbers along baseline correspond to the following sample identifiers, also used in Table 2: 2, 0805149; 3, 012309; 4, 011603; 5, 0116048; 6, 0719036; 7, 1128015; 8, 0726122; 9, 042701; 10, 040807; 11, 0512540; 12, 1009010; 13, 0408029; 14, 070409; 15, 112112; 16, 062590; 17, 0228004; 18, 0919025; 19, 0625095. BDBV, Bundibugyo virus; EBOV, Ebola virus.

isolation, will help elucidate the characteristics of filoviruses endemic to Asia and identify bat species that function as maintenance populations and reservoirs.

Acknowledgment

We thank Alison J. Peel for assistance with determination of the median fluorescence intensity cutoff and statistical advice.

This study was supported by the Duke-National University of Singapore Signature Research Program funded by the Agency of Science, Technology and Research, and the Ministry of Health, Singapore, and by grants from National University of Singapore–Global Asia Institute (NIHA-2011-1-005), the National Medical Research Council (NMRC/BNIG/2005/2013), the Ministry of Health (CDPHRG/0006/2014) in Singapore, and the US Department of Defense, Defense Threat Reduction Agency. C.C.B., E.D.L., L.Y., and S.L.S. were supported by funding from the Biological Defense Research Directorate of the Naval Medical Research Center. E.D.L. was also supported by the National Science Foundation, an East Asia and Pacific Summer Institutes Fellowship award (1515304), with collaborative support from the National University of Singapore.

Dr. Laing is a postdoctoral fellow at the Uniformed Services University and performed this work while a National Science Foundation EAPSI fellow at Duke-National University of Singapore Medical School. His research focuses on biosurveillance, batborne viruses, and antiviral immunity.

References

1. Baize S, Pannetier D, Oestereich L, Rieger T, Koivogui L, Magassouba N, et al. Emergence of Zaire Ebola virus disease

- in Guinea. *N Engl J Med.* 2014;371:1418–25. <http://dx.doi.org/10.1056/NEJMoa1404505>
2. Towner JS, Amman BR, Sealy TK, Carroll SA, Comer JA, Kemp A, et al. Isolation of genetically diverse Marburg viruses from Egyptian fruit bats. *PLoS Pathog.* 2009;5:e1000536. <http://dx.doi.org/10.1371/journal.ppat.1000536>
3. Olival KJ, Hayman DT. Filoviruses in bats: current knowledge and future directions. *Viruses.* 2014;6:1759–88. <http://dx.doi.org/10.3390/v6041759>
4. Peterson AT, Bauer JT, Mills JN. Ecologic and geographic distribution of filovirus disease. *Emerg Infect Dis.* 2004;10:40–7. <http://dx.doi.org/10.3201/eid1001.030125>
5. Olival KJ, Islam A, Yu M, Anthony SJ, Epstein JH, Khan SA, et al. Ebola virus antibodies in fruit bats, Bangladesh. *Emerg Infect Dis.* 2013;19:270–3. <http://dx.doi.org/10.3201/eid1902.120524>
6. Yuan J, Zhang Y, Li J, Zhang Y, Wang LF, Shi Z. Serological evidence of ebolavirus infection in bats, China. *Virology.* 2012;9:236. <http://dx.doi.org/10.1186/1743-422X-9-236>
7. Taniguchi S, Watanabe S, Masangkay JS, Omatsu T, Ikegami T, Alviola P, et al. Reston Ebolavirus antibodies in bats, the Philippines. *Emerg Infect Dis.* 2011;17:1559–60.
8. Jayme SI, Field HE, de Jong C, Olival KJ, Marsh G, Tagtag AM, et al. Molecular evidence of Ebola Reston virus infection in Philippine bats. *Virology.* 2015;12:107. <http://dx.doi.org/10.1186/s12985-015-0331-3>
9. Mendenhall IH, Borthwick S, Neves ES, Low D, Linster M, Liang B, et al. Identification of a lineage D betacoronavirus in cave nectar bats (*Eonycteris spelaea*) in Singapore and an overview of lineage D reservoir ecology in SE Asian bats. *Transbound Emerg Dis.* 2016. <http://dx.doi.org/10.1111/tbed.12568>
10. Chan YP, Yan L, Feng YR, Broder CC. Preparation of recombinant viral glycoproteins for novel and therapeutic antibody discovery. *Methods Mol Biol.* 2009;525:31–58, xiii. http://dx.doi.org/10.1007/978-1-59745-554-1_2
11. Peel AJ, McKinley TJ, Baker KS, Barr JA, Cramer G, Hayman DT, et al. Use of cross-reactive serological assays for detecting novel pathogens in wildlife: assessing an appropriate cutoff for henipavirus assays in African bats. *J Virol Methods.* 2013;193:295–303. <http://dx.doi.org/10.1016/j.jviromet.2013.06.030>

12. Hayman DT, Suu-Ire R, Breed AC, McEachern JA, Wang L, Wood JL, et al. Evidence of henipavirus infection in West African fruit bats. *PLoS One*. 2008;3:e2739. <http://dx.doi.org/10.1371/journal.pone.0002739>
13. Bossart KN, McEachern JA, Hickey AC, Choudhry V, Dimitrov DS, Eaton BT, et al. Neutralization assays for differential henipavirus serology using Bio-Plex protein array systems. *J Virol Methods*. 2007;142:29–40. <http://dx.doi.org/10.1016/j.jviromet.2007.01.003>
14. Yang XL, Zhang YZ, Jiang RD, Guo H, Zhang W, Li B, et al. Genetically diverse filoviruses in *Rousettus* and *Eonycteris* spp. bats, China, 2009 and 2015. *Emerg Infect Dis*. 2017;23:482–6. <http://dx.doi.org/10.3201/eid2303.161119>
15. Han BA, Schmidt JP, Alexander LW, Bowden SE, Hayman DT, Drake JM. Undiscovered bat hosts of filoviruses. *PLoS Negl Trop Dis*. 2016;10:e0004815. <http://dx.doi.org/10.1371/journal.pntd.0004815>
16. He B, Feng Y, Zhang H, Xu L, Yang W, Zhang Y, et al. Filovirus RNA in fruit bats, China. *Emerg Infect Dis*. 2015;21:1675–7. <http://dx.doi.org/10.3201/eid2109.150260>

Address for correspondence: Gavin J. D. Smith, Programme in Emerging Infectious Diseases, Duke-National University Singapore Medical School, 8 College Rd, Singapore 169857, Singapore; email: gavin.smith@duke-nus.edu.sg

September 2017: Zoonoses

- *Candidatus* *Dirofilaria hongkongensis* as Causative Agent of Human Ocular Filariasis after Travel to India
- Mucus-Activatable Shiga Toxin Genotype *stx2d* in *Escherichia coli* O157:H7
- Acute Encephalitis Syndrome and Scrub Typhus in India
- Hematophagous Endeavors, Fact and Fancy
- Processes Underlying Rabies Virus Incursions across US–Canada Border as Revealed by Whole-Genome Phylogeography
- Real-Time Whole-Genome Sequencing for Surveillance of *Listeria monocytogenes*, France
- Role of Food Insecurity in Outbreak of Anthrax Infections



among Humans and Hippopotamuses Living in a Game Reserve Area, Rural Zambia

- Bioinformatic Analyses of Whole-Genome Sequence Data in a Public Health Laboratory
- Serologic Evidence of Powassan Virus Infection in Patients with Suspected Lyme Disease
- Influenza D Virus in Animal Species in Guangdong Province, Southern China
- Seroprevalence of *Baylisascaris procyonis* Infection among Humans, Santa Barbara County, California, USA, 2014–2016
- Opiate Injection–Associated Skin, Soft Tissue, and Vascular Infections, England, UK, 1997–2016

- Risk for Death among Children with Pneumonia, Afghanistan

- Detection of *Elizabethkingia* spp. in *Culicoides* Biting Midges, Australia



- Early Evidence for Zika Virus Circulation among *Aedes aegypti* Mosquitoes, Rio de Janeiro, Brazil

- Scrub Typhus Outbreak in a Remote Primary School, Bhutan, 2014

- Scrub Typhus as a Cause of Acute Encephalitis Syndrome, Gorakhpur, Uttar Pradesh, India

- Human Infection with *Burkholderia thailandensis*, China, 2013

- *mcr-1* and *bla*_{KPC-3} in *Escherichia coli* Sequence Type 744 after Meropenem and Colistin Therapy, Portugal

- Outcomes for 2 Children after Peripartum Acquisition of Zika Virus Infection, French Polynesia, 2013–2014

- California Serogroup Virus Infection Associated with Encephalitis and Cognitive Decline, Canada, 2015

- Effects of Influenza Strain Label on Worry and Behavioral Intentionst

- Zika Virus Screening among Spanish Team Members after 2016 Rio de Janeiro, Brazil, Olympic Games

