

Fishborne Zoonotic Trematodes Transmitted by *Melanoides tuberculata* Snails, Peru

Eduardo A. Pulido-Murillo, Luis Fernando V. Furtado, Alan L. Melo, Élica M.L. Rabelo, Hudson A. Pinto

Author affiliation: Universidade Federal de Minas Gerais, Belo Horizonte, Brazil

DOI: <https://doi.org/10.3201/eid2403.172056>

We investigated the transmission of the fishborne trematodes *Centrocestus formosanus* and *Haplorchis pumilio* by *Melanoides tuberculata* snails in Peru. We report on results of experimental, morphological, and molecular approaches and discuss the potential risk for future human cases, given the existence of food habits in the country involving the ingestion of raw fish.

The World Health Organization has estimated that the number of humans infected with fishborne trematodes exceeds 18 million, and >500 million persons are at risk of infection (1). Among the causative agents of these trematodiasis are representatives of the family Heterophyidae, which are small intestinal parasites from birds and mammals, including humans (1–3). Infection by heterophyids can be considered an emerging disease because of a set of factors, including high prevalence, reported mainly in Asia; outbreaks caused by *Ascocotyle longa* trematodes in Brazil; the introduction of *Centrocestus formosanus* trematodes and *Haplorchis pumilio* flukes from Asia into the Americas; and the involvement of larvae of *Procerovum varium* flukes as causative agents of human ocular disease (3–5).

Of the 30 species of heterophyids recognized worldwide (4), 7, including *C. formosanus* and *H. pumilio*, are transmitted by the red-rimmed melania or Malaysian trumpet snail, *Melanoides tuberculata* (6). Even though human infection by these 2 heterophyids has not been reported in the Americas, the possibility of future cases must be considered, especially in countries like Peru, whose inhabitants consume ceviche, a culinary dish prepared with raw fish. Thus, the evaluation of the involvement of *M. tuberculata* snails in the transmission of heterophyids in this country is needed given the potential public health concern related to these parasites.

In this study, we collected snails in 2 areas of the central coast of Peru, the Ventanilla Wetlands Regional Conservation Area (VWRCA) (11°52'31"S; 77°8'37"W) and the Pantanos de Villa Wildlife Refuge (PVWR) (12°12'33"S; 76°59'28"W), during December 2015 and January, June, and July 2016. Identification of the collected *M. tuberculata* snail specimens

was based on conchiliological features according to previously published reports (7,8). We placed the specimens individually onto polystyrene plates containing dechlorinated water, subjected them to photostimulation, and examined them under a stereomicroscope. We examined cercariae, fluke larvae, in a light microscope after vital staining (0.05% Nile blue sulfate) and preliminarily identified 2 cercarial types, pleurolophocercous (Figure, panel A) and parapleurolophocercous (Figure, panel E). We used samples of the cercariae for experimental infection of vertebrate hosts to obtain other developmental stages for identification. Experiments were conducted in accordance with the local animal experimentation ethics committee (Comissão de Ética no Uso de Animais, Universidade Federal de Minas Gerais, protocol 20/2016).

We exposed 2 groups of *Poecilia reticulata* guppies (n = 30) individually to 50–100 cercariae of each larval type. We euthanized fish surviving at 30 days postinfection and collected metacercariae found in the gills (Figure, panels B, C) of fish infected with pleurolophocercous cercariae and in the bases of the fins (Figure, panels F and G) of fish exposed to parapleurolophocercous cercariae. We administered metacercariae orally to dexamethasone-immunosuppressed mice. Adult parasites recovered in the small intestines of mice at 6–7 days postinfection were fixed, stained, and mounted on permanent slides. We studied the morphology of the experimentally obtained stages using a light microscope for identification according to taxonomic works (2,9).

We used ethanol-fixed aliquots of cercarial types obtained in *M. tuberculata* snails for molecular characterization. We extracted DNA using the Wizard Genomic DNA Purification Kit (Promega, Madison, WI, USA) and amplified a fragment of the 28S rDNA by PCR using the primers Dig12 (forward) and 1500R (reverse) with PCR conditions as previously described (10). We purified the PCR products with 20% polyethylene glycol 8000 (Promega) and sequenced them in an ABI3730 automated sequencer using Pop-7 Polymer and the ABI BigDye v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA). We edited the sequences we obtained using ChromasPro version 2.0.1 (Technelysium Pty Ltd, South Brisbane, Queensland, Australia), compared them with data available in GenBank, and used them for phylogenetic analyses based on the maximum likelihood method using MEGA7 (<http://www.megasoftware.net/>) and Bayesian inference method using MrBayes 3.1.2 (<http://mrbayes.sourceforge.net/>). We deposited the obtained sequences in GenBank (accession nos. MG738251 and MG738252).

From the experimental infection of mice, we obtained adult parasites identified as *C. formosanus* (Figure, panel D) and *H. pumilio* (Figure, panel H). Molecular data revealed that the samples of *C. formosanus* and *H. pumilio* cercariae found in *M. tuberculata* snails from Peru are conspecific with isolates of these species from Vietnam and Thailand (99.8%–100% similarity), a finding supported by

phylogenetic analyses (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/24/3/17-2056-Techapp1.pdf>). In total, we collected 6,731 *M. tuberculata* snails, of which 112 (1.66%) were found to be infected with heterophyid cercariae. We found *C. formosanus* cercariae in 71 (1.8%) of 3,874 snails collected in VWRCA and in 29 (1.0%) of 2,857 snails collected in PVWR. We found *H. pumilio* cercariae in 12 of 2,857 (0.4%) snails from PVWR.

The presence of *C. formosanus* and *H. pumilio* trematodes in *M. tuberculata* snails from Peru reveals the need to focus attention on the possible effect of these fishborne agents on human health. The increasing number of reports of these trematodes in the Americas indicates that the geographical areas they have invaded are expanding. This finding should serve as a warning, particularly given the popularity of dishes based on raw fish.




Figure. Species of heterophyids transmitted by *Melanoides tuberculata* snails in Peru. A–D) *Centrocestus formosanus*: cercaria (pleurolophocercous type) (A), encysted metacercariae in gills of *Poecilia reticulata* (B, C), and adult parasite obtained in experimentally infected mouse (D). E–H) *Haplorchis pumilio*: cercaria (parapleurolophocercous type) (E), metacercariae found at the base of the caudal fin of *P. reticulata* (F–G), and adult recovered in experimentally infected mouse (H). Scale bars indicate 50 μ m in panels A, C, D, G, and H, 200 μ m in panels B, E, and F.

Acknowledgments

We thank Fernando Gil and Aldo López for the authorization for the development of the research in PVWR and VWRA, respectively. Thanks are due to the biologist Christian Carazas and the engineer José Junco for help in some sampling of snails; to Rosa Martínez for the use of the laboratory at Universidad Nacional Mayor de San Marcos, Peru; and to Airton Lobo for technical support. We are grateful to Sara Vanessa Brant for constructive suggestions in the earlier version of this work.

This project was supported financially by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) (process no. 23038.005297/2011-39) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Brazil (scholarship to E.A.P.M. and L.F.V.F.).

About the Author

Mr. Pulido-Murillo is a PhD student from Peru working in the Universidade Federal de Minas Gerais, Brazil. His research interests focus on morphological and molecular study of zoonotic helminths.

References

- Chai JY, Darwin Murrell K, Lymbery AJ. Fish-borne parasitic zoonoses: status and issues. *Int J Parasitol.* 2005;35:1233–54. <http://dx.doi.org/10.1016/j.ijpara.2005.07.013>
- Pearson J. Family Heterophyidae Leiper, 1909. In: Bray RA, Gibson DI, Jones A, editors. *Keys to the Trematoda*. Vol. III. London: CAB International and Natural History Museum; 2008, p. 113–142.
- Keiser J, Utzinger J. Emerging foodborne trematodiasis. *Emerg Infect Dis.* 2005;11:1507–14. <http://dx.doi.org/10.3201/eid1110.050614>
- Chai JY, Jung BK. Fishborne zoonotic heterophyid infections: an update. *Food Water Parasitol.* 2017 [Epub ahead of print]. <https://doi.org/10.1016/j.fawpar.2017.09.001>
- Arya LK, Rathinam SR, Lalitha P, Kim UR, Ghatani S, Tandon V. Trematode fluke *Procerovum varium* as cause of ocular inflammation in children, South India. *Emerg Infect Dis.* 2016;22:192–200 <https://doi.org/10.3201/eid2202.150051>. <http://dx.doi.org/10.3201/eid2202.150051>
- Pinto HA, de Melo AL. A checklist of trematodes (Platyhelminthes) transmitted by *Melanoides tuberculata* (Mollusca: Thiaridae). *Zootaxa.* 2011;2799:15–28.
- Pointier JP, David P, Jarne P. The biological control of the snail hosts of schistosomes: the role of competitor snails and biological invasions. In: Toledo R, Fried B, editors. *Biomphalaria* snails and larval trematodes. New York: Springer; 2011. p. 215–38.
- Simone LRL. Land and freshwater mollusks of Brazil. São Paulo: Editora Gráfica Bernardi & Fundação de Amparo à Pesquisa do Estado de São Paulo; 2006.
- Scholz T, Aguirre-Macedo ML, Salgado-Maldonado G. Trematodes of the family Heterophyidae (Digenea) in Mexico: a review of species and new host and geographical records. *J Nat Hist.* 2001;35:1733–72. <http://dx.doi.org/10.1080/00222930152667087>
- Tkach VV, Littlewood DTJ, Olson PD, Kinsella JM, Swiderski Z. Molecular phylogenetic analysis of the Microphalloidea Ward, 1901 (Trematoda: Digenea). *Syst Parasitol.* 2003;56:1–15. <http://dx.doi.org/10.1023/A:1025546001611>

Address for correspondence: Hudson A. Pinto, Laboratório de Biologia de Trematoda, Department of Parasitology, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, C.P. 486, 30123-970, Belo Horizonte, Minas Gerais, Brazil; email: hudsonalves13@icb.ufmg.br

LETTER

Ceftriaxone-Resistant *Neisseria gonorrhoeae*, Canada, 2017

Alan R. Katz

Author affiliation: University of Hawaii, Honolulu, Hawaii, USA

DOI: <http://dx.doi.org/10.3201/eid2403.171892>

To the Editor: I read with great interest the report by Lefebvre et al. about a *Neisseria gonorrhoeae* isolate identified in Canada demonstrating a ceftriaxone MIC of 1 mg/L (1). The authors note: “As of October 15, 2017, only 5 ceftriaxone-resistant *Neisseria gonorrhoeae* isolates had been reported worldwide (MIC range 0.5–2 mg/L).” The authors cite published reports from Spain, Japan, Australia, and France.

I would like to clarify that additional *N. gonorrhoeae* isolates have been identified with ceftriaxone MICs ≥ 0.5 mg/L. Since 1987, as part of the Gonococcal Isolate Surveillance Project, the Centers for Disease Control and Prevention has been testing *N. gonorrhoeae* isolates for ceftriaxone susceptibility. During 1987–2016, the Centers for Disease

Control and Prevention identified and reported 5 isolates with ceftriaxone MICs of 0.5 mg/L in the United States. These isolates were found in San Diego, California (1987); Cincinnati, Ohio (1992 and 1993); Philadelphia, Pennsylvania (1997); and most recently, Oklahoma City, Oklahoma (2012) (2). Therefore, although the number of *N. gonorrhoeae* isolates with ceftriaxone MICs ≥ 0.5 mg/L identified globally to date has been small, these Gonococcal Isolate Surveillance Project findings should be acknowledged. Continued and enhanced global surveillance of gonococcal isolates for antimicrobial susceptibility testing is imperative.


References

- Lefebvre B, Martin I, Demczuk W, Deshaies L, Michaud S, Labbe AC, et al. Ceftriaxone-resistant *Neisseria gonorrhoeae*, Canada, 2017 [cited 2017 Nov 19]. Epub ahead of print. <http://dx.doi.org/10.3201/eid2402.171756>
- Centers for Disease Control and Prevention. Sexually transmitted diseases surveillance 2016 [cited 2017 Nov 19]. https://www.cdc.gov/std/stats16/CDC_2016_STDS_Report-for-508WebSep21_2017_1644.pdf

Address for correspondence: Alan R. Katz, University of Hawaii at Manoa, Office of Public Health Studies, Biomedical Sciences Bldg, Rm D104M, 1960 East-West Rd, Honolulu, HI 96822, USA; email: katz@hawaii.edu

Fishborne Zoonotic Trematodes Transmitted by *Melanooides tuberculata* Snails, Peru

Technical Appendix



Technical Appendix Figure. Phylogenetic relationship between *Centrocestus formosanus* and *Haplorchis pumilio* found in Peru (shown in bold) with other heterophyids transmitted by *Melanooides tuberculata*, inferred from partial sequences of 28S rDNA (1176 bp) based on Bayesian inference (BI) and maximum likelihood (ML) analyses. For the ML analysis, nodal support was estimated by the bootstrap method (1000 replications). BI analysis was based on 2 runs, each of 4 chains based on Markov chain Monte Carlo (MCMC) analysis for 1 million generations with sampling interval of 100 generations and a burn-in of 25%. The numbers on the nodes are the posterior probability values and bootstrap for BI and ML, respectively. *Echinostoma revolutum* was used as an external group. The scale bar indicates the number of substitutions per site.