

Outbreak of *Trichinella* T9 Infections Associated with Consumption of Bear Meat, Japan

Katsushige Tada, Hiromichi Suzuki,
Yosuke Sato, Yasuyuki Morishima, Isao Nagano,
Haruhiko Ishioka, Harumi Gomi

An outbreak of trichinellosis occurred in Japan in December 2016. All case-patients had eaten undercooked bear meat, from which *Trichinella* larvae were subsequently isolated. DNA sequencing analysis of the mitochondrial genes cytochrome c-oxidase subunit 1 and internal transcribed spacer 2 confirmed that *Trichinella* T9 had caused the outbreak.

Trichinellosis is a parasitic disease caused by the *Trichinella* spp. nematode that is contracted by eating raw or undercooked meat from infected animals. Approximately 100 species of animals, including humans, can be infected (1). The most common source of human trichinellosis is meat from pigs or wild boar. A total of 65,818 human cases were reported from 41 countries during 1986–2009 (2).

In Japan, trichinellosis is rarely encountered in the clinical setting, and only 5 imported cases (1 in 1998 [3], 1 in 1999 [4], 1 in 2003 [5], 2 in 2009 [6]) have been reported during the past few decades. Three outbreaks of domestically acquired trichinellosis have been reported since 1975 (7), the last reported outbreak occurring in 1981; all were associated with bear meat consumption, but the etiologic agents were not identified at the species level. Since then, no outbreaks were reported until late 2016.

The Study

In December 2016, a previously healthy young man was referred to Tsukuba Medical Center Hospital (Tsukuba, Ibaraki Prefecture, Japan) for a fever, rash, malaise, and eosinophilia. He claimed that he had eaten a bear meat dish (Figure 1) at a restaurant in Mito, Ibaraki Prefecture, Japan, with his 4 friends, who all had similar signs and symptoms. Subsequently, a total of 32 patients who had consumed

the bear meat were reported to the Ibaraki Prefecture Mito Health Center; 28 patients had been evaluated at hospitals. Ethics approval for this research was obtained from the Institutional Review Board of Mito Kyodo General Hospital, University of Tsukuba, Mito, Japan (No. 16-69). All patients provided informed consent for their data to be included in this study.

Among the 28 patients who underwent evaluation, 21 had signs and symptoms that were compatible with trichinellosis. Each serum sample obtained from the 28 patients was tested 3 times for antibodies to *Trichinella* spp., as previously described (6). We performed antibody titer testing with ELISA using excretory–secretory (ES) antigens from *Trichinella spiralis* at the patient's initial presentation and ≥ 2 weeks after the first serum samples were obtained (online Technical Appendix, <https://wwwnc.cdc.gov/EID/article/24/8/17-2117-Techapp1.pdf>). We defined a confirmed case as illness in a patient with a history of consuming raw bear meat, clinical symptoms compatible with trichinellosis, and serologic evidence of trichinellosis. A probable case was defined as illness in a patient with a history of consuming raw bear meat, clinical symptoms compatible with trichinellosis, and a negative serologic test result (8).

In total, 19 (90.4%) patients, all symptomatic, had an antibody titer higher than the cutoff (Table 1); 2 symptomatic patients had an antibody titer lower than the cutoff (titer <200 on convalescent serologic evaluations). All 7 asymptomatic patients had negative serologic test results. Consequently, we identified 21 trichinellosis patients in our study, representing 19 confirmed and 2 probable cases.

We compiled and assessed demographic and clinical data on the 21 patients with confirmed and probable trichinellosis (Table 2). Median age was 35 years (range 23–58 years); 10 (48%) patients were female and 11 (52%) male. Thirteen patients (62%) had consumed ≥ 3 slices of infected bear meat (≈ 10 g per slice). The median incubation period was 19 days (range 6–34 days). All patients had a rash (Figure 2), 20 (95%) had a fever, 17 (81%) had myalgia, 10 (48%) had facial edema, and 9 (43%) had peripheral edema. Only 5 (24%) patients had diarrhea (range of onset day 1–16 of illness) during the outbreak. Nine (43%) patients had conjunctivitis, and 2 (10%) had uveitis.

Author affiliations: Mito Kyodo General Hospital, University of Tsukuba, Mito, Japan (K. Tada, H. Ishioka, H. Gomi); Tsukuba Medical Center Hospital, Tsukuba, Japan (H. Suzuki); Ibaraki Prefecture Mito Health Center, Mito (Y. Sato); National Institute of Infectious Diseases, Tokyo, Japan (Y. Morishima); Gifu University, Gifu, Japan (I. Nagano)

DOI: <https://doi.org/10.3201/eid2408.172117>



Figure 1. The bear meat dish implicated in an outbreak of *Trichinella* T9 infection, Japan, December 2016. Bear meat slices are marked with a circle and an arrow.

At the time of initial evaluation, the median eosinophil count was $1.0 \times 10^9/L$ (range $0.1 \times 10^9/L$ to $4.3 \times 10^9/L$), and the median creatine kinase level was 147 IU/L (range 57–786 IU/L). All patients were treated with albendazole (200 mg or 400 mg, 2×/d for 10–14 days), with or without

prednisolone. In 1 case, albendazole was changed to mebendazole because of a mild increase in the patient’s aspartate aminotransferase and alanine aminotransferase levels, which was later considered to have occurred because of trichinellosis itself. None of the patients had

Table 1. Serologic test results for 28 patients who consumed bear meat associated with *Trichinella* T9 infection, Japan, December 2016*

Patient no.	Signs and symptoms	Highest blood eosinophil count, cells/L	Initial serologic test titer	No. days postinfection	Convalescent-phase serologic test titer	No. days after first blood collection
1	Yes	7.1×10^9	<200	23	6,400	24
2	Yes	4.3×10^9	<200	23	12,800	19
3	Yes	2.7×10^9	<200	27	1,600	20
4	Yes	10.1×10^9	<200	23	800	19
5	Yes	7.8×10^9	800	24	3,200	17
6	Yes	8.8×10^9	<200	25	1,600	17
7	Yes	3.1×10^9	<200	25	1,600	17
8	Yes	1.9×10^9	800	25	6,400	18
9	Yes	11.1×10^9	<200	25	3,200	18
10	Yes	2.4×10^9	<200	24	3,200	15
11	Yes	5.0×10^9	400	12	3,200	16
12	Yes	4.2×10^9	<200	13	800	16
13	Yes	8.5×10^9	200	22	6,400	14
14	Yes	5.3×10^9	<200	20	800	15
15	Yes	10.8×10^9	200	20	3,200	14
16	Yes	2.9×10^9	<200	23	<200	24
17	Yes	4.3×10^9	<200	21	<200	16
18	Yes	1.9×10^9	<200	20	400	13
19	No	0.1×10^9	<200	22	<200	29
20	No	0.2×10^9	<200	18	<200	15
21	No	0.1×10^9	<200	19	<200	28
22	No	0.1×10^9	<200	27	<200	24
23	Yes	2.3×10^9	<200	23	6,400	15
24	Yes	2.9×10^9	<200	23	400	16
25	No	0.4×10^9	<200	25	<200	31
26	No	0.2×10^9	<200	21	<200	14
27	No	0.1×10^9	<200	11	<200	29
28	Yes	2.0×10^9	<200	27	400	13

*ELISA was performed to detect *Trichinella* antigens. The cutoff point (0.148) was 3 times the mean value of A₄₁₄ from the negative serum sample of 100 healthy persons. Of the 28 patients evaluated, 21 had signs and symptoms compatible with trichinellosis. Patients 1–15 also had elevated antibody titers; however, the antibody titers of patients 16 and 17 were not elevated. These cases were defined as probable trichinellosis, as previously described (8).

Table 2. Epidemiologic, clinical, and laboratory data for 21 symptomatic patients with probable or confirmed *Trichinella* T9 infection, Japan, December 2016*

Characteristic	Value
Median age, y (range)	35 (23–58)
Sex	
F	10 (48)
M	11 (52)
Consumed ≥ 3 slices of infected bear meat	13 (62)
Median incubation period, d (range)	19 (6–34)
Signs and symptoms	
Fever	20 (95)
Rash	21 (100)
Myalgia	17 (81)
Fatigue	9 (43)
Facial edema	10 (48)
Peripheral edema	9 (43)
Diarrhea	5 (24)
Conjunctivitis	9 (43)
Uveitis	2 (10)
Median duration from date of eating bear meat to date of blood sampling, d (range)	23 (12–27)
Median leukocyte count, cells/L (range)	7.2×10^9 (3.9×10^9 to 16.9×10^9)
Median eosinophil count, cells/L (range)	1.0×10^9 (0.1×10^9 to 4.3×10^9)
Median aspartate aminotransferase level, IU/L (range)	24 (12–41)
Median alanine aminotransferase level, IU/L (range)	22 (9–73)
Median creatine kinase level, IU/L (range)	147 (57–786)
Median C-reactive protein level, mg/L (range)	9.4 (0.4–67.5)

*Laboratory data were obtained at initial presentation. Values are no. (%) patients except as indicated.



Figure 2. Rash on the back of a patient (patient 10 in Table 1) with confirmed *Trichinella* T9 infection associated with consumption of bear meat, Japan, December 2016. Patient had onset of macular and papular, confluent, and pruritic rash with diffuse blanching on the scalp, face, chest, abdomen, back, and upper and lower extremities. Photo taken 24 days after the patient had consumed the implicated bear meat.

serious complications of trichinellosis or major adverse events during treatment.

The bear meat came from a brown bear (*Ursus arctos*) that had been hunted in Hokkaido Prefecture in November 2016. The meat had been divided into 3 blocks that were preserved in cold storage. Two of these blocks were eaten during this outbreak. The first bear meat block was brought into a restaurant in Mito. In late November, it was seared and served in thin slices with herbs (Figure 1; online Technical Appendix Figure). This bear meat was kept in cold storage and served for 2 days, after which it was preserved in a freezer. The temperature of the cold storage and the freezer were not recorded. Japan Industrial Standard (JIS B 8630) defines the temperature of refrigerated storage as not below 0°C and freezing as $\leq -20^\circ\text{C}$. The bear meat was served after being reheated for a few minutes. The second bear meat block was cooked steak-style to a medium-rare condition; 1 of the patients had eaten meat from this block (online Technical Appendix Figure). The third bear meat block was stored in a freezer without being consumed. We used this meat for the analysis of *Trichinella* spp., which was performed at the National Institute of Infectious Diseases (Tokyo, Japan).

We artificially digested the bear meat with 0.5% pepsin-0.8% HCl solution and then performed a microscopic examination on the sediment. We detected encapsulated larvae with a distinctive esophageal structure (stichosome). The density of the larvae was 84 larvae/g. For the molecular identification of the larvae, we amplified cytochrome *c*-oxidase subunit 1 (*cox1*) and internal transcribed spacer 2 (ITS2) by using PCR with primer pairs described by Kanai

et al. (9). A subsequent sequence analysis showed that both sequences (GenBank accession nos. LS361217 for *cox1* and LS361216 for ITS2) were identical to the corresponding sequences of *Trichinella* T9 (GenBank accession nos. KM357420 for *cox1* and AB255886 for ITS2).

Nine species (*T. spiralis*, *T. britovi*, *T. nativa*, *T. nelsoni*, *T. murrelli*, *T. zimbabwensis*, *T. papuae*, *T. pseudospiralis*, *T. patagoniensis*) and 3 unclassified genotypes (T6, T8, and T9) are currently recognized in the genus *Trichinella* (10). Among them, *T. spiralis* is the most common species in the world (11). The taxonomic status of *Trichinella* species in Japan has not yet been fully elucidated. A recent molecular study revealed that the *Trichinella* isolates obtained from animal specimens in Japan included *Trichinella* T9 (12) and *T. nativa* (13), but *T. spiralis* has not yet been found in Japan (9). *Trichinella* T9 has only been reported in Japan. Therefore, it is considered to be native to Japan. *Trichinella* T9 has been detected and confirmed in a brown bear (13), raccoons (14), raccoon dogs (13,14), and red foxes (13), but no cases of human infection have been reported.

In this outbreak, 2 symptomatic patients and 7 asymptomatic patients had negative serologic test results. A second blood specimen was collected from these patients 33–56 days after the consumption of the bear meat. According to the pertinent literature data (15), serum conversion has been observed up to 65 days postinfection. Thus, we need to consider the possibility of a delay in serum conversion for these 9 patients.

Conclusions

We describe an outbreak of trichinellosis that occurred because of the consumption of bear meat infected with *Trichinella* T9. Public awareness should be raised and education should be promoted to prevent further outbreaks of trichinellosis in Japan.

Acknowledgments

We thank Kenji Miyakawa for making the initial clinical diagnosis of trichinellosis and Tomoe Yukawa for diagnosing and managing the patients with uveitis of trichinellosis.

This work was funded in part by a grant to Y.M. from the Japan Agency for Medical Research and Development (grant no. 17fk0108209j0602).

About the Author

Dr. Tada is an infectious disease fellow at the Center for Global Health, Mito Kyodo General Hospital, University of Tsukuba, Mito, Japan. His primary interest is general infectious diseases and infectious diseases in emergency medicine.

References

- Pozio E. The broad spectrum of *Trichinella* hosts: from cold- to warm-blooded animals. *Vet Parasitol*. 2005;132:3–11. <http://dx.doi.org/10.1016/j.vetpar.2005.05.024>
- Murrell KD, Pozio E. Worldwide occurrence and impact of human trichinellosis, 1986–2009. *Emerg Infect Dis*. 2011;17:2194–202. <http://dx.doi.org/10.3201/eid1712.110896>
- Shiota T, Arizono N, Yoshioka T, Ishikawa Y, Fujitake J, Fujii H, et al. Imported trichinellosis with severe myositis—report of a case [in Japanese]. *Kansenshogaku Zasshi*. 1999;73:76–82. <http://dx.doi.org/10.11150/kansenshogakuzasshi1970.73.76>
- Kusuhara Y, Maeno Y, Nagase K, Taniguchi K, Torikai K, Takahashi Y. A case of mixed infection with *Schistosoma haematobium* and *Trichinella* sp. *Kansenshogaku Zasshi*. 1999;73:614–7. <http://dx.doi.org/10.11150/kansenshogakuzasshi1970.73.614>
- Nakamura T, Miura T, Nakaoka T, Nagano I, Takahashi Y, Iwamoto A. A case of trichinellosis with spontaneous remission [in Japanese]. *Kansenshogaku Zasshi*. 2003;77:839–43. <http://dx.doi.org/10.11150/kansenshogakuzasshi1970.77.839>
- Lo YC, Hung CC, Lai CS, Wu Z, Nagano I, Maeda T, et al. Human trichinosis after consumption of soft-shelled turtles, Taiwan. *Emerg Infect Dis*. 2009;15:2056–8. <http://dx.doi.org/10.3201/eid1512.090619>
- Yamaguchi T. Present status of trichinellosis in Japan. *Southeast Asian J Trop Med Public Health*. 1991;22(Suppl):295–301.
- Schellenberg RS, Tan BJ, Irvine JD, Stockdale DR, Gajadhar AA, Serhir B, et al. An outbreak of trichinellosis due to consumption of bear meat infected with *Trichinella nativa*, in 2 northern Saskatchewan communities. *J Infect Dis*. 2003;188:835–43. <http://dx.doi.org/10.1086/378094>
- Kanai Y, Nonaka N, Katakura K, Oku Y. *Trichinella nativa* and *Trichinella* T9 in the Hokkaido island, Japan. *Parasitol Int*. 2006;55:313–5. <http://dx.doi.org/10.1016/j.parint.2006.08.004>
- Pozio E, Zarlenga DS. New pieces of the *Trichinella* puzzle. *Int J Parasitol*. 2013;43:983–97. <http://dx.doi.org/10.1016/j.ijpara.2013.05.010>
- Pozio E, Darwin Murrell K. Systematics and epidemiology of *Trichinella*. *Adv Parasitol*. 2006;63:367–439. [http://dx.doi.org/10.1016/S0065-308X\(06\)63005-4](http://dx.doi.org/10.1016/S0065-308X(06)63005-4)
- Nagano I, Wu Z, Matsuo A, Pozio E, Takahashi Y. Identification of *Trichinella* isolates by polymerase chain reaction—restriction fragment length polymorphism of the mitochondrial cytochrome *c*-oxidase subunit I gene. *Int J Parasitol*. 1999;29:1113–20. [http://dx.doi.org/10.1016/S0020-7519\(99\)00060-0](http://dx.doi.org/10.1016/S0020-7519(99)00060-0)
- Kanai Y, Inoue T, Mano T, Nonaka N, Katakura K, Oku Y. Epizootiological survey of *Trichinella* spp. infection in carnivores, rodents and insectivores in Hokkaido, Japan. *Jpn J Vet Res*. 2007;54:175–82.
- Kobayashi T, Kanai Y, Ono Y, Matoba Y, Suzuki K, Okamoto M, et al. Epidemiology, histopathology, and muscle distribution of *Trichinella* T9 in feral raccoons (*Procyon lotor*) and wildlife of Japan. *Parasitol Res*. 2007;100:1287–91. <http://dx.doi.org/10.1007/s00436-006-0402-x>
- Pozio E, Varese P, Morales MA, Croppo GP, Pelliccia D, Bruschi F. Comparison of human trichinellosis caused by *Trichinella spiralis* and by *Trichinella britovi*. *Am J Trop Med Hyg*. 1993;48:568–75. <http://dx.doi.org/10.4269/ajtmh.1993.48.568>

Address for correspondence: Harumi Gomi, Center for Global Health, Mito Kyodo General Hospital, University of Tsukuba, 3-2-7, Miyamachi Mito, Ibaraki, 310-0015, Japan; email: hgomi-oky@umin.org

Outbreak of *Trichinella* T9 Infections Associated with Consumption of Bear Meat, Japan

Technical Appendix

Method for Performing ELISA Using Excretory–Secretory Antigens from *Trichinella spiralis*


ES products using as antigen for an ELISA were prepared from *T. spiralis* (ISS413) muscle-stage larvae according to the method of Gómez-Morales et al. (1). In brief, muscle-stage larvae were isolated by pepsin-HCl digestion from mice at 30 days post-infection. The larvae were washed 3 times in phosphate-buffered saline (PBS) and then incubated in Dulbecco's modified Eagle's medium at 37°C for 18 h. The medium was filtered through a 0.2- μ m YM-5 filter and concentrated 100 times in an Amicon pressure concentrating chamber (Amicon, Inc., Billerica, MA, USA).

The ELISA was performed as reported in our previous study (2,3) with slight modifications. In brief, 96-well microtiter plates (MaxiSorp, Nalge Nunc International, Tokyo, Japan) were sensitized with ES antigen at 5 μ g/ml in 0.05 M bicarbonate buffer, pH 9.6 (100 μ l/well) for 3 h at 37°C and overnight at 4°C. After the microplates were washed 3 times with PBS, they were blocked with 150 μ l of PBS containing 2% bovine serum albumin (BSA) for 2 h at 37°C. After washing with PBS-0.05% Tween 20, the microplates were probed with a diluted human serum sample (1:200–1:25,600, 100 μ l/well) in PBS containing 1% BSA for 1 h at 37°C. After washing, 100 μ l of 1:10,000-diluted goat anti-mouse IgG (Fab-specific) peroxidase-conjugate (Sigma Chemical Co., St. Louis, MO, USA) was incubated for 1 h at 37°C. For color development, 2–2'-azono-bis (3-ethylbenzthiazoline-6-sulfonic acid) (Sigma Chemical Co.) was added to each well as a substrate (0.3 mg/ml, 100 μ l/well), and the reaction was terminated after 60 min by adding 50 μ l of 1.25% sodium fluoride per well. The absorbance at 414 nm was monitored with a Multiskan JX plate reader (Labsystems, Helsinki, Finland). Since “3 \times the A_{414} means of 1:200-diluted negative sera from 100 healthy persons” was larger than “the means plus 3 standard deviations,” “3 \times the A_{414} mean of the negative sera” was determined as the cutoff

value in further experiments, resulting in a value of 0.148 for our ELISA test. Each plate contained four positive and four negative reference serum samples. Three-well repeats were measured for each diluted serum sample. The ELISA titers were shown as the highest serum dilution that yielded an A_{414} greater than the cutoff point.

References

1. Gómez-Morales MA, Ludovisi A, Amati M, Cherchi S, Pezzotti P, Pozio E. Validation of an ELISA for the diagnosis of human trichinellosis. *Clin Vaccine Immunol.* 2008;15:1723–9. [PubMed](#)
<http://dx.doi.org/10.1128/CVI.00257-08>
2. Nagano I, Pei F, Wu Z, Wu J, Cui H, Boonmars T, et al. Molecular expression of a cysteine proteinase of *Clonorchis sinensis* and its application to an enzyme-linked immunosorbent assay for immunodiagnosis of clonorchiasis. *Clin Diagn Lab Immunol.* 2004;11:411–6. [PubMed](#)
3. Lo YC, Hung CC, Lai CS, Wu Z, Nagano I, Maeda T, et al. Human trichinosis after consumption of soft-shelled turtles, Taiwan. *Emerg Infect Dis.* 2009;15:2056–8. [PubMed](#)
<http://dx.doi.org/10.3201/eid1512.090619>



Technical Appendix Figure. The relationship between the date on which the infected bear meat was consumed and the occurrence of trichinellosis. A confirmed case was defined as patients with 1) a history of consuming raw bear meat; 2) clinical symptoms compatible with trichinellosis; and 3) serologic evidence of trichinellosis. A probable case was defined as patients with 1) a history of consuming raw bear meat; 2) clinical symptoms compatible with trichinellosis; and 3) negative serologic results. Asterisk indicates that the patient consumed bear meat from a different block of meat from the other case-patients.