

Anncaliia algerae Microsporidiosis Diagnosed by Metagenomic Next-Generation Sequencing, China

Chen Liu,¹ Qin Chen,¹ Ping Fu, Yun-Ying Shi

We report a case of *Anncaliia algerae* microsporidia infection in an immunosuppressed kidney transplant recipient in China. Light microscopy and transmission electron microscopy initially failed to identify *A. algerae*, which eventually was detected by metagenomic next-generation sequencing. Our case highlights the supporting role of metagenomic sequencing in early identification of uncommon pathogens.

Anncaliia algerae is an uncommon, yet emerging microsporidian parasitic pathogen that can affect immunocompromised patients and cause fatal myositis (1,2). We report a case of *A. algerae* microsporidiosis, which was initially missed by conventional light microscopy (LM) and subsequent transmission electron microscopy (TEM) of biopsied muscle but eventually identified by metagenomic next-generation sequencing (mNGS).

The Study

In March 2021, a 45-year-old male kidney transplant recipient in China was admitted to the hospital for a 2-month history of muscle pain. He was receiving prednisone, tacrolimus, and mycophenolate mofetil for maintenance immunosuppression. The patient did not have respiratory symptoms at admission. Physical examination showed low fever and tenderness and generalized weakness in all 4 limbs. Laboratory investigations revealed serum creatine kinase level within reference range but low CD4⁺ T lymphocyte count (45 cells/ μ L; reference range 471–1,220 cells/ μ L). Serum cytomegalovirus DNA was 1.64×10^2 copies/mL. Results of tests for heavy metals, parasites, and myositis-specific autoantibodies were negative.

The patient was febrile (37.3°C) at admission. Although immunosuppressant drugs were tapered dramatically, and broad-spectrum antimicrobial drugs and ganciclovir were added, the patient remained febrile (Figure 1). Chest computed tomography (CT) imaging showed patchy irregular ground-glass opacity in the left upper lung lobe. Electromyography testing showed myogenic damage in the biceps brachii muscle. Magnetic resonance imaging of lower extremities revealed swollen soft tissue. Bronchoalveolar lavage (BAL) testing was negative for bacteria, fungi, and *Pneumocystis jirovecii* DNA.

The patient's myalgia and weakness worsened, his serum creatine kinase level increased (Appendix Figure 1, <https://wwwnc.cdc.gov/EID/article/28/7/21-2315-App1.pdf>), and watery diarrhea developed. Stool microscopy, gastroduodenoscopy, and colonoscopy revealed no specific abnormalities; repeated chest CT scans showed increased inflammatory exudation and bilateral pleural effusion.

No specific findings were reported from the initial LM of the left biceps brachii biopsy specimen except for degradation and necrosis of myofibers. We performed a second fiberoptic bronchoscopy and sent BAL fluid for untargeted mNGS via NextSeq 550 (Illumina, <https://www.illumina.com>), which revealed *P. jirovecii* (142 sequence reads) and *A. algerae* (127 sequence reads) within 48 hours of receiving the specimen (Appendix Table 1; Appendix Figure 2, panel A).

Because the previous biopsy results were negative and we were unfamiliar with *A. algerae* microsporidia, we performed a literature review and then reviewed the initial muscle biopsy again. We considered the possibility of a combined infection of *P. jirovecii* and *A. algerae*, and we consulted an infectious disease specialist who suggested adding oral sulfamethoxazole/trimethoprim (SMZ/TMP; 1,600/

Author affiliations: West China Hospital of Sichuan University, Chengdu, China (C. Liu, Q. Chen, P. Fu, Y.-Y. Shi); Chengdu 363 Hospital of Southwest Medical University, Chengdu (Q. Chen)

DOI: <https://doi.org/10.3201/eid2807.212315>

¹These first authors contributed equally to this article.




Figure 1. Clinical course of a 45-year-old patient with *Anncalia algerae* microsporidiosis infection, China. The upper section of the graph shows the body-temperature curve (black line); dotted black line indicates 37.2 °C, the upper limit of normal body temperature. Thick blue and green lines indicate medications administered; dashed thick blue line indicates a dosing frequency of every other day. Major events during the patient's course are indicated by arrows on the x-axis. Asterisk on day 34 denotes the initial light microscopy, which failed to detect *A. algerae* spores. BALF, bronchoalveolar lavage fluid; CFP/SBT, cefoperazone/sulbactam; CT, computed tomography; FB, fiberoptic bronchoscopy; GCV, ganciclovir; GI, gastrointestinal; HD, hemodialysis; LM, light microscopy; MEM, meropenem; mNGS, metagenomic next-generation sequencing; MRI, magnetic resonance imaging; MXF, moxifloxacin; PIP/SBT, piperacillin/sulbactam; SMZ/TMP, sulfamethoxazole/trimethoprim; TEM, transmission electron microscopy.

320 mg 3×/d), which might be effective against both pathogens. After SMZ/TMP treatment, the patient's temperature returned to normal for 5 successive days before climbing to 37.8°C on day 43 of admission; we added oral albendazole (400 mg 2×/d) (Figure 1), according to published cases (1,3,4).

However, the patient's condition continued to deteriorate. On day 51, he decided on comfort care and died 2 days later (Figure 1). On day 52, one day before the patient died, we discovered multiple oval organisms measuring 2–3 μm in scattered clusters under LM in the muscle biopsy sample (Figure 2, panels A–D). After the patient died, we performed mNGS using muscle tissue from the previous biopsy, which yielded 65,311 sequence reads mapped to *A. algerae* (Appendix Table 2; Appendix Figure 2, panel B). *A. algerae* was confirmed by subsequent PCR testing on muscle tissue, but PCR testing of the remaining BAL specimen yielded no findings because not enough fluid was available in the sample after previous examinations. Eventually, we identified *A. algerae* via TEM in the third sample section (Figure 1; Figure 2, panels E, F). We deposited the *A. algerae*

sequences into the National Center for Biotechnology Information Sequence Read Archive (accession nos. SRR18339014 for the BAL sample, SRR18339013 for the muscle sample).

Conclusions

A. algerae is a microsporidial species that has been reported to cause human infections since 1999 (5). Of 12 reported cases of human *A. algerae* infection (1–11), 11 were among immunocompromised patients (Table). Thus, immunodeficiency, as in this patient, appears to be a critical risk factor for *A. algerae* infection. Although the modes of *A. algerae* transmission to humans remain uncertain, waterborne transmission, either through ingestion of or exposure to spore-contaminated water, has been postulated as the most likely route (2,4,6). This patient lived near ditches in a rural area of the warm and humid Sichuan Basin and was readily exposed to waters possibly contaminated by *A. algerae* spores.

A. algerae infection in humans primarily manifests as myositis (1–11), and in reports we reviewed, 5 (62.5%) of 8 case-patients who had *A. algerae* myositis




Figure 2. Light microscopy and transmission electron microscopy of left biceps brachii muscle biopsy tissue from a 45-year-old man with microsporidiosis caused by *Anncaliia algerae*, China. A–D) Light microscopy using different stains. A) Periodic acid-Schiff stain. Scale bar indicates 10 μ m. Original magnification $\times 50$. B) Gomori methenamine silver stain. Scale bar indicates 10 μ m. Original magnification $\times 63$. C) Warthin-Starry stain. Scale bar indicates 10 μ m. Original magnification $\times 63$. D) Toluidine blue stain. Scale bar indicates 10 μ m. Arrows indicate myocytes replaced by aggregates of 2–3 μ m ovoid organisms. Original magnification $\times 63$. E, F) Transmission electron microscopy showing *Anncaliia*-like microsporidia. Scale bars indicate 500 nm. E) A mature spore with electron-dense exospore, electron-lucent endospore, and a single row of 6 to 8 polar tubule coils (arrows). Original magnification $\times 8,000$. F) Proliferating form of microsporidia showing diplokaryotic nuclei (stars) with vesiculotubular structures extending from the meront cell membrane and aggregating in the host cell cytoplasm (arrows). Original magnification $\times 3,000$.

died (Table). Because of fatality risk, early diagnosis and prompt interventions are crucial. To date, biopsy and microscopy remain the standard approaches in microsporidia identification (12), and the role of mNGS has yet to be confirmed.

Although LM is the fastest diagnostic tool for microsporidiosis, it has several limitations. First, LM is unable to identify the genus and species of microsporidia. Second, the actual turnaround time (5–7 days in our hospital) for LM varies among institutions,

Table. Clinical characteristics of 12 previously reported cases of human *Anncaliia algerae* microsporidia infection*

Case reports	Age, y/sex	Immunocompromised/ underlying conditions	Related symptoms	Positive biopsy sample sites	Treatment	Outcome
Watts et al. 2014 (1)	67/M	Y/RA	Myalgias	Vastus lateralis	Albendazole	Survived
	66/M	Y/RA	Myalgias	Vastus lateralis	NG	Died
Coyle et al. 2004 (2)	57/F	Y/RA	Myalgias	Quadriceps femoris	Albendazole	Died
Boileau et al. 2016 (3)	49/M	Y/CLL	Myalgias	Vastus lateralis	Albendazole and fumagillin	Survived
Sutrave et al. 2018 (4)	66/M	Y/GVHD	Myalgias	Vastus lateralis	Albendazole	Survived
Visvesvara et al. 1999 (5)	67/M	N/N	Eye	Cornea	Albendazole	Survived
			discomfort		and fumagillin	
Ziad et al. 2021 (6)	55/M	Y/psoriatic arthritis	Myalgias	Vastus lateralis, intercostal muscle, and tongue	Albendazole	Died
Visvesvara et al. 2005 (7)	11/M	Y/ALL	Skin lesions	Skin	NA	NA
Calí et al. 2010 (8)	69/M	Y/CLL	Hoarseness	False vocal cord	Albendazole	Died
Field et al. 2012 (9)	49/F	Y/lung transplant	Myalgias	Deltoid and tongue	NG	Died
Chacko et al. 2013 (10)	56/M	Y/kidney transplant	Myalgias	Deltoid	Albendazole	Died
Anderson et al. 2019 (11)	60/M	Y/kidney and pancreas transplant	Skin lesions	Lower extremity, finger, tongue, urine, and sputum	Albendazole	Died

*ALL, acute lymphoblastic leukemia; CLL, chronic lymphocytic leukemia; GVHD, graft-versus-host disease; NA, data not available; NG, treatment for *A. algerae* was not given because the patient was undiagnosed before death; RA, rheumatoid arthritis.

which could cause diagnostic delays. Third, the accuracy of LM diagnosis relies on laboratory conditions and microscopist experience. In addition, morphologic features of *A. algerae* spores overlap with those of other organisms, such as small yeasts, which has led to misdiagnosis under LM (1,11). Thus, familiarity with *A. algerae* spores and their appearance on histopathology preparations are crucial for rapid diagnosis. In this case, *A. algerae* spores initially were missed by the microscopist and were detected 2 weeks later during retrospective review because of the relatively long turnaround time.

TEM remains the standard technique for determining the specific microsporidia genus by identifying the ultrastructural characteristics (12). TEM examines a smaller area of tissue at one time but usually has a longer turnaround time than routine LM. TEM results are available in 1–2 days in some institutions, but turnaround time in our hospital takes ≈10–14 days.

As an unbiased, culture-free method capable of detecting all potential pathogens, untargeted mNGS enables identification of unexpected or unknown organisms (13). Compared with hypothesis-driven methods, such as PCR, shotgun mNGS is hypothesis-free, enables survey of all DNA and RNA in multiple samples en masse (13), and generally takes 24–48 hours to produce results. However, mNGS is unlikely to replace conventional diagnostic testing because of its limitations, such as high cost (US \$522 for DNA detection and \$894 for both DNA and RNA in our hospital), lack of a unified workflow, and no standard methods for interpreting results (13). Instead, mNGS can serve as a valuable adjunct tool in diagnosing uncommon or unexplained infections when conventional methods such as LM fail.

Albendazole and fumagillin have been used to treat *A. algerae* infections in previously reported cases (Table). We have easy access to albendazole, but no access to fumagillin. SMZ/TMP was reported to have no effect against *Enterocytozoon bieneusi* microsporidiosis (14), but data regarding effectiveness against *A. algerae* microsporidia were limited. Treatment was greatly delayed in this patient because of our lack of clinical experience with *A. algerae* microsporidia and the late microscopy findings. Early treatment, along with minimized immunosuppression, might be crucial for the successful management of *A. algerae* infection (1,3,4).

In conclusion, *A. algerae* microsporidia infection requires early diagnosis and prompt intervention. LM alone cannot identify microsporidia genus and species; thus, TEM or genomic sequencing are needed for

correct diagnosis. As a sensitive, culture-independent approach, mNGS could be a promising adjunct tool for the early identification of uncommon pathogens, such as *A. algerae* and other microsporidia.

Acknowledgments

We thank Song Lei for providing professional help in the transmission electron microscopy of *A. algerae* spores in muscle tissue. We also thank Teng Xu and Huan Xu for providing the mNGS methods, interpreting the mNGS results, and performing PCR tests.

This work was supported by the National Natural Science Foundation of China (grant no. 81771714) and 1.3.5 Project for Disciplines of Excellence, West China Hospital, Sichuan University (grant no. ZYJC18004).

About the Author

Mr. Liu is a doctoral student at the Department of Nephrology and Kidney Research Institute, West China Hospital of Sichuan University, Chengdu, China. His research interest is infections in renal transplant patients.




References

1. Watts MR, Chan RCF, Cheong EYL, Brammah S, Clezy KR, Tong C, et al. *Anncaliia algerae* microsporidial myositis. *Emerg Infect Dis*. 2014;20:185–91. <https://doi.org/10.3201/eid2002.131126>
2. Coyle CM, Weiss LM, Rhodes LV III, Cali A, Takvorian PM, Brown DF, et al. Fatal myositis due to the microsporidian *Brachiola algerae*, a mosquito pathogen. *N Engl J Med*. 2004;351:42–7. <https://doi.org/10.1056/NEJMoa032655>
3. Boileau M, Ferreira J, Ahmad I, Lavallée C, Qvarnstrom Y, Dufresne SF. Successful treatment of disseminated *Anncaliia algerae* microsporidial infection with combination fumagillin and albendazole. *Open Forum Infect Dis*. 2016;3:ofw158. <https://doi.org/10.1093/ofid/ofw158>
4. Suttrave G, Maundrell A, Keighley C, Jennings Z, Brammah S, Wang M-X, et al. *Anncaliia algerae* microsporidial myositis, New South Wales, Australia. *Emerg Infect Dis*. 2018;24:1528–31. <https://doi.org/10.3201/eid2408.172002>
5. Visvesvara GS, Belloso M, Moura H, Da Silva AJ, Moura IN, Leitch GJ, et al. Isolation of *Nosema algerae* from the cornea of an immunocompetent patient. *J Eukaryot Microbiol*. 1999;46:10S. PubMed
6. Ziad F, Robertson T, Watts MR, Copeland J, Chiu G, Wang D, et al. Fatal disseminated *Anncaliia algerae* myositis mimicking polymyositis in an immunocompromised patient. *Neuromuscl Disord*. 2021;31:877–80. <https://doi.org/10.1016/j.nmd.2021.06.007>
7. Visvesvara GS, Moura H, Leitch GJ, Schwartz DA, Xiao LX. Public health importance of *Brachiola algerae* (Microsporidia) – an emerging pathogen of humans. *Folia Parasitol (Praha)*. 2005;52:83–94. <https://doi.org/10.14411/fp.2005.011>
8. Cali A, Neafie R, Weiss LM, Ghosh K, Vergara RB, Gupta R, et al. Human vocal cord infection with the microsporidium *Anncaliia algerae*. *J Eukaryot*






- Microbiol. 2010;57:562-7. <https://doi.org/10.1111/j.1550-7408.2010.00510.x>
9. Field AS, Paik JY, Stark D, Qiu MR, Morey A, Plit ML, et al. Myositis due to the microsporidian *Anncaliia* (*Brachiola*) *algerae* in a lung transplant recipient. *Transpl Infect Dis.* 2012;14:169-76. <https://doi.org/10.1111/j.1399-3062.2012.00724.x>
 10. Chacko B, Trevillian P. Microsporidial myositis in a kidney transplant recipient [abstract 82]. In: Program and abstracts of Annual Scientific Meeting Transplantation Society of Australia and New Zealand. Canberra (ACT, Australia): Transplantation Society of Australia and New Zealand; 2013. p. 96.
 11. Anderson NW, Muehlenbachs A, Arif S, Bruminhent J, Deziel PJ, Razonable RR, et al. A fatal case of disseminated microsporidiosis due to *Anncaliia algerae* in a renal and pancreas allograft recipient. *Open Forum Infect Dis.* 2019;6:ofz285. <https://doi.org/10.1093/ofid/ofz285>
 12. Franzen C, Müller A. Microsporidiosis: human diseases and diagnosis. *Microbes Infect.* 2001;3:389-400. [https://doi.org/10.1016/S1286-4579\(01\)01395-8](https://doi.org/10.1016/S1286-4579(01)01395-8)
 13. Chiu CY, Miller SA. Clinical metagenomics. *Nat Rev Genet.* 2019;20:341-55. <https://doi.org/10.1038/s41576-019-0113-7>
 14. Costa SF, Weiss LM. Drug treatment of microsporidiosis. *Drug Resist Updat.* 2000;3:384-99. <https://doi.org/10.1054/drup.2000.0174>

Address for correspondence: Yun-Ying Shi, Department of Nephrology, West China Hospital of Sichuan University, No. 37 Guoxue Alley Wuhou District, Chengdu 610041, China; email: yyshi0130@163.com

Emerging Infectious Diseases Spotlight Topics



**Antimicrobial resistance • Ebola
Etymologia • Food safety • HIV-AIDS
Influenza • Lyme disease • Malaria
MERS • Pneumonia • Rabies • Ticks
Tuberculosis • Coronavirus • Zika**



**EID's spotlight topics highlight the latest articles
and information on emerging infectious disease
topics in our global community**

<https://wwwnc.cdc.gov/eid/page/spotlight-topics>

Anncaliia algerae Microsporidiosis Diagnosed by Metagenomic Next- Generation Sequencing, China

Appendix

Appendix Methods

Literature Review

Regarding *Anncaliia algerae* human infection, we searched English databases including PubMed and Web of Science from inception to March 28, 2022. Keywords used for searching were *Anncaliia algerae*, *Brachiola algerae*, and *Nosema algerae*.

Bronchoalveolar Lavage Fluid (BALF) Detection

We performed Gram staining and acid-fast staining of smears from BALF for bacteria detection and Gram staining for fungi.

Metagenomic Next-Generation Sequencing (mNGS)

Nucleic Acid Extraction, Library Preparation, and Sequencing

DNA was extracted from bronchoalveolar lavage fluid and muscle tissue by using a QIAamp UCP Pathogen DNA Kit (QIAGEN, <https://www.qiagen.com>), following the manufacturer's instructions. Human DNA was removed using Benzonase (Qiagen) and Tween20 (Sigma-Aldrich, <https://www.sigmaaldrich.com>) (1). Total RNA was extracted with a QIAamp Viral RNA Kit (Qiagen) and ribosomal RNA was removed by a Ribo-Zero rRNA Removal Kit (Illumina, <https://www.illumina.com>). cDNA was generated by using reverse transcription PCR and dNTPs (Thermo Fisher Scientific, <https://www.thermofisher.com>). Libraries were

constructed for the DNA and cDNA samples by using a Nextera XT DNA Library Prep Kit (Illumina) (2). Library was quality assessed by Qubit dsDNA High-Sensitivity (HS) Assay Kit (Thermo Fisher Scientific) followed by High Sensitivity DNA Kit (Agilent Technologies, <https://www.agilent.com>) on an Agilent 2100 Bioanalyzer. Library pools were then loaded onto an Illumina Nextseq CN500 sequencer for 75 cycles of single-end sequencing to generate ≈ 20 million reads for each library. For negative controls, we also prepared PBMC samples with 105 cells/mL from healthy donors in parallel with each batch, using the same protocol, and sterile deionized water was extracted alongside the specimens to serve as non-template controls (NTC) (2,3).

Bioinformatics Analyses

Trimmomatic (4) was used to remove low quality reads, adaptor contamination, and duplicate reads, as well as those shorter than 50 bp. Low complexity reads were removed by Kcomplexity with default parameters (5). Human sequence data were identified and excluded by mapping to a human reference genome (hg38) by using Burrows-Wheeler Aligner software (6). We designed a set of criteria similar to the National Center for Biotechnology Information (NCBI) criteria for selecting representative assembly for microorganisms (bacteria, viruses, fungi, protozoa, and other multicellular eukaryotic pathogens) from the NCBI Nucleotide and Genome databases (7). Pathogen lists were selected according to 3 references: 1 Johns Hopkins ABX Guide (https://www.hopkinsguides.com/hopkins/index/Johns_Hopkins_ABX_Guide/Pathogens); Manual of Clinical Microbiology (8); and clinical case reports or research articles published in current peer-reviewed journals (9). The final database consisted of $\approx 13,000$ genomes. Microbial reads were aligned to database with SNAP version 1.0 β .18 (M. Zaharia, unpub. data, <https://arxiv.org/abs/1111.5572>). Virus-positive detection results (DNA or RNA viruses) were defined as the coverage of ≥ 3 non-overlapping regions on the genome. A positive detection was reported for a given species or genus if the reads per million (RPM) ratio, or RPM-r was ≥ 5 ,

where the RPM-r was defined as the RPM sample/RPMNC (i.e., the RPM corresponding to a given species or genus in the clinical sample divided by the RPM in the NC/negative control) (2). In addition, to minimize cross-species misalignments among closely related microorganisms, we penalized (reduced) the RPM of microorganisms sharing a genus or family designation if the species or genus appeared in non-template controls. A penalty of 5% was used for species (10).

PCR Testing

Primers specific to *Anncaliia algerae* (NALGF1-TCA CCA GAG CCT ATG TGC AGG; NALGR2-CTT CAT AAA AAC ATC CAT CTC) were used and amplified from a 405-bp segment of the small ribosomal subunit ribonucleic acid gene, which showed 100% identity with previous Genbank entries (accession nos. HM216911, AM422905).

References

1. Amar Y, Lagkouvardos I, Silva RL, Ishola OA, Foessel BU, Kublik S, et al. Pre-digest of unprotected DNA by Benzonase improves the representation of living skin bacteria and efficiently depletes host DNA. *Microbiome*. 2021;9:123. [PubMed https://doi.org/10.1186/s40168-021-01067-0](https://doi.org/10.1186/s40168-021-01067-0)
2. Miller S, Naccache SN, Samayoa E, Messacar K, Arevalo S, Federman S, et al. Laboratory validation of a clinical metagenomic sequencing assay for pathogen detection in cerebrospinal fluid. *Genome Res*. 2019;29:831–42. [PubMed https://doi.org/10.1101/gr.238170.118](https://doi.org/10.1101/gr.238170.118)
3. Li H, Gao H, Meng H, Wang Q, Li S, Chen H, et al. Detection of pulmonary infectious pathogens from lung biopsy tissues by metagenomic next-generation sequencing. *Front Cell Infect Microbiol*. 2018;8:205. [PubMed https://doi.org/10.3389/fcimb.2018.00205](https://doi.org/10.3389/fcimb.2018.00205)
4. Xu Y, Kang L, Shen Z, Li X, Wu W, Ma W, et al. Dynamics of severe acute respiratory syndrome coronavirus 2 genome variants in the feces during convalescence. *J Genet Genomics*. 2020;47:610–7. [PubMed https://doi.org/10.1016/j.jgg.2020.10.002](https://doi.org/10.1016/j.jgg.2020.10.002)

5. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 2014;30:2114–20. [PubMed https://doi.org/10.1093/bioinformatics/btu170](https://doi.org/10.1093/bioinformatics/btu170)
6. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*. 2009;25:1754–60. [PubMed https://doi.org/10.1093/bioinformatics/btp324](https://doi.org/10.1093/bioinformatics/btp324)
7. National Center for Biotechnology Information. Assembly anomalies and other reasons a genome assembly may be excluded from RefSeq [cited 2021 Mar 15]. <https://www.ncbi.nlm.nih.gov/assembly/help/anomnotrefseq>
8. Carroll KC, Pfaller MA, Landry ML, McAdam AJ, Patel R, Richter SS, et al., editors. *Manual of Clinical Microbiology*, 12th edition. Washington, DC: ASM Press; 2019.
9. Fiorini N, Lipman DJ, Lu Z. Towards PubMed 2.0. *eLife*. 2017;6:e28801. [PubMed https://doi.org/10.7554/eLife.28801](https://doi.org/10.7554/eLife.28801)
10. Gu W, Deng X, Lee M, Sucu YD, Arevalo S, Stryke D, et al. Rapid pathogen detection by metagenomic next-generation sequencing of infected body fluids. *Nat Med*. 2021;27:115–24. [PubMed https://doi.org/10.1038/s41591-020-1105-z](https://doi.org/10.1038/s41591-020-1105-z)


Appendix Table 1. Potentially pathogenic microorganisms detected by metagenomic next-generation sequencing in bronchoalveolar lavage fluid from a patient with *Anncaliia algerae* microsporidiosis, China*

Type	Genus			Species	
	Name	Relative abundance	Sequence no.	Name	Sequence no.
G+	<i>Enterococcus</i>	33.1%	25,931	<i>E. faecium</i>	22,911
G+	<i>Peptostreptococcus</i>	2.1%	1,372	<i>P. anaerobius</i>	1,348
Fungi	<i>Pneumocystis</i>	3.2%	142	<i>P. jirovecii</i>	142
	<i>Anncaliia</i>	2.4%	127	<i>A. algerae</i>	127
	<i>Candida</i>	0.1%	9	<i>C. albicans</i>	6


*G+, gram positive bacteria.

Appendix Table 2. Potentially pathogenic microorganism(s) detected by metagenomic next-generation sequencing in muscle tissue from a patient with *Anncaliia algerae* microsporidiosis, China*

Type	Genus			Species	
	Name	Relative abundance	Sequence no.	Name	Sequence no.
Fungus	<i>Anncaliia</i>	99.9%	65,311	<i>Anncaliia algerae</i>	65,311



Appendix Figure 1. Trends of serum myohemoglobin (reference range 28–72 ng/mL) and creatine kinase (reference range 19–226 IU/L) level from a patient with *Anncaliia algerae* microsporidiosis, China.



Appendix Figure 2. Results of metagenomic next-generation sequencing of samples from a patient with *Anncaliia algerae* microsporidiosis, China. A) BALF sample; B) left biceps branchii muscle biopsy sample. BALF yielded a total of 127 aligned unique DNA reads mapped to *A. algerae* in the reference database, and the coverage of referenced *A. algerae* genome was 0.11%. Muscle biopsy yielded a total of 65,311 sequence reads mapped to *A. algerae* in the reference database, and the coverage of referenced *A. algerae* genome was 51.97%. The x-axis indicates the nucleotide position in the reference genome, ≈ 9.9 Mb. BALF, bronchoalveolar lavage fluid.