

Clostridium butyricum Bacteremia Associated with Probiotic Use, Japan

Ryuichi Minoda Sada, Hiroo Matsuo, Daisuke Motooka,
Satoshi Kutsuna, Shigeto Hamaguchi, Go Yamamoto, Akiko Ueda

Clostridium butyricum, a probiotic commonly prescribed in Asia, most notably as MIYA-BM (Miyarisan Pharmaceutical Co., Ltd.; <https://www.miyarisan.com>), occasionally leads to bacteremia. The prevalence and characteristics of *C. butyricum* bacteremia and its bacteriologic and genetic underpinnings remain unknown. We retrospectively investigated patients admitted to Osaka University Hospital during September 2011–February 2023. Whole-genome sequencing revealed 5 (0.08%) cases of *C. butyricum* bacteremia among 6,576 case-patients who had blood cultures positive for any bacteria. Four patients consumed MIYA-BM, and 1 patient consumed a different *C. butyricum*-containing probiotic. Most patients had compromised immune systems, and common symptoms included fever and abdominal distress. One patient died of nonocclusive mesenteric ischemia. Sequencing results confirmed that all identified *C. butyricum* bacteremia strains were probiotic derivatives. Our findings underscore the risk for bacteremia resulting from probiotic use, especially in hospitalized patients, necessitating judicious prescription practices.

Probiotics have emerged as agents that improve a wide range of conditions and provide essential ingredients for potential health benefits. Probiotics exhibit a diverse array of effects by engaging in competitive interactions with pathogenic microbial communities, competing for binding sites, helping exclude pathogens, and triggering activation of specific genes within and beyond the host's intestinal tract. This process, in turn, stimulates, regulates, and controls the immune response (1). Probiotics have been found to be effective not only in managing conditions

such as acute gastroenteritis (2) and irritable bowel syndrome (3) but also in preventing antibiotic-associated diarrhea (4) and even in alleviating symptoms associated with COVID-19 (5).

Clostridium butyricum is a strictly anaerobic, gram-positive, spore-forming bacillus named for its capacity to produce high amounts of butyric acid. *C. butyricum* was first isolated from the intestines of pigs by Prazmowski in 1880 (6), and several strains of *C. butyricum* have been reported from various environments in humans (7) and animals (8). *C. butyricum* has been detected in the gut of $\approx 20\%$ of human adults (9). Moreover, *C. butyricum* strains were detected in $>30\%$ of environmental samples tested (10). Some strains of *C. butyricum* are currently used as probiotics and have beneficial effects on humans and animals. One strain of *C. butyricum*, known as *C. butyricum* MIYAIRI 588 (CBM 588), can be found in pharmaceutical probiotics, such as MIYA-BM (Miyarisan Pharmaceutical Co., Ltd., <https://www.miyarisan.com>), one of the most commonly prescribed probiotics in Japan. CBM 588 has been described as a unique, nongenetically modified strain that does not naturally produce toxins (11) or cause disease owing to its susceptibility to the KM1 bacteriophage (12). Several confirmatory factors underpin this characterization: it exhibits no propensity for antibiotic resistance transfer, it is devoid of plasmids bearing mobile genetic elements, and it does not possess genes or produce substances related to clostridial toxins, including botulinum neurotoxins A, B, E, and F, or the *Clostridium perfringens* toxins α , β , and ϵ . Genomic scrutiny of CBM 588 revealed no indicators of pathogenic traits or hemolytic capabilities (13). Numerous studies have substantiated the effectiveness of CBM 588, and various animal model experiments have demonstrated its capacity to inhibit the colonization of *Clostridioides difficile* (14) and prevent enterohemorrhagic *Escherichia coli* O157 infection (15). Human studies have confirmed that

Author affiliations: Osaka University Graduate School of Medicine, Osaka, Japan (R.M. Sada, H. Matsuo, S. Kutsuna, S. Hamaguchi, G. Yamamoto); Osaka University Research Institute for Microbial Diseases, Osaka (D. Motooka); Osaka University Hospital, Suita, Osaka (A. Ueda)

DOI: <http://doi.org/10.3201/eid3004.231633>

CBM 588 prevents antibiotic-associated diarrhea (16). In the medical context in Japan, CBM 588 has been prescribed not only for its expected effectiveness as a conventional probiotic but also for the prophylaxis of the diseases we have listed.

There are, however, other strains of *C. butyricum* that are involved in infectious diseases (17–21). A few case reports have noted the development of *C. butyricum* bacteremia in patients taking probiotics, although strain definition tests using whole-genome sequencing were not conducted (22,23). Bacteremia caused by *C. butyricum* is a rare condition, and the prevalence, clinical features, and bacteriologic and genetic origins of the strains are unknown. We conducted a single-center, retrospective study of cases of bacteremia caused by *C. butyricum* in Japan to shed light on this clinical event.

Patients and Methods

Study Design

We conducted a retrospective cohort study at Osaka University Hospital, a 1,086-bed facility in Osaka, Japan. Our study followed the Strengthening the Reporting of Observational Studies in Epidemiology statement for reporting observational studies (24). The Institutional Review Board of Osaka University Hospital approved the study protocol (number 22584(T1)).

Patients and Baseline Characteristics

To identify cases of *C. butyricum* bacteremia, we reviewed all cases of positive blood culture results for any bacteria that occurred during September 19, 2011–February 5, 2023, from the Laboratory for Clinical Investigation database at Osaka University Hospital. We defined *C. butyricum* bacteremia as cases in which *C. butyricum* was detected in ≥ 1 sets of blood cultures. We used MALDI Biotyper (Bruker, <https://www.bruker.com/en>) to identify *C. butyricum* (25). The data we extracted from medical records encompassed such parameters as age; sex; conditions necessitating hospitalization; underlying diseases; placement of a central venous catheter or a peripherally inserted central catheter; presence of polymicrobial bacteremia, including identification of microorganisms other than *C. butyricum*; symptoms at onset; and the updated Charlson Comorbidity Index at the time of bacteremia diagnosis, which was evaluated for every patient (26). In addition, for patients who were prescribed MIYA-BM, we checked the MIYA-BM consumption at the point of diagnosis and confirmed the duration of MIYA-BM prescription. We also

identified whether MIYA-BM was used for specific reasons in these patients. We defined specific reasons for use of MIYA-BM as treatment for diarrhea, concurrent antibiotic use, or medical history of *C. difficile* infection (CDI), ulcerative colitis, hepatic encephalopathy, or a combination of those conditions. Our investigation involved a detailed evaluation of electronic medical records, which included symptoms of diarrhea occurring ≥ 3 times/day, concurrent antibiotic use, and medical history of CDI, ulcerative colitis, or hepatic encephalopathy. Finally, we extracted data on the etiology of bacteremia, antibiotic treatment regimens, and mortality within 90 days.

Microbiologic Information

We determined the MICs for penicillin, ampicillin, cefotaxime, ceftriaxone, cefmetazole, imipenem, meropenem, sulbactam/ampicillin, clavulanic acid/amoxicillin, tazobactam/piperacillin, clindamycin, moxifloxacin, and metronidazole for *C. butyricum* by using the agar dilution method on Brucella agar medium supplemented with 0.5% sheep's blood. Assays to gauge susceptibility followed the guidelines set by the Clinical Laboratory Standards Institute, tailored for anaerobes (28). We assessed the homogeneity of antibiotic susceptibility between the clinical strains and 3 medicinal strains from different lot numbers to evaluate the comparability of their antibiotic susceptibility.

Whole-Genome Sequencing Analysis

We conducted whole-genome analysis of all strains of *C. butyricum* obtained from blood cultures. In addition, we analyzed *C. butyricum* extracted from MIYA-BM tablets. We then investigated the genetic homology between those strains by evaluating the number of single-nucleotide polymorphisms (SNPs) or insertion/deletion genetic variants between clinical strains and the strain from the MIYA-BM tablets. Finally, we conducted a genomic comparison between clinical isolates of *C. butyricum*, the CBM 588 strain, and other strains of the same species. For the comparison, in addition to the reference strain CDC 51208, we selected 7 strains with fully sequenced genomes that are stored in a bioresource repository.

Results

We detected 5 blood culture-positive *C. butyricum* bacteremia cases (0.08%) (Table 1) from a total of 6,576 persons who had blood cultures positive for any bacteria (7,484 total clinical strains, including bacteria other than *C. butyricum*). Bacteremia developed in all 5 patients during hospitalization; 3 patients were

Table 1. Detailed clinical information on 5 patients with bacteremia caused by *Clostridium butyricum* based on a single-institute, retrospective study, Osaka University Hospital, Japan*

Category	Patient no.				
	1	2	3	4	5
Age, y/sex	68/F	81/F	77/M	53/M	19/F
Onset during hospitalization	Yes	Yes	Yes	Yes	Yes
Diseases requiring hospitalization	Chemotherapy	Immunosuppressive treatment	Post-aortic valve replacement	Simultaneous pancreas and kidney transplant	Double lung transplant
Underlying disease	Esophageal cancer; gastric cancer	Dermatomyositis	Aortic valve regurgitation; end-stage kidney disease	End-stage kidney disease; type 1 diabetes	Idiopathic pulmonary arterial hypertension
Immunosuppression	Yes	Yes	No	Yes	Yes
Charlson Comorbidity Index score	2	1	4	6	1
Central venous catheter insertion	Yes	No	Yes	No	Yes
Concurrent MIYA-BM use	Yes	No, but previously administered another probiotic with <i>C. butyricum</i>	Yes	Yes	Yes
Appropriate reason for the prescription of probiotics	Yes (concomitant antibiotic use)	NA	Yes (concomitant antibiotic use)	No	No
Duration of use of probiotics, d	8	NA	12	91	30
Polymicrobial bacteremia, microorganisms other than <i>C. butyricum</i>	Yes (MSSA)	Yes (<i>Enterococcus faecium</i> /MRCNS)	None	None	None
Symptoms of onset	Fever and diarrhea	Fever and diarrhea	Fever and abdominal pain, septic shock	Fever and abdominal pain	Fever and diarrhea
Diagnosis	Enterocolitis	Enterocolitis	NOMI	Duodenal perforation	Enterocolitis
Antibiotics	CMZ	CTR	MEM	MEM	VCM
90-d mortality	Alive	Alive	Died	Alive	Alive

*CMZ, cefmetazole; CTR, ceftriaxone; MEM, meropenem; MRCNS, methicillin-resistant coagulase-negative staphylococci; MSSA, methicillin-resistant *Staphylococcus aureus*; NA, not applicable; NOMI, nonocclusive mesenteric ischemia; VCM, vancomycin.

women and 2 were men. Four patients were immunocompromised: 2 had undergone transplantation, 1 was undergoing chemotherapy for esophageal and gastric cancers, and 1 was receiving multiple immunosuppressive treatments for dermatomyositis. Two of the 5 patients also had end-stage kidney disease and were on dialysis. The Charlson Comorbidity Index scores ranged from 1 to 6 points for each patient. Three patients underwent catheterization with either a central venous catheter or a peripherally inserted central catheter. Four patients were taking prescribed MIYA-BM at the time of bacteremia diagnosis, and 1 patient (no. 2) had been prescribed a different probiotic containing *C. butyricum* 1 month before the diagnosis of bacteremia. All 4 patients taking MIYA-BM were prescribed it >1 week prior to hospitalization, and MIYA-BM was discontinued following the diagnosis of bacteremia in all these patients. Despite a detailed review of the medical records, we were unable to identify the specific reason for prescribing probiotics in 2 patients. All 5 patients had fever and abdominal symptoms, such as diarrhea and pain. One

patient (no. 3) with nonocclusive mesenteric ischemia died within 90 days.

A consistent pattern of antibiotic susceptibility was observed in all clinical strains (Table 2). Moreover, those results were consistent with those of previous reports on the antibiotic susceptibility of *C. butyricum*. *C. butyricum* has been reported to be susceptible to penicillin, ampicillin, cefmetazole, imipenem, meropenem, clavulanic acid/amoxicillin, tazobactam/piperacillin, clindamycin, moxifloxacin, and metronidazole but resistant to cefotaxime and ceftriaxone (11,29,30).

Whole-genome analysis of all 5 patient clinical strains revealed that they either exhibited complete homology or had a maximum divergence of only 19 mutations relative to CBM 588, which was extracted from the MIYA-BM tablets. This result indicates that all clinical strains had the same clone as the CBM 588 extracted from MIYA-BM (Table 3) (31–34). We performed genetic annotation of the detected mutations (Appendix Table, <https://wwwnc.cdc.gov/EID/article/30/4/23-1633-App1.pdf>). We performed phylogenetic analysis of

Table 2. Antimicrobial drug susceptibility of clinical bacterial strains from 5 patients who tested positive for *Clostridium butyricum* in a single-institute, retrospective study, Osaka University Hospital, Japan, and 3 medicinal strains from different lot numbers of *C. butyricum* MIYAIRI 588 strain

Category	Patient strains					Medicinal strains of CBM 588		
	1	2	3	4	5	No. 1	No. 2	No. 3
Patient no.	114–4	129–32	180–11	181–16	216–41			
Strain no.						No. 1	No. 2	No. 3
Antimicrobial drug								
Penicillin	0.25	0.25	0.5	0.5	0.25	0.25	0.25	0.25
Ampicillin	0.25	0.25	0.25	0.25	0.25	0.12	0.25	0.25
Cefotaxime	32	32	32	32	32	32	32	32
Ceftriaxone	8	8	16	8	16	8	16	8
Cefmetazole	≤4	≤4	8	≤4	≤4	≤4	≤4	≤4
Imipenem	1	1	2	1	1	1	1	1
Meropenem	≤0.12	≤0.12	0.5	≤0.12	≤0.12	≤0.12	≤0.12	≤0.12
Sulbactam/ampicillin	≤2	≤2	≤2	≤2	≤2	≤2	≤2	≤2
Clavulanic acid/amoxicillin	0.25	0.25	0.5	0.25	0.25	0.12	0.25	0.12
Tazobactam/piperacillin	≤8	≤8	≤8	≤8	≤8	≤8	≤8	≤8
Clindamycin	0.5	0.25	0.5	0.5	0.25	0.25	0.5	0.25
Moxifloxacin	≤0.5	≤0.5	≤0.5	≤0.5	≤0.5	≤0.5	≤0.5	≤0.5
Metronidazole	≤2	≤2	≤2	≤2	≤2	≤2	≤2	≤2

C. butyricum by using the Type (Strain) Genome Server (35). All clinical isolates and probiotics strain were clustered on the same branches (Figure). Average nucleotide identity scores of clinical isolates against those of the probiotics strain were higher than against those of reference strains. This analysis further corroborated the genetic homology between all the clinical strains and the CBM 588 strain.

Discussion

Our single-center, retrospective study determined that the prevalence of *C. butyricum* bacteremia was 0.08% among all cases with bacteria-positive whole-blood cultures and that all clinical strains were derived from the CBM 588 strain. Bacteremia developed in all patients during hospitalization. Out of 5 cases, 4 had received immunosuppressive treatment and 2 had intra-abdominal issues (1 case of esophageal and gastric cancer and 1 case of post-pancreas and kidney transplantation).

Ishikawa et al. reported a case series of 11 cases of *C. butyricum* bacteremia, including 3 self-experienced cases and 8 cases from a literature review (23). The study revealed that at least 8 cases developed bacteremia during their hospitalization for conditions unrelated to the bacteremia itself. Furthermore, most

patients had intra-abdominal issues at the time of developing bacteremia. In 3 cases, *C. butyricum* bacteremia developed after intra-abdominal surgery. Among the 8 cases without intra-abdominal surgery, 6 cases occurred after various intra-abdominal conditions (2 cases of Crohn's disease, 2 cases of gastrointestinal ulcers, 1 case of biliary tract infection, and 1 case of non-obstructive mesenteric ischemia). Our study results align with previous findings, emphasizing the need for vigilant monitoring of bacteremia development associated with probiotic use in patients with intra-abdominal issues or those undergoing immunosuppressive therapy during their hospitalization.

Our study revealed a high degree of genetic similarity between the strains of *C. butyricum* extracted from MIYA-BM tablets and clinical strains identified through genetic analysis, strongly supporting the definition of probiotic-related bacteremia in all our cases. Reports on probiotic-related bacteremia are scarce. Although systematic reviews of cases of bacteremia after probiotic use have been reported (36), to the best of our knowledge, no studies have evaluated the genetic similarities among these reports. Our study offers evidence supporting a direct causal relationship between probiotic prescription and bacteremia. Nonetheless, the patients we identified as nos. 1 and

Table 3. Results of whole-genome sequencing of *Clostridium butyricum* obtained from blood culture from 5 patients who tested positive for *Clostridium butyricum* in a single-institute, retrospective study, Osaka University Hospital, Japan

Category	Patient strains				
	1	2	3	4	5
Patient no.	114–4	129–32	180–11	181–16	216–41
Strain no.					
Average nucleotide identity* against CBM 588 strains	99.986	99.947	99.949	99.943	99.946
All variants†	50	40	63	65	81
Variants not on rRNA region‡	19	1	2	1	0

*Calculated using FastANI (31).

†Number of all variants in coding genes, which were called and annotated by GATK HaplotypeCaller (32) and snpEff (33) with annotation information from DFAST (34).

‡Number of variants after excluding variants on rRNA region.




Figure. Phylogenetic tree reflecting the relationship between *Clostridium butyricum* MIYAIRI 588, clinical isolates of *C. butyricum*, and 8 reference strains based on data from a single-institute, retrospective study, Osaka University Hospital, Japan. Note: 114-4, 129-32, 180-11, 181-16, and 216-41 represent strain numbers of clinical isolates of *C. butyricum*. MIYAIRI 588 indicates *C. butyricum* MIYAIRI 588. DSM10702 (GCA_014131795.1), NBRC 13949 (GCA_006742065.1), DSM 10702 (GCA_000409755.1), NBRC 13949 (GCA_007992895.1), NBRC 3315 (GCA_007992895.1), ATCC 43755 (GCA_011017415.1), CDC 51208 (GCA_001886875.1), and NBRC 3858 (GCA_007992875.1) represent 8 reference strains. ANI was calculated using FastANI (31). ANI, average nucleotide identity.

2 present lingering challenges. We observed 19 differences in terms of SNPs between the strains found in the blood culture of patient 1 and the CBM 588 strain, which was relatively higher than that of the other patients. However, it is common to evaluate strain dissimilarity using fewer than 100 SNPs. Notably, rapidly growing bacteria, such as *Helicobacter pylori*, can accumulate ≈ 30 SNPs within 6 months of acute infection (37). In fact, some studies have established a genetic similarity cutoff of 80 for carbapenem-resistant *Klebsiella pneumoniae* (38), suggesting that the genetic dissimilarity observed in this case could be reasonably acceptable. We also considered the possibility that long-term oral administration of probiotics in the past could have led to genetic mutations in the CBM 588 strain within the bodies of patients we examined. Patient 2, who had been prescribed a different probiotic containing *C. butyricum* 1 month before the diagnosis of bacteremia, developed bacteremia caused by the CBM 588 strain. We considered 2 possibilities for this observation: the patient had previously taken MIYA-BM and it had colonized in the patient's gastrointestinal tract, leading to an infection; or the *C. butyricum* present in the probiotics the patient was taking had genetic similarities to the CBM 588 strain.

Our findings also bring to light the potential adverse effects related to the inappropriate prescribing of probiotics. In all cases where MIYA-BM was

prescribed, probiotics were administered for >1 week. However, after a comprehensive review of the detailed medical records, we were unable to identify the appropriate reasons for prescribing probiotics in half of the cases. Probiotics exhibit various therapeutic and preventive effects in different medical conditions, such as averting antibiotic-associated diarrhea (39) and CDI (40), preventing hepatic encephalopathy in patients with liver cirrhosis (41), and managing symptoms in patients with ulcerative colitis (42). Although probiotics may demonstrate effectiveness in such specialized clinical scenarios, those scenarios were not observed in the cases we studied, in which probiotics appeared to have been prescribed indiscriminately over an extended period.

One limitation of our study was that it was a single-center, retrospective investigation. Multi-center studies are needed to elucidate the prevalence of *C. butyricum* bacteremia and the genetic origin of the strains. Another limitation was that patient 1 showed improvement with ceftriaxone use, although *C. butyricum* is resistant to it. There is a possibility of contamination resulting from such factors as polymicrobial bacteremia and the absence of central venous catheterization. However, it cannot be ruled out that patients with concurrent sacral pressure ulcers are at risk of developing polymicrobial bacteremia, including *C. butyricum* bacteremia. Also, the duration

of probiotic use for each case patient was based on information documented in their medical records, and the precise prescription durations were not always clear. However, the actual prescription periods must exceed the durations documented in the medical records, because the recorded periods represent at least the minimum assessable timeframe. Moreover, although specific reasons for probiotic prescription were not evident in the medical records, unique justifications may have existed. Nevertheless, it is crucial to note that none of the patients had a history of prior antibiotic use, CDI, irritable bowel syndrome, or liver cirrhosis. Hence, the need for prolonged administration exceeding 2 weeks for therapeutic purposes seems unlikely.

In conclusion, our study demonstrates that all clinical strains of *C. butyricum* identified in the positive blood cultures of the 5 cases we analyzed were derived from the strain found in probiotics. Although this type of bacteremia is rare, careful monitoring is essential when bacteremia is caused by probiotics. Clinicians must avoid long-term, inappropriate prescription of probiotics for hospitalized patients with multiple comorbidities, including immunosuppressive treatment and intraabdominal problems, to prevent bacteremia caused by probiotics.

This research was conducted as part of the All-Osaka U Research in “The Nippon Foundation– Osaka University Infectious Disease Response Project.”

About the Author

Dr Sada is an endowed chair associate professor of the Department of Transformative Protection to Infectious Disease, Graduate School of Medicine, Osaka University, Osaka, Japan. His research interests include the epidemiology of bacteremia, infections caused by rare bacteria, and immunodeficiency-related infections.

References

- Kerry RG, Patra JK, Gouda S, Park Y, Shin HS, Das G. Benefaction of probiotics for human health: A review. *Yao Wu Shi Pin Fen Xi*. 2018;26:927–39. <https://doi.org/10.1016/j.jfda.2018.01.002>
- Collinson S, Deans A, Padua-Zamora A, Gregorio GV, Li C, Dans LF, et al. Probiotics for treating acute infectious diarrhoea. *Cochrane Database Syst Rev*. 2020;12:CD003048.
- Zhang T, Zhang C, Zhang J, Sun F, Duan L. Efficacy of probiotics for irritable bowel syndrome: a systematic review and network meta-analysis. *Front Cell Infect Microbiol*. 2022;12:859967. <https://doi.org/10.3389/fcimb.2022.859967>
- Hempel S, Newberry SJ, Maher AR, Wang Z, Miles JN, Shanman R, et al. Probiotics for the prevention and treatment of antibiotic-associated diarrhea: a systematic review and meta-analysis. *JAMA*. 2012;307:1959–69. <https://doi.org/10.1001/jama.2012.3507>
- Zhu J, Pitre T, Ching C, Zeraatkar D, Gruchy S. Safety and efficacy of probiotic supplements as adjunctive therapies in patients with COVID-19: A systematic review and meta-analysis. *PLoS One*. 2023;18:e0278356. <https://doi.org/10.1371/journal.pone.0278356>
- Kerry RG, Patra JK, Gouda S, Park Y, Shin HS, Das G. Benefaction of probiotics for human health: A review. *J Food Drug Anal*. 2018;26:927–939.
- Mountzouris KC, McCartney AL, Gibson GR. Intestinal microflora of human infants and current trends for its nutritional modulation. *Br J Nutr*. 2002;87:405–20. <https://doi.org/10.1079/BJN2002563>
- Tran NT, Li Z, Ma H, Zhang Y, Zheng H, Gong Y, et al. *Clostridium butyricum*: a promising probiotic confers positive health benefits in aquatic animals. *Rev Aquacult*. 2020;12:2573–89. <https://doi.org/10.1111/raq.12459>
- Finegold SM, Sutter VL, Mathisen GE. Normal indigenous intestinal flora. In: Hentges DJ, editor. *Human Intestinal Microflora in Health and Disease*. New York: Elsevier Inc. 1983. p. 1:3–31. <https://doi.org/10.1016/C2012-0-01555-4>
- Ghoddusi HB, Sherburn R. Preliminary study on the isolation of *Clostridium butyricum* strains from natural sources in the UK and screening the isolates for presence of the type E botulinum toxin gene. *Int J Food Microbiol*. 2010;142:202–6. <https://doi.org/10.1016/j.ijfoodmicro.2010.06.028>
- Isa K, Oka K, Beauchamp N, Sato M, Wada K, Ohtani K, et al. Safety assessment of the *Clostridium butyricum* MIYAIRI 588[®] probiotic strain including evaluation of antimicrobial sensitivity and presence of *Clostridium* toxin genes in vitro and teratogenicity in vivo. *Hum Exp Toxicol*. 2016;35:818–32. <https://doi.org/10.1177/0960327115607372>
- Maeda A, Ishii K, Tanaka M, Mikami Y, Arai T. KM1, a bacteriophage of *Clostridium butyricum*. *Microbiology*. 1986;132:2271–5. <https://doi.org/10.1099/00221287-132-8-2271>
- Oka K, McCartney E, Ariyoshi T, Kudo H, Vilá B, de Jong L, et al. In vivo safety evaluation of the *Clostridium butyricum* MIYAIRI 588 strain in broilers, piglets, and turkeys. *Toxicol Res Appl*. 2019;3. <https://doi.org/10.1177/2397847319826955>
- Hagihara M, Ariyoshi T, Kuroki Y, Eguchi S, Higashi S, Mori T, et al. *Clostridium butyricum* enhances colonization resistance against *Clostridioides difficile* by metabolic and immune modulation. *Sci Rep*. 2021;11:15007. <https://doi.org/10.1038/s41598-021-94572-z>
- Takahashi M, Taguchi H, Yamaguchi H, Osaki T, Sakazaki R, Kamiya S. Antagonistic interaction between *Clostridium butyricum* and enterohemorrhagic *Escherichia coli* O157:H7 [in Japanese]. *Kansenshogaku Zasshi*. 1999;73:7–14. <https://doi.org/10.11150/kansenshogakuzasshi1970.73.7>
- Seki H, Shiohara M, Matsumura T, Miyagawa N, Tanaka M, Komiyama A, et al. Prevention of antibiotic-associated diarrhea in children by *Clostridium butyricum* MIYAIRI. *Pediatr Int*. 2003;45:86–90. <https://doi.org/10.1046/j.1442-200X.2003.01671.x>
- Muldrew KL. Rapidly fatal postlaparoscopic liver infection from the rarely isolated species *Clostridium butyricum*. *Case Rep Infect Dis*. 2020;2020:1839456. <https://doi.org/10.1155/2020/1839456>
- Smith MF, Borriello SP, Clayden GS, Casewell MW. Clinical and bacteriological findings in necrotising enterocolitis: a controlled study. *J Infect*. 1980;2:23–31. [https://doi.org/10.1016/S0163-4453\(80\)91727-2](https://doi.org/10.1016/S0163-4453(80)91727-2)
- Sato Y, Kujirai D, Emoto K, Yagami T, Yamada T, Izumi M, et al. Necrotizing enterocolitis associated with *Clostridium butyricum* in a Japanese man. *Acute Med Surg*. 2018;5:194–8. <https://doi.org/10.1002/ams2.329>

20. Cassir N, Benamar S, La Scola B. *Clostridium butyricum*: from beneficial to a new emerging pathogen. *Clin Microbiol Infect*. 2016;22:37–45. <https://doi.org/10.1016/j.cmi.2015.10.014>
21. Cassir N, Benamar S, Khalil JB, Croce O, Saint-Faust M, Jacquot A, et al. *Clostridium butyricum* strains and dysbiosis linked to necrotizing enterocolitis in preterm neonates. *Clin Infect Dis*. 2015;61:1107–15. <https://doi.org/10.1093/cid/civ468>
22. Shimura M, Mizuma M, Nakagawa K, Aoki S, Miura T, Takadate T, et al. Probiotic-related bacteremia after major hepatectomy for biliary cancer: a report of two cases. *Surg Case Rep*. 2021;7:133. <https://doi.org/10.1186/s40792-021-01216-5>
23. Ishikawa K, Hasegawa R, Shibutani K, Mikami Y, Kawai F, Matsuo T, et al. Probiotic-related *Clostridium butyricum* bacteremia: a case report and literature review. *Anaerobe*. 2023;83:102770. <https://doi.org/10.1016/j.anaerobe.2023.102770>
24. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Ann Intern Med*. 2007;147:573–7. <https://doi.org/10.7326/0003-4819-147-8-200710160-00010>
25. Sulaiman IM, Miranda N, Simpson S. MALDI-TOF mass spectrometry and 16S rRNA gene sequence analysis for the identification of foodborne *Clostridium* spp. *J AOAC Int*. 2021;104:1381–8. <https://doi.org/10.1093/jaoacint/qsab070>
26. Quan H, Li B, Couris CM, Fushimi K, Graham P, Hider P, et al. Updating and validating the Charlson Comorbidity Index and score for risk adjustment in hospital discharge abstracts using data from 6 countries. *Am J Epidemiol*. 2011;173:676–82. <https://doi.org/10.1093/aje/kwq433>
27. Wilkins T, Sequoia J. Probiotics for gastrointestinal conditions: a summary of the evidence. *Am Fam Physician*. 2017;96:170–8.
28. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing, 33rd ed (M100-ED33). Wayne (PA): The Institute; 2023.
29. Mory F, Lozniewski A, Bland S, Sedallian A, Grollier G, Girard-Pipau F, et al. Survey of anaerobic susceptibility patterns: a French multicentre study. *Int J Antimicrob Agents*. 1998;10:229–36. [https://doi.org/10.1016/S0924-8579\(98\)00041-7](https://doi.org/10.1016/S0924-8579(98)00041-7)
30. Hecht DW. Anaerobes: antibiotic resistance, clinical significance, and the role of susceptibility testing. *Anaerobe*. 2006;12:115–21. <https://doi.org/10.1016/j.anaerobe.2005.10.004>
31. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun*. 2018;9:5114. <https://doi.org/10.1038/s41467-018-07641-9>
32. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytisky A, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res*. 2010;20:1297–303. <https://doi.org/10.1101/gr.107524.110>
33. Cingolani P, Platts A, Wang L, Coon M, Nguyen T, Wang L, et al. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly (Austin)*. 2012;6:80–92. <https://doi.org/10.4161/fly.19695>
34. Tanizawa Y, Fujisawa T, Nakamura Y. DFAST: a flexible prokaryotic genome annotation pipeline for faster genome publication. *Bioinformatics*. 2018;34:1037–9. <https://doi.org/10.1093/bioinformatics/btx713>
35. Meier-Kolthoff JP, Göker M. TYGS is an automated high-throughput platform for state-of-the-art genome-based taxonomy. *Nat Commun*. 2019;10:2182. <https://doi.org/10.1038/s41467-019-10210-3>
36. Costa RL, Moreira J, Lorenzo A, Lamas CC. Infectious complications following probiotic ingestion: a potentially underestimated problem? A systematic review of reports and case series. *BMC Complement Altern Med*. 2018;18:329. <https://doi.org/10.1186/s12906-018-2394-3>
37. Linz B, Windsor HM, McGraw JJ, Hansen LM, Gajewski JP, Tomsho LP, et al. A mutation burst during the acute phase of *Helicobacter pylori* infection in humans and rhesus macaques. *Nat Commun*. 2014;5:4165. <https://doi.org/10.1038/ncomms5165>
38. Hassoun-Kheir N, Snitser O, Hussein K, Rabino G, Eluk O, Warman S, et al. Concordance between epidemiological evaluation of probability of transmission and whole genome sequence relatedness among hospitalized patients acquiring *Klebsiella pneumoniae* carbapenemase-producing *Klebsiella pneumoniae*. *Clin Microbiol Infect*. 2021;27:468.e1–7. <https://doi.org/10.1016/j.cmi.2020.04.017>
39. Goldenberg JZ, Lytvyn L, Steurich J, Parkin P, Mahant S, Johnston BC. Probiotics for the prevention of pediatric antibiotic-associated diarrhea. *Cochrane Database Syst Rev*. 2015;12:CD004827. <https://doi.org/10.1002/14651858.CD004827.pub4>
40. Goldenberg JZ, Ma SS, Saxton JD, Martzen MR, Vandvik PO, Thorlund K, et al. Probiotics for the prevention of *Clostridium difficile*-associated diarrhea in adults and children. *Cochrane Database Syst Rev*. 2013;5:CD006095. <https://doi.org/10.1002/14651858.CD006095.pub>
41. Xu J, Ma R, Chen LF, Zhao LJ, Chen K, Zhang RB. Effects of probiotic therapy on hepatic encephalopathy in patients with liver cirrhosis: an updated meta-analysis of six randomized controlled trials. *Hepatobiliary Pancreat Dis Int*. 2014;13:354–60. [https://doi.org/10.1016/S1499-3872\(14\)60280-0](https://doi.org/10.1016/S1499-3872(14)60280-0)
42. Naidoo K, Gordon M, Fagbemi AO, Thomas AG, Akobeng AK. Probiotics for maintenance of remission in ulcerative colitis. *Cochrane Database Syst Rev*. 2011;12:CD007443. <https://doi.org/10.1002/14651858.CD007443.pub2>

Address for correspondence: Ryuichi Minoda Sada, Department of Transformative Protection to Infectious Disease, Graduate School of Medicine, Osaka University, Osaka, Japan; email: sadao@cider.osaka-u.ac.jp

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

Single-Center, Retrospective Study Showing *Clostridium butyricum* Bacteremia Associated with Probiotic Use, Japan

Appendix

Appendix Table. Summary of Genetic Mutation Locations and Types of Variants on Genes

strain	variants name	contig name	position	Reference sequence	Alternative sequence	gene_ID	product	Nucleic acid change	Amino acid change	frequency
129-32	129-32_1	contig2	228589	C	T	MGA_3444	hypothetical protein	c.880C>T	p.Arg294*	0.65
180-11	180-11_1	contig2	391259	A	C	MGA_3588	2-hydroxyacid dehydrogenase	c.200A>C	p.Lys67Thr	0.20
	180-11_2	contig3	7	T	TA	MGA_3995	hypothetical protein	c.757dupT	p.Ter253fs	1.00
181-16	181-16_1	contig2	348564	A	T	MGA_3554	AraC family transcriptional regulator	c.801T>A	p.Phe267Leu	1.00
114-4	114-4_01	contig1	113797	C	T	MGA_97	hypothetical protein	c.3865C>T	p.Pro1289Ser	1.00
	114-4_02	contig1	198552	G	A	MGA_154	transposase	c.1432G>A	p.Asp478Asn	1.00
	114-4_04	contig1	427855	A	G	MGA_323	transposase	c.315T>C	p.Asp105Asp	1.00
	114-4_07	contig1	1642093	A	G	MGA_1474	hypothetical protein	c.332A>G	p.Asn111Ser	1.00
	114-4_08	contig1	1660116	G	A	MGA_1489	tryptophan synthase β chain	c.1162G>A	p.Glu388Lys	1.00
	114-4_10	contig1	1819090	T	C	MGA_1625	transposase	c.690T>C	p.Tyr230Tyr	1.00
	114-4_13	contig1	2203803	G	A	MGA_1967	hypothetical protein	c.5G>A	p.Cys2Tyr	1.00
	114-4_14	contig1	2350242	T	C	MGA_2102	hypothetical protein	c.323A>G	p.Glu108Gly	1.00
	114-4_15	contig1	2438117	C	T	MGA_2183	dihydroorotate dehydrogenase B (NAD(+))%2C electron transfer subunit	c.500G>A	p.Gly167Asp	1.00
	114-4_17	contig1	2756750	C	T	MGA_2447	transcriptional regulator	c.217G>A	p.Asp73Asn	1.00
	114-4_18	contig1	3050309	G	A	MGA_2716	hypothetical protein	c.230C>T	p.Thr77Ile	1.00
	114-4_19	contig1	3076183	A	G	MGA_2738	DNA binding response regulator	c.48T>C	p.Ile16Ile	1.00
	114-4_20	contig1	3238758	C	T	MGA_2878	ATPase AAA	c.1405G>A	p.Val469Ile	1.00
	114-4_21	contig1	3269699	G	A	MGA_2905	hypothetical protein	c.2584C>T	p.Pro862Ser	1.00
	114-4_22	contig1	3438483	A	G	MGA_3066	hypothetical protein	c.315A>G	p.Lys105Lys	1.00
	114-4_27	contig1	3512050	A	G	MGA_3130	isoleucine-tRNA ligase	c.1926T>C	p.Phe642Phe	0.99

strain	variants name	contig name	position	Reference sequence	Alternative sequence	gene_ID	product	Nucleic acid change	Amino acid change	frequency
	114-4_28	contig2	333470	T	C	MGA_3541	PTS maltose transporter subunit IIBC	c.343T>C	p.Ser115Pro	1.00
	114-4_32	contig2	724815	C	T	MGA_3944	oxidoreductase	c.553G>A	p.Glu185Lys	1.00
	114-4_33	contig3	106664	G	A	MGA_4088	membrane metallo endopeptidase	c.1316C>T	p.Ser439Leu	1.00